

National Inventory Report

Iceland 2007

Submitted under the United Nations Framework
Convention on Climate Change



Authors:

Birna Sigrún Hallsdóttir, Environment and Food Agency

Rob Kamsma, Environment and Food Agency

Jón Guðmundsson, Agricultural University of Iceland

Preface

The United Nations Framework Convention on Climate Change (UNFCCC) was adopted in 1992 and entered into force in 1994. According to Articles 4 and 12 of the Convention, Parties are required to develop and to submit annually to the UNFCCC national inventories of anthropogenic emissions by sources and removals by sinks of all greenhouse gases not controlled by the Montreal Protocol.

To comply with this requirement, Iceland has prepared a National Inventory Report (NIR) for the year 2007. The NIR together with the associated Common Reporting Format tables (CRF) is Iceland's contribution to this round of reporting under the Convention, and covers emissions and removals in the period 1990 – 2005.

The NIR is written by the Environment and Food Agency of Iceland (EFA), with a major contribution by the Agricultural University of Iceland (AUI), under the responsibility of the Ministry for the Environment.

Ministry for the Environment, Reykjavík, June 2007

EXECUTIVE SUMMARY	9
1 INTRODUCTION	12
1.1 Background information	12
1.2 Institutional arrangement	14
1.3 Process of inventory preparation	16
1.4 Methodologies and data sources	16
1.5 Key source categories	16
1.6 Quality assurance and quality control (QA/QC).....	17
1.7 Uncertainty evaluation	17
1.8 General assessment of the completeness	18
1.9 Planned and implemented improvements	18
2 TRENDS IN GREENHOUSE GAS EMISSIONS	21
2.1 Emission trends for aggregated greenhouse gas emissions	21
2.2 Emission trends by gas.....	22
2.2.1 Carbon dioxide (CO ₂)	23
2.2.2 Methane (CH ₄).....	26
2.2.3 Nitrous oxide (N ₂ O).....	27
2.2.4 Perfluorcarbons	28
2.2.5 Hydrofluorocarbons (HFCs)	29
2.2.6 Sulphur hexafluorid (SF ₆).....	30
2.3 Emission trends by source	30
2.3.1 Energy	31
2.3.2 Industrial processes.....	33
2.3.3 Solvent and other product use.....	35
2.3.4 Agriculture	36
2.3.5 Waste.....	36
2.3.6 Geothermal energy.....	37
2.4 Emission trends for indirect greenhouse gases and SO ₂	38
2.4.1 Nitrogen oxides (NO _x)	38
2.4.2 Non-methane volatile organic compounds (NMVOC).....	39
2.4.3 Carbon monoxide (CO).....	40
2.4.4 Sulphur dioxide (SO ₂).....	40
3 ENERGY	42
3.1 Overview.....	42
3.1.1 Methodology	42
3.1.2 Completeness	42
3.2 Stationary fuel combustion	43
3.2.1 Energy industries, manufacturing industries, commercial/institutional and residential fuel combustion.....	43
3.3 Mobile combustion	45
3.3.1 Construction sector	45
3.3.2 Road vehicles	45
3.3.3 Fishing.....	46
3.3.4 Civil aviation.....	47
3.3.5 National navigation.....	47

3.4	International bunker fuels	48
3.5	Cross-cutting issues	48
3.5.1	Sectoral versus reference approach.....	48
3.5.2	Feedstock and non-energy use of fuels.....	48
4	INDUSTRIAL PROCESSES.....	49
4.1	Overview.....	49
4.1.1	Methodology.....	49
4.1.2	Completeness.....	49
4.2	Mineral Products.....	50
4.2.1	Cement Production (2A1).....	50
4.2.2	Road paving with asphalt.....	51
4.2.3	Mineral Wool Production	52
4.3	Chemical industry	52
4.4	Metal Production.....	52
4.4.1	Ferroalloys	52
4.4.2	Aluminium Production.....	53
4.5	Emissions from Substitutes for Ozone Depleting Substances – HFCs (2F).....	54
5	SOLVENT AND OTHER PRODUCT USE	57
6	AGRICULTURE	58
6.1	Overview.....	58
6.1.1	Methodology.....	58
6.1.2	Completeness.....	58
6.2	Enteric Fermentation.....	58
6.3	Manure management.....	59
6.4	Emissions from Agricultural Soils – N ₂ O (4D)	61
6.4.1	Description.....	61
6.4.2	Methodological issues.....	61
7	LULUCF	64
7.1	Overview.....	64
7.2	Data Sources	64
7.2.1	NYTJALAND- Icelandic farmland database: Geographical database on condition of farming land.....	65
7.2.2	Vegetation maps.....	66
7.2.3	Cities, towns and villages	67
7.2.4	Unified dataset	68
7.2.5	Land use changes	71
7.2.6	Land use definitions and the classification system and their correspondence to the LULUCF categories.....	71
7.2.7	Uncertainties QA/QC.....	71
7.2.8	Planned improvements regarding land use identification.....	71
7.2.9	Completeness and method	72
7.2.10	Key sources/sink and key areas	75
7.3	Forest land.....	77
7.3.1	Carbon stock changes (5A).....	77
7.3.2	Other emissions (5(I), 5 (II), 5(III)).....	78
7.3.3	Land converted to forest land.	78

7.3.4	Methodological issues.....	79
7.3.5	Emission/removal factors.....	79
7.3.6	Uncertainties QA/QC.....	79
7.3.7	Recalculations.....	80
7.3.8	Planned improvements regarding Forest land.....	80
7.4	Cropland.....	80
7.4.1	Carbon stock changes (5B).....	80
7.4.2	Other emissions (5(I), 5 (II), 5(III), 5(IV)).....	81
7.4.3	Land converted to cropland.....	81
7.4.4	Emission factors.....	81
7.4.5	Uncertainty QA/QC.....	81
7.4.6	Planned improvements regarding cropland.....	82
7.5	Grassland.....	82
7.5.1	Carbon stock changes (5C).....	82
7.5.2	Other emissions (5(IV)).....	83
7.5.3	Emission factors.....	83
7.5.4	Land converted to grassland.....	83
7.5.5	Uncertainty QA/QC.....	84
7.5.6	Planned improvements regarding grassland.....	84
7.6	Wetland.....	85
7.6.1	Carbon stock changes (5D).....	85
7.6.2	Other emissions (5).....	85
7.6.3	Emission factors.....	85
7.6.4	Land converted to wetland.....	85
7.6.5	Uncertainty QA/QC.....	86
7.6.6	Planned improvements regarding Wetland.....	86
7.7	Settlements.....	86
7.7.1	Carbon stock changes (5E).....	86
7.7.2	Other emissions (5).....	86
7.7.3	Land converted to settlement.....	86
7.7.4	Planned improvements regarding Settlement.....	86
7.8	Other land (5, 5F).....	86
7.8.1	Planned improvements regarding other land.....	87
7.9	Other (5).....	87
7.9.1	Harvested Wood Products.....	87
7.9.2	Grassland organic soil.....	87
7.9.3	Revegetation.....	87
7.10	Biomass burning (5V).....	93
7.11	Planned improvements of emission/removal data for LULUCF.....	93
8	WASTE.....	95
8.1	Overview.....	95
8.1.1	Methodology.....	95
8.1.2	Completeness.....	95
8.2	Solid waste disposal sites.....	95
8.2.1	Activity data.....	97
8.2.2	Emission factors.....	98

8.3	Emission from Wastewater Handling (6B).....	101
8.3.1	Methodological issues.....	101
8.4	Waste incineration	103
9	OTHER – GEOTHERMAL ENERGY	105
9.1	Overview.....	105
9.2	Methodology	105
	REFERENCES	106
	ANNEX I: KEY SOURCES.....	109
	ANNEX II: QUANTITATIVE UNCERTAINTY.....	110
	Annex III: CRF TABLES FOR YEAR 2005	111

EXECUTIVE SUMMARY

Kyoto accounting:

For 2005, Iceland's total greenhouse gas emissions (without LULUCF) were estimated to be 3.705 Gg CO₂-equivalents. Iceland's total emissions in 2005 were 10,5% above 1990 levels and 1% above the 1990 levels when activities under Article 3, paragraph 3 and 4 of the Kyoto Protocol are accounted for. Emissions that could fall under Decision 14/CP.7 amounted to 440 Gg in 2005.

Background

The 1992 United Nations Framework Convention on Climate Change (UNFCCC) requires that the Parties report annually on their greenhouse gas emissions by sources and removals by sinks. In response to these requirements, Iceland has prepared the present National Inventory Report (NIR).

The IPCC Good Practice Guidance, the Revised 1996 Guidelines and national estimation methods are used in producing the greenhouse gas emissions inventory. The responsibility of producing the emissions data lies with the Environment and Food Agency, which compiles and maintains the greenhouse gas inventory. Emissions and removals from the LULUCF sector are compiled by the Agricultural University of Iceland. The national inventory and reporting system is continually being developed and improved.

Iceland is a party of the UNFCCC and acceded to the Kyoto Protocol on May 23rd, 2002. Earlier that year the government adopted a climate change policy that was formulated in close cooperation between several ministries. The aim of the policy is to curb emissions of greenhouse gases so they will not exceed the limits of Iceland's obligations under the Kyoto Protocol. A second objective is to increase the level of carbon sequestration resulting from afforestation and revegetation programs. In February 2007 a new climate change strategy was adopted by the Icelandic government. The strategy sets forth a long-term vision for the reduction of net emissions of greenhouse gases by 50-75% until the year 2050, using 1990 emissions figures as a baseline.

The Kyoto Protocol commits Annex I Parties to individual, legally binding targets for their greenhouse gas emissions in the "commitment period" 2008-2012. Iceland's obligations according to the Kyoto Protocol are as follows:

- For the first commitment period, from 2008 to 2012, the greenhouse gas emissions shall not increase more than 10% from the level of emissions in 1990.
- Decision 14/CP.7 on the "Impact of single projects on emissions in the commitment period" allows Iceland to report certain industrial process' carbon dioxide emissions separately and not include them in national totals; to the extent they would cause Iceland to exceed its assigned amount. For the first

commitment period, from 2008 to 2012, the mean annual carbon dioxide emissions falling under decision 14/CP.7 shall not exceed 1,600,000 tons.

Trends in emissions and removals

In 1990, the total emissions of greenhouse gases in Iceland were 3.352 Gg of CO₂-equivalents. In 2005 total emissions were 3.705 Gg CO₂-equivalents. This is an increase of 10,5% over the time period. Total emissions show a decrease between 1990 and 1994, with an exception in 1993, and an increase thereafter.

A summary of the Icelandic national emissions for 1990, 2004 and 2005 is presented in Table ES1 (without LULUCF). Empty cells indicate emissions not occurring.

Table ES1. Emissions of greenhouse gases during 1990, 2004 and 2005 in Gg CO₂-eq.

	1990	2004	2005	Changes 90-05	Changes 04-05
CO ₂	2151	2863	2872	34%	0,3%
CH ₄	413	411	417	1%	1,2%
N ₂ O	363	302	309	-15%	2%
HFC 32		0,1	0,2		98%
HFC 125		22,4	30,1		35%
HFC 134a		11,5	10,9		-6%
HFC 143a		24,3	35,5		46%
HFC 152		0,04	0,05		31%
CF ₄	355	33	22	-94%	-32%
C ₂ F ₆	65	6	4	-94%	-32%
SF ₆	5	5	5	0%	0%
Total Emissions	3352	3678	3705	10,5%	0,7%
CO ₂ emissions 'fulfilling' 14/CP.7		455	440		-3%
Total emissions, excluding CO₂ emissions 'fulfilling' 14/CP.7¹	3352	3223	3265	-3%	1%

The largest contributor of greenhouse gas emissions in Iceland is the energy sector, followed by industrial processes, then agriculture, waste and solvent and other product use. From 1990 to 2005 the contribution of the energy sector to the total emissions increased from 51% to 54%. The contribution of industrial processes decreased from 26% in 1990 to around 17 - 19% in the period 1992 to 1997. The contribution of industrial processes increased again after 1997 and was 25% in 2005.


¹ Decision 14/CP.7 allows Iceland to report certain industrial process carbon dioxide emissions separately and not include them in national totals; to the extent they would cause Iceland to exceed its assigned amount.

Table ES2. Total emissions of greenhouse gases by source 1990, 2004 and 2005, Gg CO₂-eq.

	1990	2004	2005
Energy	1704	1961	1992
Industrial Processes	867	949	944
Emissions fulfilling 14/CP.7*	-	455	440
Solvent Use	6	3	3
Agriculture	568	472	475
Waste	141	167	167
Geothermal Energy	67	124	123
Total without LULUCF	3352	3678	3705
LULUCF	2090	1784	1755

* Industrial process carbon dioxide emissions that could fall under Decision 14/CP.7 are included in national totals

The distribution of the total greenhouse gas emissions over the UNFCCC sectors (including geothermal energy and excluding LULUCF) in 2005 is shown in Figure ES1. Emissions from the energy sector account for 54% of the national total emissions, industrial processes account for 25% and agriculture for 13%. The waste sector accounts for 4%, geothermal energy for 3% and solvent and other product use for 0,1%.


Figure ES1. Emissions of greenhouse gases by UNFCCC sector in 2005

1 INTRODUCTION

1.1 Background information

The 1992 United Nations Framework Convention on Climate Change (UNFCCC) was ratified by Iceland in 1993 and entered into force in 1994. One of the requirements under the Convention is that Parties are to report their national anthropogenic emissions by sources and removals by sinks of all greenhouse gases not controlled by the Montreal Protocol, using methodologies agreed upon by the Conference of the Parties to the Convention (COP).

In 1995 the Government of Iceland adopted an implementation strategy based on the commitments in the Framework Convention. The domestic implementation strategy was revised in 2002, based on the commitments in the Kyoto Protocol and the provisions of the Marrakech Accords. In February 2007 a new climate change policy was adopted by the Icelandic government. Iceland acceded to the Kyoto Protocol on May 23rd 2002. The Kyoto Protocol commits Annex I Parties to individual, legally binding targets for their greenhouse gas emissions in the "commitment period" 2008-2012. Iceland's obligations according to the Kyoto Protocol are as follows:

- For the first commitment period, from 2008 to 2012, the greenhouse gas emissions shall not increase more than 10% from the level of emissions in 1990.
- Decision 14/CP.7 on the "Impact of single project on emissions in the commitment period" allows Iceland to report certain industrial process' carbon dioxide emissions separately and not include them in national totals; to the extent they would cause Iceland to exceed its assigned amount. For the first commitment period, from 2008 to 2012, the mean annual carbon dioxide emissions falling under decision 14/CP.7 shall not exceed 1,600,000 tons.

A new climate change strategy was adopted by the Icelandic government in February 2007². The Ministry for the Environment formulated the strategy in close collaboration with the ministries of Transport and Communications, Fisheries, Finance, Agriculture, Industry and Commerce, Foreign Affairs and the Prime Minister's Office. The long-term vision of the strategy is to reduce net greenhouse gas emissions in Iceland by 50 – 75% by 2050, compared to 1990 levels. In the shorter term, Iceland aims to ensure that emissions of greenhouse gases will not exceed Iceland's obligations under the Kyoto Protocol in the first commitment period. The strategy contains provision for measures in six sectors in curbing and reducing GHG emissions, as well as provisions to increase carbon sequestration resulting from afforestation and revegetation programs.

The greenhouse gas emissions profile for Iceland is unusual in many respects. Firstly, electricity production and space heating are based on renewable energy sources resulting in very low emissions from these sectors. Secondly, more than 80% of emissions from the energy sector stem from mobile sources (transport, mobile machinery and fishing

² http://eng.umhverfissraduneyti.is/media/PDF_skrar/Stefnumorkun_i_loftslagsmalum_en.pdf

vessels). Thirdly, emissions from the LULUCF sector are relatively high. Recent research has indicated that there are significant emissions of carbon dioxide from drained wetlands. These emissions can be attributed to government-sponsored drainage of wetlands in the latter half of the 20th Century, which was largely ceased by 1990. These emissions of CO₂ continue for a long time after drainage. The fourth distinctive feature is that individual sources of industrial process emissions have a significant proportional impact on emissions at the national level. Most noticeable are increased emissions from aluminum production associated with the expanded production capacity of this industry. This last aspect of Iceland's emission profile made it difficult to set meaningful targets for Iceland during the Kyoto Protocol negotiations. This fact was acknowledged in Decision 1/CP.3 paragraph 5(d), which established a process for considering the issue and taking appropriate action. This process was completed with Decision 14/CP.7 on the Impact of single projects on emissions in the commitment period.

The problem associated with the significant proportional impact of single projects on emissions is fundamentally a problem of scale. In small economies, single projects can dominate the changes in emissions from year to year. When the impact of such projects becomes several times larger than the combined effects of available greenhouse gas abatement measures, it becomes very difficult for the party involved to adopt quantified emissions limitations. It does not take a large source to strongly influence the total emissions from Iceland. A single aluminium plant can add more than 15% to the country's total greenhouse gas emissions. A plant of the same size would have negligible effect on emissions in most industrialized countries. Decision 14/CP.7 sets a threshold for significant proportional impact of single projects at 5% of total carbon dioxide emissions of a party in 1990. Projects exceeding this threshold shall be reported separately and carbon dioxide emissions from them not included in national totals to the extent that they would cause the party to exceed its assigned amount. Iceland can therefore not transfer assigned amount units to other Parties through international emissions trading. The total amount that can be reported separately under this decision is set at 1.6 million tons of carbon dioxide. The scope of Decision 14/CP.7 is explicitly limited to small economies, defined as economies emitting less than 0.05% of total Annex I carbon dioxide emissions in 1990. In addition to the criteria above, which relate to the fundamental problem of scale, additional criteria are included that relate to the nature of the project and the emission savings resulting from it. Only projects where renewable energy is used and where this use of renewable energy results in a reduction in greenhouse gas emissions per unit of production will be eligible. The use of best environmental practice and best available technology is also required. It should be underlined that the decision only applies to carbon dioxide emissions from industrial processes. Other emissions, such as energy emissions or process emissions of other gases, such as PFCs, will not be affected.

The Government of Iceland notified the Conference of the Parties with a letter, dated October 17th 2002, of its intention to avail itself of the provisions of Decision 14/CP.7. Since Decision 14/CP.7 only applies for the first commitment period, emissions that are stated in this report as emissions that could fall under Decision 14/CP.7 are not excluded

from national totals. On the other hand, they show the scope Iceland has for increased emissions.

The present report together with the associated Common Reporting Format tables (CRF) is Iceland's contribution to this round of reporting under the Convention, and covers emissions and removals in the period 1990 – 2005. The methodology used in calculating the emissions is according to the revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories as set out by the IPCC Good Practice Guidance and Good Practice Guidance for Land Use, Land-Use Change and Forestry (GPG -LULUCF)(IPCC 2003), to the extent possible.

The greenhouse gases included in the national inventory are the following: carbon dioxide (CO₂), methane (CH₄), nitrous oxide (N₂O), hydrofluorocarbons (HFCs), perfluorocarbons (PFCs) and sulphur hexafluoride (SF₆). Emissions of the precursors NO_x, NMVOC and CO as well as SO₂ are also included, in compliance with the reporting guidelines.

1.2 Institutional arrangement

The Environment and Food Agency of Iceland (EFA), an agency under the Ministry for the Environment, has overall responsibility for the national inventory. EFA compiles and maintains the greenhouse gas emission inventory, except LULUCF which is compiled by the Agricultural University of Iceland (AUI). EFA reports to the Ministry for the Environment, which reports to the Convention. Figure 1.1 illustrates the flow of information and allocation of responsibilities.




Figure 1.1 Information flow and distribution of responsibilities in the Icelandic emission inventory system for reporting to the UNFCCC

A new law on emissions of greenhouse gases was passed by the Icelandic legislature, Althing, in March 2007. The stated purpose of the law is to create conditions for Icelandic authorities to comply with international obligations in limiting emissions of greenhouse gases. The law covers the national system for the estimation of greenhouse gas emissions and removals by sinks, the establishment of a national registry, emission permits and the duty of companies to report relevant information to the authorities.

The law specifies that the EFA is the responsible authority for the national accounting as well as the inventory of emissions and removals of greenhouse gases according to Iceland's international obligations. The EFA shall, in accordance with the legislation, produce instructions on the preparation of data and other information for the national inventory.

1.3 Process of inventory preparation

The EFA collects the bulk of data necessary to run the general emission model, i.e. activity data and emission factors. Activity data is collected from various institutions and companies, as well as by EFA directly. The National Energy Forecast Committee (NEFC) collects annual information on fuel sales from the oil companies. This information has been provided on an informal basis. Since sales statistics were not provided by all the oil companies for the years 2003 – 2005, fuel use by sector has been estimated by the NEFC. The Icelandic Association of Farmers (IAF), on the behalf of the Ministry of Agriculture, is responsible for assessing the size of the animal population each year. On request from the EFA, the IAF also accounts for young animals that are mostly excluded from national statistics on animal population. Statistics Iceland provides information on imports of solvents, the use of fertilizers in agriculture and on the import and export of fuels. The EFA collects various additional data directly. Annually a questionnaire is sent out to the industry regarding imports, use of feedstock, and production and process specific information. Importers of HFCs submit reports on their annual imports by different types of HFCs to the EFA. EFA also estimates activity data with regard to waste. Emission factors are mainly taken from the revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories, since limited information is available from measurements of emissions in Iceland.

AUI receives information on revegetated areas from the Soil Conservation Service of Iceland and information on forests and afforestation from the Icelandic Forest Service. The AUI assesses other land use categories on basis of its own geographical database and available supplementary land use information. AUI then calculates emissions and removals for the LULUCF sector and reports to the EFA.

1.4 Methodologies and data sources

The estimation methods of all greenhouse gases are harmonized with the IPCC Guidelines for National Greenhouse Gas Inventories and are, to the extent possible, in accordance with IPCC's Good Practice Guidance.

The general emission model is based on the equation:

$$\text{Emission (E)} = \text{Activity level (A)} \cdot \text{Emission Factor (EF)}$$

The model includes the greenhouse gases and in addition the precursors and indirect greenhouse gases NO_x, SO₂, NMVOC and CO as well as some other pollutants (POPs).

Methodologies and data sources for LULUCF are described in Chapter 7.

1.5 Key source categories

According to the IPCC definition, a key source category is one that is prioritized within the national inventory system because its estimate has a significant influence on a country's total inventory of direct greenhouse gases in terms of the absolute level of emissions, the trend in emissions, or both. In the Icelandic Emission Inventory key source categories are identified by means of Tier 1 method.

A key source analysis was prepared for this round of reporting. The Table below lists the 16 identified key sources. Tables showing key source analysis (trend and level assessment) can be found in Annex I. The key source analysis shows almost the same result as last year. The only change is that CO₂ emissions from stationary coal combustion are now a key source in trend instead of level.

Table 1.1 Key sources

IPCC SOURCE CATEGORIES	Direct GHG	Key source	
		Level	Trend
ENERGY SECTOR			
Mobile combustion: fishing	CO ₂	v	v
Mobile combustion: road vehicles	CO ₂	v	v
Mobile combustion: road vehicles	N ₂ O		v
Mobile combustion: construction	CO ₂	v	v
CO ₂ emissions from stationary combustion, oil	CO ₂	v	v
CO ₂ emissions from stationary combustion, coal	CO ₂		v
INDUSTRIAL PROCESSES			
CO ₂ emissions from Ferroalloys	CO ₂	v	v
CO ₂ emissions from cement production	CO ₂	v	
CO ₂ emissions from aluminium production	CO ₂	v	v
PFC emissions from aluminium production	PFC		v
Emissions from substitutes for Ozone Depleting Substances	HFC	v	v
AGRICULTURE			
CH ₄ emissions from enteric fermentation	CH ₄	v	v
Direct N ₂ O emissions from agricultural soils	N ₂ O	v	v
Indirect N ₂ O emissions from Nitrogen used in agriculture	N ₂ O	v	v
WASTE			
CH ₄ emissions from solid waste disposal sites	CH ₄	v	v
GEOTHERMAL ENERGY			
CO ₂ emissions from geothermal energy utilisation	CO ₂	v	v

1.6 Quality assurance and quality control (QA/QC)

A QA/QC plan is under preparation. Calculations and units have been checked internally within the EFA and some checks have been made regarding several key sources. Data consistency between years is also checked.

1.7 Uncertainty evaluation

For this round of reporting a preliminary estimate of the quantitative uncertainty of the Icelandic emission inventory has been prepared. The uncertainty estimate has revealed that the total uncertainty of the Icelandic inventory (excluding LULUCF) is 7,5%. The results of the uncertainty estimate can be found in Annex II.

1.8 General assessment of the completeness

An assessment of the completeness of the emission inventory should, according to the IPCC's Good Practice Guidance, address the issues of spatial, temporal and sectoral coverage along with all underlying source categories and activities.

In terms of spatial coverage, the emission reported under the UNFCCC covers all activities within Iceland's jurisdiction.

In the case of temporal coverage, CRF tables are reported for the whole time series from 1990 to 2005.

With regard to sectoral coverage the few sources listed in Table 9 of the CRF are not estimated.

The main sources not estimated are:

- Emissions of CO₂ and CH₄ from distribution of oil products (1B2a v)
- Only the potential emissions of HFCs are estimated and SF₆ emissions are not estimated but held constant over the whole time series (2F)
- The emissions/removals of many LULUCF components are not estimated (see chapter 7.2.9). Most important are probably the emissions from degraded grassland and emissions due to biomass burning.

The reason for not including the above activities/gases in the present submission is lack of data, and/or that additional work was impossible due to time constraints in the preparation of the emission inventory.

1.9 Planned and implemented improvements

In 2004 the UNFCCC secretariat coordinated an in-country review of the 2004 greenhouse gas inventory submission of Iceland, in accordance with decision 19/CP.8 of the Conference of the Parties. The review took place from 20th to 24th of September 2004 in Iceland. The review was based on a complete set of common reporting format tables for the years 1990-2002, as well as a national inventory report from 2004. The expert review team concluded that the Icelandic emissions inventory is largely complete and consistent with the UNFCCC reporting guidelines. However, the expert review team noted some departures from the UNFCCC guidelines in that not all CO₂ emissions/removals from the Industrial Processes and LUCF sectors are included and the lack of data on actual emissions of HFCs and SF₆. There were other gaps in the data, for instance, with regard to emissions from wastewater handling, N₂O and CH₄ emissions from fuel combustion of various combustion sources, CO₂ and N₂O emissions from solvent and other product use, and CO₂ emissions and removals from soils. A centralized review of the 2005 greenhouse gas inventory submissions of Iceland, took place from 10th to 15th of October 2005 in Bonn. The main findings of that review were the same as from the in-country review from the year before.

The expert review teams pointed out the following issues of high importance for improving the inventory:

- The establishment of a more robust institutional and legal framework for fulfilling the reporting requirements under the UNFCCC, for example, as a basis for the preparation of the national energy balance;
- The reporting of all LULUCF activities and industrial process emissions in accordance with the UNFCCC reporting guidelines;
- Closing other estimation and reporting gaps in the inventory (e.g. emissions from wastewater handling, and actual emissions from HFCs and SF₆);
- Improving the quality of the activity data (e.g. in the LULUCF and Waste sectors);
- Further implementation of the IPCC good practice guidance (e.g. the use of additional country-specific methodologies for key sources such as road transportation; quantitative estimation of uncertainties for total and sectoral emissions, as well as for the main key sources; and the introduction of a more advanced quality assurance/quality control system, including verification of information provided by industry, including the fishing industry);
- Improvement of the transparency of the inventory by providing more detailed information in the national inventory report (e.g. on recalculations, on the choice of methodologies, on details of country-specific methodologies and on activity data, and references to background material);
- Improvement of consistency with other national and international data sets (e.g. waste-related activity data, cement production data) and of explanation for inconsistencies, if any.

Based on the review reports, some important improvements have already been implemented, while others are to be implemented. A list of planned and implemented improvements can be seen in Table 1.2.

Table 1.2 Planned and implemented improvements
Implemented improvements:

- The reporting of Industrial Processes falling under Decision 14/CP.7 are now as suggested by the ERT.
- N₂O and CH₄ emissions from fuel combustion of various combustion sources have been estimated.
- N₂O emissions from solvent and other product use have been estimated.
- The Ministry for the Environment, in close cooperation with other relevant ministries has established an institutional and legal framework to further strengthen Icelandic climate change policy.
- Preliminary estimate of the quantitative uncertainty.
- A quality assurance/quality control system is under development.
- Activity data and methodologies in the waste sector have been improved. Emissions from wastewater handling are now included and emissions from solid waste disposal sites are now estimated with the FOD method.
- Afforestation and revegetation prior to 1990 is now reported, as well as liming. All liming is reported under cropland.
- Cultivation of histosols is not reported separately but included in emissions from other drained organic soils.
- Prior to its 2006 submission Iceland did not submit the new CRF LULUCF tables. The LULUCF reporting is thus much more extensive than before. This includes coverage of categorized land use and land use change information on all areas of the country. In many cases, though, information on land use prior to 2004 is not available and thus reported land use for years before 2004 are the same as 2004.

Planned and ongoing improvements:

- Iceland has until now not prepared a national energy balance. Following the recommendations from the In-country review team, Iceland will start preparing annually a national energy balance.
- Estimate actual emissions of HFCs and SF₆.
- Construction of geographically identifiable land use database covering the whole country will start in 2007, including extensive sampling on present and previous land use.
- Ongoing new forest inventory will improve both estimates of Forest land area and Carbon stock changes. Similar effort regarding Revegetation is being prepared.

Improvements under consideration:

- Improve methodologies to estimate emissions from road transportation.
- Develop country-specific emission factor for enteric fermentation
- Revise country-specific N excretion factors.
- Improvements of QA/QC for LULUCF.
- Revision of LULUCF emission/removal factors, emphasizing key sources and aiming at higher tier levels.

2 TRENDS IN GREENHOUSE GAS EMISSIONS

2.1 Emission trends for aggregated greenhouse gas emissions

The total amount of greenhouse gases emitted in Iceland during the period 1990 – 2005 is presented in the following tables, expressed in terms of contribution by gases and by sources.

Table 2.1 below presents emission figures for all direct greenhouse gases, expressed in CO₂-equivalents along with the percentage change indicated for both the time period 1990 – 2005 and 2004 – 2005.

Table 2.1. Emissions of greenhouse gases in Iceland during the period 1990 – 2005 (without LULUCF). Empty cells indicate emissions not occurring. Units: Gg CO₂-eq

Gas	Year															
	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
CO ₂	2151	2072	2200	2304	2266	2299	2388	2483	2493	2696	2745	2747	2842	2765	2863	2872
CH ₄	413	411	407	408	413	407	415	421	428	429	423	429	417	416	411	416
N ₂ O	363	353	332	341	346	342	360	358	356	376	352	345	311	302	302	309
HFC 32									0	0	0	0	0	0	0	0
HFC 125						11	12	11	27	23	15	23	16	26	22	30
HFC 134a			1	2	3	4	6	7	8	8	6	7	4	13	12	11
HFC 143a						10	10	19	29	28	12	24	16	29	24	35
HFC 152a						0	0	0	0	0	0	0	0	0	0	0
CF ₄	355	295	131	63	38	50	21	70	152	147	108	78	61	51	33	22
C ₂ F ₆	65	54	24	12	7	9	4	13	28	27	20	14	11	9	6	4
SF ₆	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5
Total	3352	3189	3099	3135	3078	3138	3222	3387	3526	3739	3684	3671	3684	3618	3678	3705
CO ₂ emissions that could fall under 14/CP.7									108	115	273	404	441	451	455	440
Total, without CO₂ emissions that could fall under 14/CP.7*									3418	3624	3411	3267	3242	3167	3223	3265

	1990	2004	2005	Changes 90-05	Changes 04-05
CO ₂	2151	2863	2872	34%	0,3%
CH ₄	413	411	417	1%	1,2%
N ₂ O	363	302	309	-15%	2%
HFC 32		0,1	0,2		98%
HFC 125		22,4	30,1		35%
HFC 134a		11,5	10,9		-6%
HFC 143a		24,3	35,5		46%
HFC 152		0,04	0,05		31%
CF ₄	355	33	22	-94%	-32%
C ₂ F ₆	65	6	4	-94%	-32%
SF ₆	5	5	5	0%	0%
Total Emissions	3352	3678	3705	10,5%	0,7%
CO ₂ emissions 'fulfilling' 14/CP.7		455	440		-3%
Total emissions, excluding CO₂ emissions 'fulfilling' 14/CP.7*	3352	3223	3265	-3%	1%

*Decision 14/CP.7 allows Iceland to report certain industrial process carbon dioxide emissions separately

As mentioned in Chapter 1.1 industrial process CO₂ emissions that fulfill Decision 14/CP.7 shall be reported separately and shall not be included in national totals, to the extent they would cause a Party to exceed its assigned amount. In this report, emissions that are stated as emissions that could fall under Decision 14/CP.7 show the scope for increased emissions. Since this decision only applies for the first commitment period, they are not excluded from national totals.

In 1990, the total emissions of greenhouse gases (excluding LULUCF) in Iceland were 3.352 Gg of CO₂- equivalents. In 2005 total emissions were 3.705 Gg CO₂-equivalents. This implies an increase of 10,5% over the time period. Total emissions show a decrease between 1990 and 1994, with an exception in 1993, and an increase thereafter. So far, 1999 has been the year with the highest emissions recorded.

Iceland has experienced economic growth since 1990, which explains the general growth in emissions. This has resulted in higher emissions from most sources, but in particular from transport and industrial processes.

Since 1990 the number of private cars has been increasing much faster than the population. Also the number of passengers using the public transport system has declined. More traffic is thus not mainly due to population growth, but much rather because a larger share of the population owns and uses private cars for their daily travel.

During the late nineties large-scale industry expanded in Iceland. The existing aluminium plant and the ferroalloys industry experienced enlargement in 1997 and 1999, and in 1998 a new aluminium plant was established.

Nitrous oxide emissions have decreased since 1990, despite the fact that nitrous oxide emissions from road transport have increased. This is due to a decrease in animal livestock and because fertilizer production in Iceland was terminated in 2001.

Before 1992 there were no imports of HFCs, but since then, imports have increased rapidly in response to the phase-out of CFCs and HCFCs. The potential emissions of HFCs have risen from 0,5 Gg CO₂-equivalent in 1992 to 76,7g CO₂-equivalent in 2005.

The overall increasing trend of greenhouse gas emissions has to some extent been counteracted by decreased emissions of PFCs, caused by improved technology and process control in the aluminium industry.

2.2 Emission trends by gas

As shown in Figure 2.1, the largest contributor by far to the total GHG emissions is CO₂ (78), followed by CH₄ (11%) and N₂O (8) and then by the fluorinated gases PFCs, HFCs and SF₆ (3%). In 1990 the share of CO₂ was lower than in 2004 or (64%), the share of CH₄ and N₂O about the same (12% and 11% respectively) but the share of fluorinated gases was higher (13%).




Figure 2.1 Distribution of emissions of greenhouse gases by gas in 2005

Figure 2.2 illustrates the percentage change in emissions of greenhouse gases by gas in Iceland from 1990 to 2005 compared with 1990.




Figure 2.2 Percentage changes in emissions of greenhouse gases by gas 1990 – 2005 compared with 1990

2.2.1 Carbon dioxide (CO₂)

Fisheries, road transport and industrial processes are the three main sources of CO₂ emissions in Iceland. Since emissions from the electricity generation and space heating are very low because they are generated from renewable energy sources, emissions from stationary combustion are dominated by industrial sources. Thereof the fishmeal industry is by far the largest user of fossil fuels. Emissions from mobile sources in the construction industry are also significant. Emissions from geothermal energy

exploitation are moderate. ‘Other sources’ consist mainly of emissions from coal combustion in the cement industry and other transportation than road transportation. Table 2.2 lists CO₂ emissions from each source category for the period 1990 – 2005. Figure 2.3 illustrates the distribution of CO₂ emissions by main source categories, and Figure 2.4 shows the percentage change in emissions of CO₂ by source from 1990 to 2005 compared with 1990.

Table 2.2 Emissions of CO₂ by sector 1990 – 200, Gg.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Fishing	655	676	740	770	760	772	828	810	781	765	720	640	705	662	712	703
Road vehicles	509	527	540	537	544	534	514	545	552	577	589	595	604	628	636	673
Stationary combustion, oil	237	168	241	253	229	219	270	285	261	264	203	245	275	228	191	168
Industrial processes	393	359	362	410	411	427	426	485	512	659	766	804	822	824	846	835
Construction	121	115	107	116	118	149	144	174	175	192	197	192	180	197	243	254
Geothermal energy	67	67	67	67	67	82	82	71	94	123	163	154	159	138	124	123
Other	169	160	142	152	138	117	124	114	117	115	107	116	97	87	111	115
Total	2151	2072	2200	2304	2266	2299	2388	2483	2493	2696	2745	2747	2842	2765	2863	2872

In 2005 the total CO₂ emissions in Iceland were 2.872 Gg. This implies an increase of about 0,3% from the preceding year but an increase of about 34% from 1990. Emissions from stationary oil combustion decreased by 12% from 2004 to 2005. This is mainly due to decreased emissions from the fish meal industry. Emissions from construction rose by 5% and emissions from road vehicles rose by 6%. Emissions from other sources were around the same in 2005 as in 2004.

The increase in CO₂ emissions between 1990 and 2004 can be explained by increased emissions from industrial processes (113%), road transport (32%), geothermal energy utilization (85%), the construction sector (111%), and from fishing (7%). Since 1990 and in particular after 1995 Iceland has experienced economic growth, which partly explains the general growth in emissions. During the late nineties energy intensive industry expanded. The existing aluminium plant and the ferroalloys industry experienced enlargement in 1997 and 1999, and in 1998 a new aluminium plant was built. The economic growth and the expansion of energy intensive industry have resulted in higher emissions from most sources, but in particular from the industrial processes sector as well as from the construction sector. Emissions from the construction sector have risen, particularly in recent years, due to increased activity related to the construction of Iceland’s largest hydropower plant. Since 1990 the vehicle fleet in Iceland has increased by 50%. This has led to increased emissions from road transportation, a trend that is still ongoing. Furthermore the latest trend has been towards larger passenger cars which consume more fuel. Since 1999 the average fuel consumption of newly registered passenger cars has increased by over 6%. Emissions from fishing rose from 1990 to 1996 because a substantial portion of the fishing fleet was operating in distant fishing grounds. From 1996 the emissions decreased again reaching 1990 levels in 2001. Emissions increased again by 10% between 2001 and 2002, but in 2003 they dropped to 1990 levels.

In 2005 emissions were 7% over 1990 levels. Annual changes in emissions reflect the inherent nature of the fishing industry. Emissions from other sources decreased from 1990 to 2003, but rose again in 2004 and 2005 though they were still 32% below the 1990 level. This is mainly due to changes in the cement industry where production had been slowly decreasing since 1990. Due to the construction of a hydropower plant (resulting in more demand of cement) production increased again in 2004 and 2005, though the major part of the cement used in this project is imported. Emissions from both domestic flight and navigation have declined significantly since 1990.




Figure 2.3 Distribution of CO₂ emissions by source in 2005




Figure 2.4 Percentage changes in emissions of CO₂ by major sources 1990 – 2005, compared with 1990

2.2.2 Methane (CH₄)

As can be seen from Table 2.3 and Figure 2.5, about 61% and 37% of the emissions of methane in 2005 originated from waste treatment and agriculture respectively. The emissions from agriculture have decreased by 14% since 1990, but the emissions from waste treatment increased from 1990 to 2001. This is due to an increased amount of waste generated and increased ratio of landfilled wastes in managed waste disposal sites. The emissions from landfills have decreased slightly since 2001, due to increased methane recovery. In 2005, emissions from waste treatment were 39% above the 1990 level.

In whole emissions of methane have been relatively stable over the period.

Table 2.3 Emissions of CH₄ by sector 1990 – 2004, Gg

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Agriculture	14,0	13,7	13,3	13,2	13,3	12,8	13,0	13,1	13,3	13,2	12,6	12,6	12,3	12,1	11,9	12,0
Waste	5,5	5,7	5,8	6,0	6,1	6,3	6,5	6,7	6,9	7,1	7,3	7,7	7,4	7,6	7,5	7,6
Other	0,3	0,3	0,3	0,3	0,3	0,2	0,3	0,2	0,2	0,2	0,2	0,2	0,2	0,2	0,2	0,2
Total	19,7	19,6	19,4	19,4	19,7	19,4	19,8	20,1	20,4	20,4	20,1	20,4	19,9	19,8	19,6	19,8




Figure 2.5 Distribution of CH₄ emissions by source in 2005




Figure 2.6 Percentage changes in emissions of CH₄ by major sources 1990 – 2005, compared to 1990

2.2.3 Nitrous oxide (N₂O)

As can be seen from Table 2.4 and Figure 2.7 agriculture accounts for around 72% of N₂O emissions in Iceland, with agricultural soils as the most prominent contributor. The second most important source is road transport, which has increased rapidly after the use of catalytic converters in all new vehicles became obligatory in 1995.

The overall nitrous oxide emissions decreased by 15% from 1990 to 2005, due to a decrease in the number of animal livestock and because fertilizer production in Iceland was terminated in 2001.

Table 2.4 Emissions of N₂O by sector 1990 – 2005, Gg

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Agriculture	0,9	0,9	0,8	0,8	0,8	0,8	0,8	0,8	0,8	0,9	0,8	0,8	0,8	0,7	0,7	0,7
Road traffic	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,1	0,1	0,1	0,1	0,1	0,1	0,1	0,1	0,1
Other	0,3	0,3	0,2	0,3	0,3	0,3	0,3	0,3	0,3	0,3	0,2	0,2	0,1	0,1	0,2	0,2
Total	1,2	1,1	1,1	1,1	1,1	1,1	1,2	1,2	1,1	1,2	1,1	1,1	1,0	1,0	1,0	1,0




Figure 2.7 Distribution of N₂O emissions by source in 2005




Figure 2.8 Changes in N₂O emission for major sources between 1990 and 2005

2.2.4 Perfluorocarbons


The emissions of the perfluorocarbons, tetrafluoromethane (CF₄) and hexafluoroethane (C₂F₆) from the aluminium industry were 22,1 and 4,0 Gg CO₂-equivalents respectively in 2005.

Total PFC emissions decreased by 94% in the period of 1990 – 2005. As can be seen from Figure 2.9 the emissions decreased steadily from 1990 to 1996 with the exception of 1995. In 1997 and 1998 the emissions rose again due to expansion of the single existing aluminium plant in 1997 and the establishment of a new aluminium plant in 1998. Since 1998 the emissions show a steady downward trend. PFCs reduction is achieved through

improved technology and process control, which has led to a 98% decrease in the amount of PFCs emitted per tonne of aluminium produced during the period of 1990 – 2005.

Table 2.5 Emissions of PFCs by species 1990 – 2005, Gg CO₂-equivalent

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
CF ₄	355	295	131	63	38	50	21	70	152	147	108	78	61	51	33	22
C ₂ F ₆	65	54	24	12	7	9	4	13	28	27	20	14	11	9	6	4
Total	423	348	155	75	45	59	25	82	180	173	127	92	73	60	39	26


Figure 2.9 Emissions of PFCs from 1990 to 2005, Gg CO₂-equivalent

2.2.5 Hydrofluorocarbons (HFCs)

The total potential emissions of HFCs, used as substitutes for ozone depleting substances, amounted to 76,7 CO₂-equivalents in 2005. The import of HFCs started in 1992 and increased until 1998. Since then annual imports have ranged between 30 and 77 Gg CO₂-equivalents. Sufficient data is not yet available to calculate actual emissions. This means that only potential emissions, based on imports, are estimated. The potential method is likely to overstate emissions, since chemicals used e.g. in refrigerators are emitted over a period of several years. The application category refrigeration contributes by far the largest part of HFCs emissions but foam blowing is also thought to be a minor source.

Table 2.6 Emissions of HFCs by species 1990 – 2005, Gg CO₂-equivalent

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
HFC 32	-	-	-	-	-	-	-	-	0,0	0,0	0,1	0,0	0,0	0,1	0,1	0,2
HFC 125	-	-	-	-	-	10,8	11,7	11,1	27,1	23,5	14,5	23,2	15,7	26,3	22,4	30,1
HFC 134a	-	-	0,5	1,6	3,1	4,1	6,5	7,1	8,0	8,2	6,0	6,8	3,8	13,4	11,5	10,9
HFC 143a	-	-	-	-	-	10,0	10,3	19,0	28,6	27,6	11,6	23,8	15,6	29,4	24,3	35,5
HFC 152a	-	-	-	-	-	0,1	0,1	0,2	0,1	0,1	0,1	0,1	0,0	0,1	0,0	0,0
Total	-	-	0,5	1,6	3,1	25,0	28,6	37,5	63,9	59,4	32,3	53,8	35,2	69,3	58,4	76,7




Figure 2.10 Potential emissions of HFCs by species 1990 – 2005, Gg CO₂-eq

2.2.6 Sulphur hexafluorid (SF₆)

Sulphur hexafluoride emissions are not estimated but held constant over the whole time series. The largest source of SF₆ emissions is thought to be leakages from electrical equipment.

2.3 Emission trends by source

The largest contributor of greenhouse gas emissions (without LULUCF) in Iceland is the energy sector, followed by industrial processes, agriculture, waste, geothermal energy and solvent and other product use. From 1990 to 2005 the contribution of the energy sector to the total net emissions increased from 52% to 54% respectively. The contribution of industrial processes was 26% in 1990 and 25% in 2005.

Table 2.7 Total emissions of greenhouse gases by sources (without LULUCF) in Iceland 1990 – 2005, Gg CO₂-equivalents

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Energy	1704	1658	1783	1845	1808	1820	1911	1968	1931	1971	1873	1845	1917	1862	1961	1992
Industrial Processes	867	760	566	536	509	559	535	652	798	934	950	971	936	960	949	944
Solvent Use	6	5	5	5	4	5	5	5	5	5	5	4	4	4	3	3
Agriculture	568	554	530	534	541	520	533	532	538	544	525	522	499	483	472	475
Waste	141	145	148	149	151	152	156	159	160	163	168	175	169	171	167	167
Geothermal Energy	67	67	67	67	67	82	82	71	94	123	163	154	159	138	124	123
Total	3352	3189	3099	3135	3078	3138	3222	3387	3526	3739	3684	3671	3684	3618	3678	3705

The distribution of the total greenhouse gas emissions over the UNFCCC sectors (including geothermal energy and excluding LULUCF) in 2005 is shown in Figure 2.11. Emissions from the energy sector account for 54% of the national total emissions, industrial processes account for 25% and agriculture for 13%. The waste sector accounts for 4%, geothermal energy for 3% and solvent and other product use for 0,1%.




Figure 2.11 Emissions of greenhouse gases by UNFCCC sector in 2005




Figure 2.12 Percentage changes in emissions of total greenhouse gas emissions by UNFCCC source categories during the period 1990 – 2005, compared to 1990

2.3.1 Energy

The energy sector in Iceland is unique in many ways. In 2000 the per capita energy use was close to 500 MJ, which is high compared to other industrial countries, but the proportion of domestic renewable energy in the total energy budget is 70%, which is a much higher share than in most other countries. The cool climate and sparse population calls for high energy use for space heating and transport. Iceland relies heavily on geothermal energy for space heating and on hydropower for electricity production.

The total emissions of greenhouse gases from the energy sector over the period of 1990 – 2005 are listed in Table 2.8. Figure 2.14 shows the distribution of emissions in 2005 by different source categories. The percentage change in the various source categories in the energy sector between 1990 and 2005, compared with 1990 are illustrated in Figure 2.15.

Table 2.8 Total emissions of greenhouse gases from the energy sector in 1990 – 2005, Gg CO₂-eq.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Energy industries	21	22	21	23	22	25	20	16	38	21	15	15	15	14	20	23
Manufacturing ind. & constr.	377	301	353	381	360	378	418	491	465	492	445	477	477	451	482	475
Transport	608	620	630	631	634	615	605	624	627	657	660	670	674	698	710	754
Other sectors	698	715	779	811	792	803	867	838	801	801	753	683	750	698	749	740
Fugitive emissions	NE															
Total	1704	1658	1783	1845	1808	1820	1911	1968	1931	1971	1873	1845	1917	1862	1961	1992




Figure 2.14 Greenhouse gas emissions in the energy sector 2005, distributed by source categories




Figure 2.15 Percentage changes in emissions in various source categories in the energy sector during the period 1990 – 2005, compared to 1990

As can be seen from Table 2.8 and Figure 2.15 emissions from all source categories except energy industries have increased during the period. The peak in the energy industries in 1998 was due to unusual weather condition during the winter of 1997/1998, which led to unfavourable water conditions for the hydropower reservoirs. This created a shortage of electricity which was met by burning oil for electricity and heat production.

Increased emissions from the manufacturing industries and construction source category are explained by the increased activity in the construction sector during the period.

The fisheries dominate the 'other sector'. Emissions from fisheries rose from 1990 to 1996 because a substantial portion of the fishing fleet was operating in distant fishing grounds. From 1996, the emissions decreased again reaching 1990 levels in 2001. Emissions increased again by 10% between 2001 and 2002. In 2003 emissions again reached the 1990 level. In 2005 emissions were 7% over the 1990 level. Annual changes are inherent in the nature of fisheries.

Since 1990 the vehicle fleet in Iceland has increased by 50%. This has led to increased emissions from the transport sector. The latest trend has been towards larger passenger cars, which consume more fuel. Since 1999 the average fuel consumption of newly registered passenger cars has increased by over 6%. A decrease in navigation and aviation has however compensated the effect of rising emissions in the transport sector to some extent.

2.3.2 Industrial processes

Production of raw materials is the main source of industrial process related emissions for both CO₂ and other greenhouse gases such as N₂O and PFCs. The industrial process

sector accounts for about 25% of the national greenhouse gas emissions. As can be seen from Figure 2.16 and Table 2.9 emissions decreased from 1990 to 1996, mainly because of decrease in PFC-emissions. During the late nineties large-scale industry expanded in Iceland. The existing aluminium plant and the ferroalloys industry experienced enlargement in 1997 and 1999, and in 1998 a new aluminium plant was established. This led again to an increase in industrial process emissions.

Table 2.9 Total greenhouse gas emissions from industrial processes 1990 – 2005, Gg CO₂-eq.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Mineral Products	52	49	46	40	37	38	42	47	54	62	66	59	41	33	51	55
Chemical Industry	49	47	42	44	45	43	50	42	36	37	19	17	0,5	0,5	0,4	NO
Metal Production	760	659	472	445	417	448	409	520	638	700	827	826	854	851	833	806
- Ferroalloys	203	171	182	231	225	238	227	249	192	250	354	370	389	389	387	371
- Aluminium	556	488	289	214	193	210	182	271	446	520	473	466	465	461	446	435
- Aluminium CO ₂	136	139	134	139	148	151	157	189	266	347	346	374	393	402	407	409
- Aluminium PFC	420	348	155	75	45	59	25	82	180	173	127	92	73	60	39	26
Other production	NE															
Consumption of HFCs and SF ₆	5	5	6	7	8	30	34	43	69	65	38	59	41	75	64	82
Total	866	760	565	536	508	559	535	651	798	933	949	971	935	959	948	943
Emissions fulfilling 14/CP.7									108	115	273	404	441	451	455	440




Figure 2.16 Total greenhouse gas emissions in the industrial process sector during the period from 1990 – 2005, Gg CO₂-eq.

The main category within the industrial process sector is metal production, which accounted for 88% of the sector's emissions in 1990 and 85% in 2005. Aluminium production is the main source within the metal production category, accounting for 46% of the total industrial process emissions. The production technology in both existing plants is based on using prebake anode cells. The main energy source is electricity, and industrial process CO₂ is mainly due to the anodes that are consumed during the

electrolysis. In addition, the production of aluminium gives rise to emissions of PFCs. From 1990 to 1996 PFC emissions were reduced by 94%. Because of the enlargement of the existing aluminium plant in 1997 and the establishment of a new aluminium plant in 1998, emissions increased again from 1997 to 1999 but have decreased since. In 2005 the emissions had decreased by 94% from the 1990 level. The reduction in PFC emissions was effectuated by improved technology and process control. PFC emissions per tonne aluminium produced went from 4,78 tonnes CO₂-equivalents in 1990, to 0,10 tonnes CO₂-equivalents in 2005.

Production of ferroalloys is another major source of emissions, accounting for 39% of the industrial processes emissions in 2005. CO₂ is emitted due to the use of coal and coke as reducing agents and from the consumption of electrodes. In 1998 a power shortage caused a temporary closure of the ferroalloy plant, resulting in exceptionally low emissions that year. In 1999, however, the existing plant was expanded and emissions have therefore increased considerably.

Production of minerals is the sector's second most important category, accounting for 6% of the emissions in 2005. Cement production is the dominant contributor. Cement is produced in one plant in Iceland, emitting CO₂ derived from carbon in the shell sand used as raw material in the process. Emissions from the cement industry reached a peak in 2000 but declined until 2003, partly because imports of cement. In 2004 and 2005 emissions increased again. This can be explained by increased activity related to the construction of Iceland's largest hydropower plant.

Production of fertilizers used to be the main contributor to the process emissions from the chemical industry. The production was terminated in 2001. Silicon production was terminated in 2004 and therefore no chemical industry was operated in Iceland in 2005.

Imports of HFCs started in 1992 and increased until 1998. Since then annual imports have been between 30 and 77 Gg CO₂-equivalents. Sufficient data is not available to calculate actual emissions. This means that only potential emissions, based on imports, are estimated. The potential method is likely to overstate emissions, since the chemicals used, e.g. in refrigerators, are emitted over a period of several years. The application category refrigeration contributes by far the largest part of HFCs emissions but foam blowing is also thought to be a source.

Sulphur hexafluoride emissions are not estimated but held constant over the whole time series. The largest source of SF₆ emissions is thought to be leakages from electrical equipment.

2.3.3 Solvent and other product use

The use of solvents and products containing solvents leads to emissions of non-methane volatile organic compounds (NMVOC), which are regarded as indirect greenhouse gases. The NMVOC emissions will over a period of time oxidize to CO₂ in the atmosphere. This conversion has not been estimated. The only emissions of direct GHG reported here

are due to use of N_2O , mainly for medical purposes but also to a smaller extent for car racing. Those emissions were 3 Gg CO_2 -equivalents in 2005 and have declined by 45% since 1990.

2.3.4 Agriculture

As can be seen from Table 2.10 and Figure 2.17 the emissions from agriculture decreased from 1990 to 2005. This was mainly due to a decreasing number of livestock.

Table 2.10 Total greenhouse gas emissions from agriculture in 1990 – 2005, Gg CO_2 -eq.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Enteric Fermentation	270	264	256	254	256	247	250	253	256	254	243	242	237	233	229	231
Manure Management	56	55	53	53	53	50	51	51	52	51	49	49	47	46	46	46
Agricultural Soils	241	236	221	227	233	222	233	228	230	238	234	231	215	204	198	198
Total	568	555	530	534	541	520	533	532	538	544	526	522	499	483	472	475

Greenhouse gas emissions from agriculture comprise emissions of methane and nitrous oxide. The greenhouse gas emissions from the agricultural sector accounted for 13% of the overall greenhouse gas emissions in 2005. The largest sources for agricultural greenhouse gas emissions are CH_4 from enteric fermentation and N_2O from agricultural soils.




Figure 2.17 Total greenhouse gas emissions from agriculture from 1990 – 2005, Gg CO_2 -eq.

2.3.5 Waste

As can be seen from Table 2.11 and Figure 2.18 the amount of greenhouse gases (CH_4) from landfills increased steadily from 1990 to 2001. From 2002 to 2005 a minor decrease in emissions occurred. From 1990 to 2005 the emissions rose by 36%. There are two reasons for this, increasing amounts of waste being landfilled and a larger

percentage of that waste being landfilled in managed waste disposal sites. The amount of landfilled waste increased by 37% over the period. Methane recovery started at the largest operating landfill site in 1997, and the amount recovered has increased steadily since then.

Emissions from wastewater handling have increased constantly since 1990 because total number of inhabitants connected to wastewater facilities has increased in the time period.

Emissions from waste incineration have decreased constantly since 1990 because total amount of waste being incinerated in Iceland has decreased. A higher percentage of the waste has concurrently been incinerated with energy recovery and is thus reported under 1A1a (public electricity and heat production).

Table 2.11 Emissions from the waste sector from 1990 – 2005, Gg CO₂-eq.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Landfills	114	118	121	124	127	131	136	139	142	146	151	158	150	153	152	154
Wastewater Handling	8	8	8	8	8	8	8	8	9	9	10	10	13	13	13	13
Waste Incineration	19	19	18	15	14	13	11	11	9	8	7	7	6	5	2	0,03
	140	144	147	148	150	151	155	158	160	163	169	174	168	171	167	167




Figure 2.18 Emissions of greenhouse gases in the waste sector 1990 – 2005, Gg CO₂-eq.

2.3.6 Geothermal energy

Iceland relies heavily on geothermal energy for space heating and to some extent for electricity production. Researches indicate that CO₂ emissions associated with the utilization of geothermal energy in Iceland constitute a net increase in emissions. The emissions are though considerably less extensive than from fossil fuel power plants. Table 2.12 shows the emissions from geothermal energy from 1990 to 2005.

Table 2.12 Emissions from geothermal energy from 1990 – 2005, Gg CO₂-eq.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total	67	67	67	67	67	82	82	71	94	123	163	154	159	138	124	123

2.4 Emission trends for indirect greenhouse gases and SO₂

Nitrogen oxides (NO_x), non-methane volatile organic compounds (NMVOC) and carbon monoxide (CO) have an indirect effect on climate through their influence on greenhouse gases, especially ozone. Sulphur dioxide (SO₂) affects climate by increasing the level of aerosols that have in turn a cooling effect on the atmosphere.

2.4.1 Nitrogen oxides (NO_x)

As can be seen in Figure 2.19, the main sources of nitrogen oxides in Iceland are fishing, transport and manufacturing industry and construction. The NO_x emissions from fishing rose from 1990 to 1996 when a substantial portion of the fishing fleet was operating in distant fishing grounds. From 1996 emissions decreased, reaching the 1990 levels in 2001. In 2005 emissions were 7% above the 1990 level. Annual changes are inherent in the nature of fisheries. Emissions from transport are dominated by road transport. These emissions have decreased rapidly after the use of catalytic converters in all new vehicles became obligatory in 1995. The rise in emissions from the manufacturing industries and construction are dominated by increased activity in the construction sector during the period. Total NO_x emissions show, like the emissions from fishing, an increase until 1996 and then a decrease until 2001. Emissions have been rising again since 2001. The emissions in 2005 were 7% above the 1990 level.




Figure 2.19 Emissions of NOx by sector 1990 – 2005, Gg

2.4.2 Non-methane volatile organic compounds (NMVOC)

As can be seen in Figure 2.20 the main sources of non-methane volatile organic compounds are transport and solvent use. Emissions from transport are dominated by road transport. These emissions have decreased rapidly after the use of catalytic converters in all new vehicles became obligatory in 1995. Emissions from solvent use vary between 2 Gg and 4 Gg during the period with no obvious trend. The total emissions show a downward trend from 1994 to 2003 with exception of 2000 and 2001. The emissions in 2005 were 34% below the 1990 level.




Figure 2.20 Emissions of NMVOC by sector 1990 – 2005, Gg

2.4.3 Carbon monoxide (CO)

As can be seen in Figure 2.21, transport is the prominent contributor to CO emissions in Iceland. Emissions from transport are dominated by road transport. These emissions have decreased rapidly after the use of catalytic converters in all new vehicles became obligatory in 1995. Total CO emissions show, like the emissions from transport, a rapid decrease after 1990. The emissions in 2005 were 57% below the 1990 level.




Figure 2.21 Emissions of CO by sector 1990 – 2005, Gg

2.4.4 Sulphur dioxide (SO₂)

Geothermal energy exploitation is by far the largest source of sulphur emissions in Iceland and has increased by 160% since 1990 due to increased activity in this field. Other significant sources of sulphur dioxide in Iceland are industrial processes and manufacturing industry and construction, as can be seen in Figure 2.22. Emissions from industrial processes are dominated by metal production. Until 1996 industrial process sulphur dioxide emissions were relatively stable. During the late nineties the metal industry expanded. The existing aluminium plant and the ferroalloys industry experienced enlargement in 1997 and 1999, and in 1998 a new aluminium plant was established. This led to increased emissions of sulphur dioxide. The fishmeal industry is the main contributor to sulphur dioxide emissions in the sector ‘manufacturing industries and construction’. Emissions from the fishmeal industry increased generally from 1990 to 1997 but have declined since and were 22% below the 1990 level in 2005.

Total SO₂ emissions in 2005 were 106% above the 1990 level.




Figure 2.22 Emissions of SO₂ by sector 1990 – 2005, Gg

3 ENERGY

3.1 Overview

The energy sector in Iceland is unique in many ways. In 2000 the per capita energy use was nearly 500 MJ, which is high compared to other industrial countries. However, the proportion of domestic renewable energy in the total energy budget is 70%, which is a much higher share than in most other countries. The cool climate and sparse population calls for high energy use for space heating and transport. Iceland relies heavily on its geothermal energy sources for space heating and on hydropower for its electricity production. Emissions from hydropower reservoirs are included in the LULUCF sector. Emissions from geothermal energy are discussed in Chapter 9.

The energy sector accounts for 54% of the GHG emissions in Iceland. Emissions increased by 17% from 1990 to 2005. From 2004 to 2005 the emissions increased by 2%. Fisheries and road traffic are the sector's largest single contributors. Combustion in the manufacturing industries and construction is also an important source.

3.1.1 Methodology

The calculation of greenhouse gas emissions from fuel combustion activities is based on the methodologies suggested by the IPCC Guidelines and the Good Practice Guidance. Fuel combustion activities are divided into two main categories: stationary and mobile combustion. Stationary combustion includes energy industries, manufacturing industries and the other sector (residential and commercial/institutional sector). Mobile combustion includes civil aviation, road transport, navigation, fishing, mobile combustion in construction and international bunkers. The methodology applied for each source category is described below.

The key source analysis performed for 2005 has revealed, as indicated in Table 1.1, that in terms of total level and/or trend uncertainty the key sources in the Energy sector are the following:

- Stationary combustion: oil – CO₂ (1A1, 1A2, 1A4)
- Stationary combustion: coal – CO₂ (1A2f)
- Mobile combustion: construction – CO₂ (1A2f)
- Mobile combustion: road vehicles – CO₂ (1A3b)
- Mobile combustion: road vehicles – N₂O (1A3b)
- Mobile combustion: fishing – CO₂ (1A4c)

3.1.2 Completeness

Table 3.1 gives an overview of the IPCC source categories included in this chapter and presents the status of emission estimates from all sub-sources in the energy sector.

Table 3.1 Energy - completeness

Sector	Greenhouse gases						Other gases			
	CO ₂	CH ₄	N ₂ O	HFC	PFC	SF ₆	Nox	CO	NMVOC	SO ₂
Energy industries										
Public electricity and heat production	X	X	X	NA	NA	NA	X	X	X	X
Petroleum refining	NOT OCCURRING									
Manufacture of Solid Fuels and other energy industries	NOT OCCURRING									
Manufacturing Industries and Construction										
Iron and Steel	X	X	X	NA	NA	NA	X	X	X	X
Non-ferrous metals	X	X	X	NA	NA	NA	X	X	X	X
Chemicals	X	X	X	NA	NA	NA	X	X	X	X
Pulp, paper and print	NOT OCCURRING									
Food Processing, Beverages and Tobacco	X	X	X	NA	NA	NA	X	X	X	X
Other	X	X	X	NA	NA	NA	X	X	X	X
Transport										
Civil Aviation	X	X	X	NA	NA	NA	X	X	X	X
Road Transportation	X	X	X	NA	NA	NA	X	X	X	X
Railways	NOT OCCURRING									
Navigation	X	X	X	NA	NA	NA	X	X	X	X
Other Transportation	NOT OCCURRING									
Other Sector										
Commercial/Institutional	X	X	X	NA	NA	NA	X	X	X	X
Residential	X	X	X	NA	NA	NA	X	X	X	X
Agriculture/Forestry/Fisheries	X	X	X	NA	NA	NA	X	X	X	X
Other										
Stationary	X	X	X	NA	NA	NA	X	X	X	X
Mobile	NOT OCCURRING									
Fugitive Emissions from Fuels										
Solid Fuels	NOT OCCURRING									
Oil and Natural Gas	NE	NE	NE	NA	NA	NA	NE	NE	NE	NE
International Transport										
Aviation	X	X	X	NA	NA	NA	X	X	X	X
Marine	X	X	X	NA	NA	NA	X	X	X	X

3.2 Stationary fuel combustion

3.2.1 Energy industries, manufacturing industries, commercial/institutional and residential fuel combustion

This source refers to emissions of GHG from combustion of fuels in the energy industries, in the manufacturing industries and construction and in the 'other sector' (commercial/institutional and residential fuel combustion activities). The key source analysis shows that CO₂ emissions from stationary oil combustion constitute a key source in both level and trend and CO₂ emissions from stationary coal combustion constitute a key source in trend.

Emissions from fuel combustion are estimated at the sectoral level. They are calculated by multiplying energy use by source and sector with pollutant specific emission factors. Activity data is provided by the National Energy Forecast Committee (NEFC), which collects data from the oil companies on fuel sales by sector. Since not all oil companies provided sales statistics for the years 2003 to 2005, fuel use by sector has been estimated by the NEFC. Since Iceland relies heavily on geothermal energy for space heating and hydropower for electricity production, emissions from those sectors are relatively low. Emissions in this key source originate predominantly from the combustions in the manufacturing industries, and the fishmeal industry in particular. Emissions from waste incineration with energy recovery are reported under 'energy industries' and a description of the method is in Chapter 8.

Activity data

Total use of different oil products is based on the NEFCs annual sales statistics for fossil fuels. The data is, with the exception of 2003 to 2005, considered reliable since all oil companies report their sales statistics. There is thus a given total, which usage in the different sectors must sum up to. There is not a clear distinction between the energy industries sector and residential sector in fuel sales statistics. The National Energy Authority (NEA) has on request by the Environment and Food Agency (EFA) divided the fuel consumption between the two sectors. The EFA collects consumption data from all major industry installations and the consumption in the fishmeal industry is from 1990 to 2002 estimated from production statistics, but for 2003 to 2005 they are based on data provided by the industry.

Emission factors

The CO₂ emission factors (EF) used reflect the average carbon content of fossil fuels. They are taken from the revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories and are presented in Table 3.2. SO₂ emissions are calculated from the S-content of the fuels. Emission factors for other pollutants are taken from Table 1-15 to 1-19 of the revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories: Reference Manual. Where EF were not available the default EF from Tables 1-7 to 1-11 in the Reference Manual were used.

Table 3.2 Emission factors for CO₂ from stationary combustion

	NCV [TJ/kt]	Carbon EF [t C/TJ]	Fraction oxidised	CO ₂ EF [t CO ₂ /t fuel]
Kerosene (heating)	44,75	19,60	0,99	3,18
Gas / Diesel Oil	43,33	20,20	0,99	3,18
Residual fuel oil	40,19	21,10	0,99	3,08
Coking Coal	28,05	25,80	0,98	2,60

Uncertainty

The preliminary estimate of quantitative uncertainty has revealed that the uncertainty of CO₂ emissions from stationary oil combustion is 7% and for coal combustion 11%.

Recalculations

In this submission combustion of clinical waste is included for the first time. Due to this a minor change is observed from 2002 to 2004. Also the combustion of petroleum coke in the cement industry is now reported in 2004.

3.3 Mobile combustion

3.3.1 Construction sector

Emissions are calculated by multiplying energy use with a pollutant specific emission factor. Activity data is provided by the NEFC, which collects data on fuel sales by sector.

Activity data

Total use of oil products in the construction sector is based on the NEFC's annual sales statistics for fossil fuels. The data is, with the exception of 2003 to 2005, considered reliable since all the oil companies have reported their sales statistics. In some instances oil, which is reported to fall under vehicle usage, is actually used for machinery and vice versa. This is, however, very minimal and the deviation is believed to level out.

Emission factors

The CO₂ emission factors used reflect the average carbon content of fossil fuels. The emission factors for other pollutants are taken from Table 1-49 in the revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories: Reference Manual. EF for CO₂ and N₂O are presented in Table 3.3.

Table 3.3 Emission factors for CO₂ and N₂O from combustion in the construction sector

	NCV [TJ/kt]	Carbon EF [t C/TJ]	Fraction oxidised	CO ₂ EF [t CO ₂ /t fuel]	N ₂ O EF [t N ₂ O/kt fuel]
Gas / Diesel Oil	43,33	20,20	0,99	3,18	1,3

Uncertainties

The preliminary estimate of quantitative uncertainty has revealed that the uncertainty of CO₂ emissions from construction is 11%.

3.3.2 Road vehicles

Emissions from road traffic are estimated by multiplying the fuel use by type of fuel and vehicle, and fuel and vehicle pollutant specific emission factors. Activity data is provided by the NEFC, which collects data on fuel sales by sector.

Activity data

Total use of diesel oil and gasoline are based on the NEFC's annual sales statistics for fossil fuels. The data is, with the exception of 2003 to 2005, considered reliable since all the oil companies have reported sales statistics. The EFA estimates how fuel sale is

divided between the different types of vehicles, but the method used is considered to be inaccurate.

Emission factors

For CO₂ the standard emission factors based on carbon content of the fuels are used. Emission factors for CH₄ and N₂O depend upon vehicle type and emission control. They are taken from the revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories and are presented in Table 3.4.

Table 3.4 Emission factors for greenhouse gases from European vehicles, g/kg fuel

	CH ₄	N ₂ O	CO ₂
Passenger car – gasoline, uncontrolled	0,8	0,06	3070
Passenger car – gasoline, non catalyst control	1,1	0,08	3070
Passenger car – gasoline, three way catalyst	0,3	0,8	3070
Light duty vehicle – gasoline	0,8	0,06	3070
Heavy duty vehicle – gasoline	0,7	0,04	3070
Passenger car – diesel	0,08	0,2	3180
Light duty vehicle – diesel	0,06	0,2	3180
Heavy duty vehicle – diesel	0,2	0,1	3180

Uncertainties

The preliminary estimate of quantitative uncertainty has revealed that the uncertainty of CO₂ emissions from road vehicles is 11%. For N₂O, both activity data and emission factors are highly uncertain. The uncertainty of N₂O emissions from road vehicles is 206%.

3.3.3 Fishing

Emissions are calculated by multiplying energy use with a pollutant specific emission factor. Activity data is provided by the NEFC, which collects data on fuel sales by sector.

Activity data

Total use of residual fuel oil and gas/diesel oil for the fishing is based on the NEFC's annual sales statistics for fossil fuels. The data is, with the exception of 2003 and 2004 considered reliable since all oil companies reported their sales statistics.

Emission factors

The emission factors are taken from the revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories for ocean-going ships and are presented in Table 3.5.

Table 3.5 Emission factors for CO₂, CH₄ and N₂O for ocean going ships

	NCV [TJ/kt]	Carbon EF [t C/TJ]	Fraction oxidised	EF CO ₂ [t CO ₂ /t]	EF N ₂ O [kg N ₂ O/TJ]	N ₂ O EF [kg N ₂ O/t]	EF CH ₄ [kg CH ₄ /TJ]	EF CH ₄ [kg CH ₄ /t]
Gas / Diesel Oil	43,33	20,20	0,99	3,18	2	0,086	7	0,30
Residual fuel oil	40,19	22,00	0,99	3,08	2	0,084	7	0,28

Uncertainties

The preliminary estimate of quantitative uncertainty has revealed that the uncertainty of CO₂ emissions from fishing is 5%.

3.3.4 Civil aviation

Emissions are calculated by using the Tier 1 method, thus multiplying energy use with a pollutant specific emission factor. Activity data is provided by the NEFC, which collects data on fuel sales by sector.

Activity data

Total use of jet kerosene and gasoline is based on the NEFC's annual sales statistics for fossil fuels. The data is, with the exception of 2003 to 2005 considered reliable since all the oil companies reported their sales statistics.

Emission factors

The emission factors are taken from the revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories and are presented in Table 3.6. Emissions of SO₂ are calculated from S-content in the fuels.

Table 3.6 Emission factors for CO₂ and other pollutants for aviation

	NCV [TJ/kt]	Carbon EF [t C/TJ]	Fraction oxidised	EF CO ₂ [t CO ₂ /t]	NO _x [kg/TJ]	CH ₄ [kg/TJ]	NMVOC [kg/TJ]	CO [kg/TJ]	N ₂ O [kg/TJ]
Jet kerosene	44,59	19,50	0,99	3,16	300	0,5	50	100	2
Gasoline	44,80	18,90	0,99	3,07	300	0,5	50	100	2

3.3.5 National navigation

Emissions are calculated by multiplying energy use with a pollutant specific emission factor. Activity data is provided by the National Energy Forecast Committee (NEFC), which collects data on fuel sales by sector.

Activity data

Total use of residual fuel oil and gas/diesel oil for national navigation is based on the NEFC's annual sales statistics for fossil fuels. The data is, with the exception of 2003 to 2005 considered reliable since all the oil companies reported their sales statistics.

Emission factors

The emission factors are taken from the revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories for ocean-going ships and are presented in Table 3.5.

3.4 International bunker fuels

Emissions from international aviation and marine bunker fuels are excluded from national totals as required according to the IPCC Guidelines. These emissions are presented separately for informational purposes and can be seen in Table 3.6.

In 2005, greenhouse gas emissions from ships and aircraft in international traffic bunkered in Iceland amounted to a total of 593 Gg CO₂-equivalents, which corresponds to about 16% of the total Icelandic greenhouse gas emissions. Greenhouse gas emissions from marine and aviation bunkers increased by around 85% from 1990 to 2005, and between 2004 and 2005 emissions decreased by 2%.

Looking at these two categories separately, it can be seen that greenhouse gas emissions from international marine bunkers increased by 83% from 1990 to 2005, while emissions from aircrafts increased by 85% during the same period. Between 2004 and 2005 emissions from marine bunkers decreased by 20% while emissions from aviation bunkers increased by 10%.

Table 3.7 Greenhouse gas emissions from international aviation and marine bunkers, Gg CO₂-eq.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Marine	100	38	61	98	94	146	125	150	178	166	221	151	209	181	229	183
Aviation	222	224	205	197	216	238	274	295	341	367	411	352	313	333	374	411
Total	322	262	266	296	310	384	399	445	520	532	632	503	522	514	603	593

Emissions are calculated by multiplying energy use with pollutant specific emission factors. Activity data is provided by the NEFC, which collects data on fuel sales by sector. These data distinguish between national and international usage. The data is with the exception of 2003 to 2005 considered reliable since all oil companies selling oil products report those statistics. The emission factors for marine bunkers are taken from the revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories for ocean-going ships and are presented in Table 3.5 above. Emission factors for aviation bunkers are also taken from the IPCC Guidelines and presented in Table 3.6 above.

3.5 Cross-cutting issues

3.5.1 Sectoral versus reference approach

This section will be completed for the next submission.

3.5.2 Feedstock and non-energy use of fuels

Emissions from the use of feedstocks are according to the Good Practice Guidance accounted for in the industrial processes sector in the Icelandic inventory. This includes all use of petroleum coke, other bituminous coal and coke oven coke.

Iceland uses a carbon storage factor of 1 for bitumen and 0,5 for lubricants for the non-energy use in the Reference Approach, CRF Table 1(A)d.

4 INDUSTRIAL PROCESSES

4.1 Overview

Production of raw materials is the main source of industrial process related emissions for CO₂, N₂O and PFCs. Emissions also occur because of use of HFCs as substitutes for ozone depleting substances. The industrial process sector accounted for 25% of the GHG emissions in Iceland in 2005. Emissions decreased from 1990 to 1996, mainly due to reduction in PFC emissions. In 1996 emissions were 38% below the 1990 level. Due to the expansion of energy intensive industry, emissions rose rapidly from 1996 to 1999, when they were 8% above the 1990 level. In 2005 emissions from the industrial processes sector were 9% above the 1990 level. The main category within the industrial process sector is metal production, which accounted for 85% of the sector's emissions in 2005.

4.1.1 Methodology

The calculation of greenhouse gas emissions from industrial processes is based on the methodologies suggested by the IPCC Guidelines and the Good Practice Guidance.

The key source analysis performed for 2005 has revealed, as indicated in Table 1.1, that in terms of total level and/or trend uncertainty the key sources in the Industrial Processes Sector are the following:

- Emissions from Cement Production – CO₂ (2A1)
- Emissions from Ferroalloys – CO₂ (2C2)
- Emissions from Aluminium Production – CO₂ (2C3)
- Emissions from Aluminium Production – PFCs (2C3)
- Emissions from Substitutes for Ozone Depleting Substances – HFCs (2F)

4.1.2 Completeness

Table 4.1 gives an overview of the IPCC source categories included in this chapter and presents the status of emission estimates from all sub-sources in the industry sector.

Table 4.1 Industrial Processes - Completeness

Sector	Greenhouse gases						Other gases			
	CO ₂	CH ₄	N ₂ O	HFC	PFC	SF ₆	NO _x	CO	NM VOC	SO ₂
Mineral Products:										
Cement Production	X	NE	NE	NA	NA	NA	NE	NE	NE	IE
Lime Production	NOT OCCURRING									
Limestone and Dolomite Use	NOT OCCURRING									
Soda Ash Production and Use	NOT OCCURRING									
Asphalt Roofing	NOT OCCURRING									
Road Paving with Asphalt	NE	NE	NE	NA	NA	NA	X	X	X	X
Other (Mineral Wool Production)	X	NE	NE	NA	NA	NA	NE	X	NE	X
Chemical Industry										
Ammonia Production	NOT OCCURRING									
Nitric Acid Production	NOT OCCURRING									
Adipic Acid Production	NOT OCCURRING									
Carbide Production	NOT OCCURRING									
Other (Silicium Production – until 2004)*	X	NE	NE	NA	NA	NA	X	NE	NE	NE
Other (Fertilizer Production – until 2001)*	NA	NE	X	NA	NA	NA	X	NE	NE	NE
Metal Production										
Iron and Steel Production	NOT OCCURRING									
Ferroalloys Production	X	X	NA	NA	NA	NA	X	X	X	X
Aluminium Production	X	NE	NE	NA	X	NA	NE	NE	NE	X
SF ₆ used in aluminium/magnesium foundries	NOT OCCURRING									
Other	NOT OCCURRING									
Other Production										
Pulp and Paper	NOT OCCURRING									
Food and Drink	NE	NA	NA	NA	NA	NA	NA	NA	NE	NA
Production of Halocarbons and SF₆	NOT OCCURRING									
Consumption of Halocarbons and SF₆	NA	NA	NA	X	NO	X	NA	NA	NA	NA
Other	NOT OCCURRING									

* Fertilizer production was terminated in 2001 and Silicium production was terminated in 2004

4.2 Mineral Products

4.2.1 Cement Production (2A1)

Emissions of CO₂ originate from the calcination of the raw material calcium carbonate, which comes from shell sand in the production process. The resulting calcium oxide is heated to form clinker and then crushed to form cement. Emissions are calculated according to the Tier 2 method based on clinker production data and data on the CaO content of the clinker. Cement Kiln Dust (CKD) is non-calcined dust produced in the kiln. CKD may be partly or completely recycled to the kiln. Any CKD that is not recycled can be considered lost to the system in terms of CO₂ emissions. Emissions are thus corrected with plant specific cement kiln dust correction factor.

Activity data

Process specific data on clinker production, CaO content of the clinker and non-recycled CKD are collected by the EFA directly from the cement production plant. The data is considered reliable. Data on clinker production is only available for the years 2003 to 2005. Historical clinker production data has been calculated as 85% of cement production.

Table 4.2 Clinker production and CO₂ emissions from cement production from 1990 – 2005.

Year	Clinker production [kt]	CO ₂ emissions [kt]
1990	114,100	51,6
1991	106,174	48,0
1992	99,800	45,1
1993	86,419	39,1
1994	80,856	36,5
1995	81,514	36,8
1996	90,325	40,8
1997	100,625	45,5
1998	117,684	53,2
1999	133,647	60,4
2000	142,604	64,4
2001	127,660	57,7
2002	84,684	39,4
2003	75,314	32,1
2004	104,829	49,8
2005	99,170	53,9

Emission factors

It has been estimated by the cement production plant that CaO content of the clinker is 63%. The corrected emission factor for CO₂ is thus 0,495. For CKD it is 7,5% for all years except 2005 when it is 110%.

Uncertainties

The preliminary estimate of quantitative uncertainty has revealed that the uncertainty of CO₂ emissions from cement production is 6,5%.

4.2.2 Road paving with asphalt

Asphalt road surfaces are composed of compacted aggregate and asphalt binder. Gases are emitted from the asphalt plant, the road surfacing operations and from the subsequent road surface. Information on the amount of asphalt produced come from Statistics Iceland. Emission factor for SO₂, NO_x, CO and NMVOC are taken from Table 2-4, IPCC Guidelines, Reference Manual.

4.2.3 Mineral Wool Production

Emissions of CO₂ and SO₂ are calculated from the amount of shell sand and electrodes used in the production process. Emissions of CO are based on measurements that were made at the single operating plant in 2000.

4.3 Chemical industry

The only chemical industry that has existed in Iceland is the production of silicium and fertilizer. The fertilizer production plant was closed down in 2001 and the silicium production plant was closed down in 2004.

At the silicium production plant silicium containing sludge is burned to remove organic material. Emissions of CO₂ and NO_x are estimated on the basis of C-content and N-content of the sludge. Emissions also occur from the use of soda ash at the production process and those emissions are reported here.

When the fertilizer production plant was operational it reported its emissions of NO_x and N₂O to the Environment and Food Agency.

4.4 Metal Production

4.4.1 Ferroalloys

Emissions of CO₂ originate from the use of coal and coke as reducing agent, as well as from consumption of electrodes. Emissions are calculated according to the Tier 1 method based on the consumption of reducing agents and electrodes and emission factors from the IPCC Guidelines.

Activity data

The consumption of reducing agents and electrodes are collected by the EFA directly from the single operating ferroalloys production plant. The data is thus considered reliable.

Emission factors

For CO₂, the standard emission factors based on carbon content of the reducing agents and electrodes are used. They are taken from the revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories and are presented in Table 4.3. Values for NCV are from NEA. Emission factors for CH₄, NO_x and NMVOC are taken from Tables 1-7, 1-9 and 1-11 in the IPCC Guidelines, Reference Manual. Emissions of SO₂ are calculated from the sulphur content of the reducing agents and electrodes. The emission factor for CO is taken from Table 2-16 in the IPCC Guidelines, Reference Manual.

Table 4.3 Emission factors for CO₂ from production of ferroalloys

	NCV [TJ/kt]	Carbon EF [t C/TJ]	Fraction oxidised	CO ₂ EF [t CO ₂ /t input]
Other Bituminous Coal	28,00	25,80	0,98	2,60
Coke Oven Coke	28,00	29,50	0,98	2,97
Electrodes	28,00	32,14	0,98	3,23

Uncertainties

The preliminary estimate of quantitative uncertainty has revealed that the uncertainty of CO₂ emissions from ferroalloys production is 11%.

4.4.2 Aluminium Production

Primary aluminium production results in emissions of CO₂ and PFCs. CO₂ emissions originate from the consumption of electrodes. Emissions are calculated according to the Tier 1 method based on the quantity of electrodes used in the process and the emission factors from the IPCC Guidelines.

PFCs are produced during anode effects (AE) in the prebake cells, when the voltage of the cells increases from the normal 4 – 5 V to 25 – 40 V. Emissions of PFCs are dependent on the number of anode effects and their intensity and duration. Anode effect characteristics are different from plant to plant. Emission factors are calculated according to the Tier 2 Slope Method. Default coefficients are taken from the IPCC Good Practice Guidance for Centre Worked Prebaked Technology. Emission factors are calculated with the following formula:

$$EF \text{ (kg CF}_4 \text{ or C}_2\text{F}_6 \text{ per tonne of Al)} = \text{Slope} \cdot \text{AE min/cell day}$$

Emissions are then calculated by multiplying the emission factors with the amount of aluminium produced.

Activity data

The EFA collects annual process specific data from the two operating aluminium plants. The data is considered reliable.

Emission factors

For CO₂, the standard emission factors based on carbon content of the electrodes are used. They are taken from the revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories and are presented in Table 4.2. The default coefficients for the calculation of PFC emissions are taken from the IPCC Good Practice Guidance for Centre Worked Prebaked Technology (0,14 for CF₄ and 0,018 for C₂F₆). The emissions calculated in this way seem to fit well to the measurements that have been performed at both plants. The measurements took place in 1997 at Iceland's first aluminium plant and in 2001 at the newer one.

Table 4.4 Emission factors CO₂ from aluminium production

	NCV [TJ/kt]	Carbon EF [t C/TJ]	Fraction oxidised	CO ₂ EF [t CO ₂ /t input]
Electrodes	31,35	31,42	0,98	3,54

Table 4.5 Aluminium production, AE, CO₂ and PFC emissions from 1990 – 2005.

Year	Aluminium production [kt]	CO ₂ emissions [kt]	AE Andoe Effect [min/cell day]	PFC emissions [kt CO ₂ -eq]	CO ₂ [t/t Al]	PFC [t/t Al]
1990	87,839	136,5	4,44	419,6	1,55	4,78
1991	89,217	139,3	3,63	348,3	1,56	3,90
1992	90,045	134,2	1,60	155,3	1,49	1,72
1993	94,152	139,0	0,74	74,9	1,48	0,80
1994	98,595	148,0	0,42	44,6	1,50	0,45
1995	100,198	150,7	0,55	58,84	1,50	0,59
1996	103,362	157,0	0,23	25,2	1,52	0,24
1997	123,562	188,9	0,62	82,4	1,53	0,67
1998	173,869	265,5	10,90	180,1	1,53	1,04
1999	222,014	347,2	2,17	173,2	1,56	0,78
2000	226,362	345,5	1,13	127,2	1,53	0,56
2001	244,148	373,9	0,71	91,7	1,53	0,38
2002	264,107	392,6	0,56	72,5	1,49	0,27
2003	266,611	401,6	0,40	59,8	1,51	0,22
2004	271,384	407,3	0,25	38,6	1,50	0,14
2005	272,488	408,7	0,22	26,1	1,50	0,10

Uncertainties

The preliminary estimate of quantitative uncertainty has revealed that the uncertainty of CO₂ emissions from aluminium production is 11%.

The emission factors for calculating PFC emissions have more uncertainty but still seem to fit well to the measurements that have been performed so far at the aluminium production plants. The preliminary estimate of quantitative uncertainty has revealed that the uncertainty of PFC emissions from aluminium production is 9% for CF₄ and 23% for C₂F₆.

Recalculations

The amount of LPG used at one aluminium plant in 2003 and 2004 were reversed. This was corrected in this submission.

4.5 Emissions from Substitutes for Ozone Depleting Substances – HFCs (2F)

HFCs are used as substitutes for the ozone depleting substances (CFCs and HCFCs) which being phased out by the Montreal Protocol. There is no production of HFCs in Iceland but they are imported for use in stationary and mobile air-conditioning systems, metered dose inhalers and foam blowing. HFC is banned in other aerosols, solvents and

fire extinguishers. The HFCs used in significant quantities in Iceland are HFC-23, HFC-32, HFC-125, HFC-134a, HFC-143a and HFC-152a.

The import of HFCs started in 1992 and increased until 1998. The annual imports have since stayed between 30 and 77 Gg CO₂-equivalents. Sufficient data are still not available to calculate actual emissions in most applications, meaning that only potential emissions, based on registered imports, are estimated. However, estimates of HFC import in equipment are now being ascertained and will be included in potential emissions next year. In 2005 the potential emissions of HFCs were about 2% of national total greenhouse gas emissions (without LULUCF). This source category is a key source in both level and trend.

Method

Potential emissions of HFCs (sector 2F) are calculated using the Tier 1b methodology which considers the import, export and destruction of chemicals in bulk and in equipment without time lag.

Data on imported and exported bulk are reported directly to the EFA each year. The data are considered reliable. There is no destruction of HFCs in Iceland, although small amounts are exported every year for destruction at a facility in Denmark.

Activity data

Information on the import of chemicals in bulk is reported directly to the EFA. The importers are required to report on the type and amount of HFC they are importing in order to release the chemicals from the customs agency. At present, there is no registration of HFC in imported refrigeration equipment or vehicles.

Uncertainty

The quantitative uncertainty has not been evaluated. The activity data are considered reliable. Emissions are likely to be overestimated since only potential emissions are calculated.

Planned improvements

Considerable progress has been made towards improving estimations for this source for 2006 estimates. Much has been done already to assess the amount of HFC imported in refrigeration equipment and in mobile air-conditioning systems.

It now is believed that the levels of HFCs imported in mobile air-conditioning systems (MACs) are higher than previously assumed. A voluntary survey of car importers was conducted this year and the resulting data, along with import statistics, will be used to extrapolate data for levels of imports in MACs. According to the survey the import of

vehicles with mobile air conditioning (MAC) systems has increased in recent years. Preliminary data suggest that 8,5 thousands tonnes of HFC-134a (CO₂-equivalents) were imported in vehicles in 2005 alone. Considering that 77 thousand tonnes of HFC (CO₂-equivalents) were imported in bulk in 2005, the amount imported in MAC systems is considerable and will be included in next year's inventory.

Data on HFCs in refrigeration equipment will be estimated from import statistics, based on land of origin and type of refrigerator.

A considerably smaller source is the use of HFCs in aerosols, or metered dose inhalers. Their import is registered by the Icelandic Medicines Control Agency. This category will be moved to Tier 2 by next submission.

5 SOLVENT AND OTHER PRODUCT USE

The use of solvents and products containing solvents leads to emissions of non-methane volatile organic compounds (NMVOC), which are regarded as indirect greenhouse gases. Estimates of NMVOC emissions are based on data on imports of solvents combined, with expert estimates based on surveys. The NMVOC emissions will over a period of time oxidize to CO₂ in the atmosphere. This conversion has not been estimated.

The only emissions of direct GHG reported under the sector solvent and other product use are due to use of N₂O, mainly for medical purposes, and also, to a smaller extent, for car racing. Data on sold amounts are collected directly by the Environment and Food Agency.

6 AGRICULTURE

6.1 Overview

Icelanders are self-sufficient in all major livestock products such as meat, milk and eggs. Traditional livestock production is grassland-based and all the native breeds, i.e. of dairy cattle, sheep, horses and goats are of ancient Nordic origin, one of each species. These animals are generally smaller than in Europe. Beef production, however, is partly based on imported breeds. The more intensive agricultural sector, pork and poultry production is based on imported breeds.

6.1.1 Methodology

The calculation of greenhouse gas emissions from agriculture is based on the methodologies suggested by the IPCC Guidelines and the Good Practice Guidance.

As indicated in Table 1.1, the key source analysis performed for 2005 has revealed that in terms of total level and/or trend uncertainty the key sources in the agriculture sector are as follows:

- Emissions from Enteric Fermentation – CH₄ (4A)
- Direct Emissions from Agricultural Soils – N₂O (4D1)
- Indirect Emissions from Agricultural Soils – N₂O (4D2)

6.1.2 Completeness

Table 6.1 gives an overview of the IPCC source categories included in this chapter and presents the status of emission estimates from all sub-sources in the agricultural sector.

Table 6.1 Agriculture - completeness

Sector	Greenhouse gases		
	CO ₂	CH ₄	N ₂ O
Enteric Fermentation	NA	X	NA
Manure Management	NA	X	X
Rice Cultivation	Not Occurring		
Agricultural Soils			
Direct emissions	NA	NE	X
Animal Production	NA	NE	X
Indirect emissions	NA	NE	X
Prescribed burning of Savannas	Not Occurring		
Field burning of agricultural residues	Not Occurring		
Other	Not Occurring		

6.2 Enteric Fermentation

The production of CH₄ by enteric fermentation in animals varies with digestive systems and feed intake. Ruminants such as cattle and sheep produce the largest amount of methane. However, enteric fermentation in pseudo-ruminants (e.g. horses) and monogastric animals (e.g. pigs) is also of significance. The methodology for calculating

methane from enteric fermentation is in accordance with the Tier 1 method. Both the population levels and emission factors by type of animal are used to calculate the emissions.

Activity data

The Icelandic Association of Farmers (IAF) is in charge of assessing the size of the animal population each year. On request from the EFA, the IAF also accounts for young animals, but those are mostly excluded from national statistics on animal population. The data is considered relatively reliable.

Emission factors

Emission factors are taken from the IPCC Guidelines. They are presented in Table 6.2. The emission factors are likely to be too high, since domestic animals in Iceland are generally smaller (sheep, horses) than in other European countries.

Table 6.2 Emission factors for CH₄ from enteric fermentation

	kg CH ₄ per head per year
Dairy cattle	100
Non-dairy cattle	48
Sheep	8
Goats	5
Horses	18
Swine	1,5

Uncertainties

The preliminary estimate of quantitative uncertainty has revealed that the uncertainty of CH₄ emissions from enteric fermentation is 54%.

Planned improvements

Develop country-specific emission factors from feed intake according to the Tier 2 method, in particular for the special Icelandic livestock.

6.3 Manure management

Manure management is responsible for methane and nitrous oxide emissions. Methane is produced during the anaerobic decomposition of the manure, while nitrous oxide is produced during storage and treatment of the manure before it is used as fertilizer.

CH₄ emissions from manure management were estimated according to the IPCC Tier 1 methodology. Population levels for each kind of animal as well as its specific emission factor are used to calculate the emissions. The animal population size is collected, as mentioned before, from the Icelandic Association of Farmers (IAF).

Emission factors are taken from the IPCC Guidelines. They are presented in Table 6.3, but are likely to be overstated, as domestic animals in Iceland are generally smaller (sheep, horses) than in other European countries.

Table 6.3 Emission factors for CH₄ from manure management

	kg CH ₄ per head per year
Dairy cattle	14
Non-dairy cattle	6
Sheep	0,19
Goats	0,12
Horses	1,4
Swine	3
Poultry	0,078

In order to calculate N₂O emissions from manure management, the default IPCC methodology was used, according to the following equation.

$$E = \sum_S \left(\sum_T (N_T \cdot Nex_T \cdot MS_{T,S}) \right) \cdot EF_S$$

where E is N₂O emissions, T is the animal species index, S is the manure management system index, N_T is the livestock population, Nex_T is the annual average N excretion per head of species, $MS_{T,S}$ is the fraction of total annual excretion for each livestock species that is managed in system S and EF_S is the N₂O emission factor for system S .

The emission factors for N excretion are country-specific nitrogen factors. They are presented in Table 6.5. Emission factors for N₂O-N/N are those suggested by the IPCC Guidelines. The treatment of animal manure in different management system per animal species was estimated by the Agricultural University of Iceland. There have been some changes in the manure management practices over the time series. For example the share of liquid systems for cattle is believed to have increased from 46% in 1990 to 53% in 2005. The shares of manure management systems per animal species differ therefore for the period 1990 – 2005. The situation in 2005 is reflected in Table 6.4.

Table 6.4 Manure management systems

Manure management systems	Liquid system	Solid storage and dry lot	Pasture/range/paddock
Dairy cows	53%	13%	34%
Other cattle	53%	13%	34%
Sheep	17%	41%	42%
Goats	17%	41%	42%
Horse	0%	17%	83%
Swine	100%	0%	0%
Poultry	0%	100%	0%

6.4 Emissions from Agricultural Soils – N₂O (4D)

6.4.1 Description

Three sources of N₂O from agricultural soils are distinguished in the IPCC methodology:

- Direct emissions from agricultural soils (applying for Iceland: use of synthetic fertilizers, applied animal manure, crop residue, cultivation of soils (NE)). This is a key source in both level and trend.
- Direct soil emissions from production of animals
- N₂O emissions indirectly induced by agricultural activities (N losses by volatilization, leaching and runoff). This is key source in level.

6.4.2 Methodological issues

The methodology for calculating N₂O from agricultural soil is in accordance with the Tier 1b method.

Use of synthetic fertilizer

The direct emissions of N₂O from the use of synthetic fertilizers are calculated from data on annual usage of fertilizers and their nitrogen content, multiplied by the IPCC default emission factor. The emissions are corrected for ammonia that volatilizes during application.

Manure applied to soil

It is assumed that all animal excreta that are not deposited during grazing are used as manure. The total amount of nitrogen in manure is estimated from the number of animals and the country-specific nitrogen factors for each kind of animal, presented in Table 6.5. They are taken from Óskarsson, M. and Eggertsson, M. (1991).

Table 6.5 Nitrogen excretion factors

	kg N per head per year
Dairy cattle	60
Non-dairy cattle	33,6
Sheep	5,76
Goats	5,76*
Horses	28,8
Swine	13,3
Poultry	0,42

* N-excretion from goats are assumed to be the same as by sheep

Crop residue

This source is negligible.

Cultivation of organic histosols

This source is not estimated separately but included under emission from organic Grassland soils.

Direct soil emission from animal production

The fraction of the total amount of animal manure produced, which is deposited on pastures during grazing, is set to be 40 - 45% and differs between years. The Agricultural University of Iceland has estimated the proportion of excreted nitrogen from different types of livestock subject to different types of animal waste management systems. The level of animal manure deposited on pastures has been changing slightly due to changes in farming practices.

N losses by volatilization

Atmospheric deposition of nitrogen compounds fertilizes soils and surface waters, and enhances biogenic N₂O formation. Climate and the type of fertilizer influence the ammonia volatilization. The IPCC default values for volatilization are used (10% for synthetic fertilizers and 20% for animal manure).

N₂O from leaching and runoff

A considerable amount of nitrogen from fertilizers is lost from agricultural soils through leaching and runoff. Fertilizer nitrogen in ground water and surface waters enhances biogenic production of N₂O as the nitrogen undergoes nitrification and denitrification. The IPCC default value of 30% is used.

Emission factors

The IPCC default emission factor of 0,0125 kg N₂O-N/kg N has been used for all sources of direct N₂O emissions from agricultural soils, except for the emissions of N₂O from animal production which are calculated using the IPCC default factor of 0,02 kg N₂O-N/kg N.

The IPCC default emission factor of 0,025 kg N₂O-N/kg N is used for leaching and runoff.

Uncertainties

The preliminary estimate of quantitative uncertainty has revealed that the uncertainty of direct N₂O emissions from agricultural soils is 102% and the uncertainty of indirect emissions from Nitrogen used in agriculture is 102%.

Planned improvements

Revise country-specific N excretion factors.

Recalculations

The amount of produced barley in 2004 was revised as well as the amount synthetic fertilizer. In this submission the amount of synthetic fertilizer used in forestry and revegetation are reported under LULUCF and the corresponding amount therefore subtracted from the agriculture sector. Prior to this submission all emissions resulting from the use of synthetic fertilizer has been reported under agriculture.

7 LULUCF

7.1 Overview

This chapter provides estimates of emissions and removals from Land Use, Land-Use Change and Forestry (LULUCF) and documentation of the implementation of guidelines given in “Good Practice Guidance for Estimating and Reporting of Emissions and Removals from LULUCF” (IPCC, 2004). The LULUCF reporting is for second year according to the new CRF LULUCF tables. This section was written by the Agricultural University of Iceland.

For the first time the sectors expert (AUI) for LULUCF was able to access the UNFCCC CRF Reporter (version 3.1.) program as tool for preparing the CRF. This new working procedure has lead to some development in structure of the information reported. The main changes are; that emission of CO₂ from organic soil in Grassland is included in general CRF structure, other emissions from Grassland organic soils still needs modification of CRF; organic soil in Forest land is now reported separately involving CO₂ and N₂O emission; fertilizers used in forestry and revegetation activities are now reported separately; usage of notation keys is more complete and consistent and was reviewed for the years 1990-2004. As in 2004-2006 submission QA/QC is still only qualitative.

Land use information was obtained through same accumulation and processing of existing data as in the 2004 inventory submitted 2006. The processing of land use data is described in chapter 7.2

7.2 Data Sources

The changes from the former CRF for LUCF to the new CFR for LULUCF have complex implications regarding data needed to prepare the tables. The most important change is the need for country-wide land use information, not limited to the area previously reported on. For many land use categories division of area on organic and mineral soil is also requested. The criteria for which land use categories, an estimate for Area on organic soil is requested and which not is not always obvious.

No systematic sampling or recording has been carried out country-wide. Definitions of land use categories only exist for some categories and might in many cases be overlapping. There is a need to clearly define categories and there are cases of overlapping within categories. Information on land use is also in many cases inconsistent.

For several years Iceland’s Agricultural Research Institute, (now merged into the Agricultural University of Iceland) has been compiling a geographical database on vegetation types and grazing land condition on all farmlands. The need to control grazing pressure in accordance with ground tolerance in order to prevent erosion has been the main driver for creating this database. This work has been based on remote sensing by satellite images, existing maps of erosion and vegetation cover and various other sources.

Extensive ground truthing has resulted in a level of approximately 85% scoring in categorisation on less than 0.05 ha resolution.

Based on this investment and estimation of present and future needs for such information the Ministry of Agriculture has decided to establish a database where land use categories can to large extent be identified geographically. The work will be carried out by AUI.

The present database on farmland condition has also made it possible to produce a crude estimate of land use categories needed for CRF_LULUCF. This estimate was done for the 2004 inventory submitted 2006. The same estimate is applied to this year's submission with minor adaptations. Due to the high uncertainty of information on area of land use categories and planned improvements no attempt is made in this year inventory to estimate backward the changes in each land use category.

As explained in the following chapters the accuracy of land use information is not high but is expected to improve considerably in next years. At present information on previous land use is not available. Thus for this submission reported land use for years prior to the year 2005 is the same as for the year 2005 with the exception that changes in afforestation are reported backward as previously reported and other emission not changed. All afforestation is assumed to have been on Grassland and Revegetation and is not assumed to involve land use changes. Afforestation on organic soil is also assumed to have been on Grassland organic soil

The below description explains how information on total area of individual land use categories was obtained.

There has not been any systematic data sampling on land use in Iceland covering the whole country. Data on land use in many of the larger municipalities which have finished their municipal area planning is available. On the other hand the land use categories used in these municipal plans are often incongruous with the categories required for this inventory. These municipal plans are therefore not directly applicable to the inventory.

In preparing the inventory several databases and information sources were used to estimate coverage of the main land use categories. These data sources and their compilation are described below.

7.2.1 NYTJALAND- Icelandic farmland database: Geographical database on condition of farming land.

The Agricultural Research Institute in Iceland in cooperation with other institutes has for several years been working on a geographical database on the condition of vegetation on all farms in Iceland. The mapping is now nearly finished and approximately 60% of the country, there of 70% of the lowlands below 400 m a.s.l., has been covered. This geographical database is based on remote sensing using both *Landsat 7* and *Spot 5* images. The categorization used divides the vegetation cover into ten classes and in addition includes lakes and glaciers as classes. The definitions of categories are not the

same as required for CRF LULUCF. The classes used in NYTJALAND are listed in Table 7.1

The pixel size in this database is 15*15 m and the reference scale is 1:30000. The data was simplified by merging areas of less than 10 pixels to nearest neighbour area, thus leaving 0.225 ha as minimum mapping unit.

<i>NYTJALAND Class</i> <i>(Icelandic name in brackets)</i>	<i>Short description</i>	<i>Converted to CRF category</i>
Cultivated land (ræktað land)	All cultivated land including hayfields and cropland.	Cropland
Grassland (Graslendi)	Land with perennial grasses as dominating vegetation including drained peat-land where upland vegetation has become dominating.	Grassland
Richly vegetated heath land (Ríkt mólendi)	Heath land with rich vegetation, good grazing plants common, dwarf shrubs often dominating, and mosses common.	Grassland
Poorly vegetated heath land (Rýrt mólendi)	Heath land with lower grazing values than richly vegetated heath land often dominated by less valuable grazing plants and dwarf shrubs, mosses and lichens apparent.	Grassland
Moss land (Mosi)	Land where moss covers more than 2/3 of the total plant cover. Other vegetation includes grasses and dwarf shrubs.	Grassland
Shrubs and forest (Kjarr og skóglendi)	Land covered to more than 50% of vertical projection with trees or shrubs higher than 50 cm	Grassland
Semi-wetland- wetland upland ecotone- (Hálfdeigja)	Land where vegetation is mixture of upland and wetland species. Carex and Equisetum species common also often dwarf shrubs. Soil is generally wet but without standing water. This category includes drained land where vegetation not yet dominated by upland species.	Wetland
Wetland (votlendi)	Mires and fens. Variability of vegetation is high but mires are dominated by Carex and Equisetum species and often also shrubs.	Wetland
Partly vegetated land (hálf gróið)	Land where vegetation cover is from 20-50% generally infertile areas often on gravel soil. Both areas where the vegetation is retreating and in progress can be included in this class	Other land
Sparsely vegetated land (Litt gróið)	Many types of surfaces are included in this class with the common criteria of less than 20 % vegetation cover in vertical projection.	Other land
Lakes and rivers	Lakes and rivers	Wetland
Glaciers	Glaciers	Other land

Table 7.1 Land cover classes of NYTJALAND database and their transformation to CRF land use classes

7.2.2 Vegetation maps

To compensate for areas not covered by NYTJALAND database simplified vegetation, maps from Iceland Museum of Natural History were used. The scale of these maps is 1:500000 and vegetation is categorised to 5 classes plus water and glaciers. These maps

are of considerably less resolution and accuracy than the NYTJALAND database and their classes are listed in Table 7.2

<i>Vegetation class</i>	<i>Converted to CRF land use category</i>
Grassland -Heath land	Grassland
Wetland	Wetland
Shrubs and forests	Grassland
Moss land	Grassland
Sparsely vegetated land	Other land
Lakes and rivers	Wetland
Glaciers	Other land

Table 7.2. Categories of 1:500000 vegetation maps from Iceland Museum of Natural History and the conversion of those to CRF_LULUCF categories

7.2.3 Cities, towns and villages

Data on area covered by cities, towns and villages were obtained from IS50 database of National Land Survey of Iceland available in 1:50000 scale. This area was converted to CRF_LULUCF category settlement.

7.2.4 Unified dataset




Figure 7.1 First approach to land use map of Iceland

The above described geographical databases were merged into one dataset, providing a first approach to a geographically identifiable land use database covering four of the Good Practice Guidance for Land Use, Land-Use Change and Forestry (GPG - LULUCF)(IPCC 2003) suggested land use categories, i.e. grassland, wetland, settlement and other land. Two subcategories, lakes and rivers and glaciers are included. A map of Iceland showing this first approach is in Figure 7.1

Besides lacking two of the main categories suggested in the GPG-LULUCF, there are many uncertainties regarding the definition of categories. Further refinement of this database could therefore result in large areas to be transposed from one category to another. To better meet the requirements of the inventory many of the categories need to be further subdivided, most importantly into managed and unmanaged areas.

To improve the quality of the information provided several modifications of the land use categorization were made. Some of these changes are geographically identifiable but others not.

The modifications that were made are:

- a) Subdivision of class “Partly vegetated land” of the NYTJALAND database and reclassification of some of these subcategories. Part of this land cover

class does better fit to the category ‘grassland’ than ‘other land’. Some of the land in this category has been reclassified as such because it is eroded to a certain stage. All erosion in Iceland was mapped in the years 1993-1995, in a large project which was carried out in cooperation between the Iceland Agricultural Research Institute and the Iceland Soil Conservation (Arnalds et al. 2001) By comparing the category ‘partly vegetated land’ and all erosion classes connected to vegetated land the areas inside this class where vegetation is retreating but still should be classified as grassland can be identified. These areas were added to the ‘grassland’ category and removed from the category ‘other land’. In areas covered by the NYTJALAND database these transformations are geographically identifiable, in other areas not.

- b) Lava fields from historic time (from year 875 AC) covered with mosses was moved from category ‘grassland’ to ‘other land’. By comparing the geological maps from the Icelandic Museum of Natural History to land referred to as ‘moss land’ from both the NYTJALAND database as well as vegetation maps, these areas can be identified geographically.
- c) Drained wetlands are in this inventory classified as grassland although LULUCF GPG would allow it to be also classified as cropland or managed wetland. Part of the drained wetlands is already included under the grassland category but some areas are under the categories ‘semi-wetland’ or ‘wetland’. To correct for this, all ditches with a 100m buffer zone, from the map layer “lakes and rivers” of the National Land Survey of Iceland database were added to the ‘grassland’ category and subsequently removed from the ‘wetland’ category. As the ditches layer of National Land Survey of Iceland database is incomplete only covering 10.000 km length of ditches of which the total length is estimated 32.700 km (Óskarsson 1998) the results of the re-estimate was extrapolated to these 32.700 km of ditches, this additional area estimate was added to the Grassland category and subsequently removed from the ‘wetland’ category. This improved categorization is only partly geographically identifiable.
- d) Land for roads is not included in the Settlement as it is in IS50 database of National Land Survey of Iceland used in compiling the unified dataset. To correct for that a layer of all roads with a 15 m buffer zone was added to the unified dataset and overlaps to each category in that dataset were added to the Settlement category and subsequently removed from the relevant category.
- e) The category Forest land is not in the unified dataset. Information on total area of forests both plantations and native birch shrub- and woodland are available from Icelandic Forest Research. All tree plantations fulfil the country specific definition of forest but only 37% of the native shrub- and woodland does so. Total area of defined forest was set as total area of Forest land category and subsequently removed from the grassland category. The area estimate for Forest land is explained in chapter 7.3 .
- f) The Cropland category in the NYTJALAND database is incomplete and does only cover part of the country. This information has to be screen digitized to the database and that work is not finished. The area already mapped is

subsequently removed from the relevant land use category. To compensate for the incompleteness in mapping the total area of cropland is corrected according to information from Iceland Agricultural Statistics and the difference in area removed from the Grassland category.

- g) Revegetated land is not reported in this year submission as a land use category, although information on some areas is available and calculation is based on the size of land subjected to revegetation. It can be argued that revegetated land should be categorized separately in order to increase transparency and completeness of the land use recording. Thus classification of land subjected to revegetation is still to be decided.
- h) The area of drained peat land has been estimated (Óskarsson 1998), and used in the grassland subcategory, 'drained organic soil'. The areas drained are only partly geographically identifiable. The estimate based on maps from National Land Survey of Iceland is considered less accurate.
- i) The area of hydro-reservoirs used for electrical power plants was obtained from the National Power Company (Landsvirkjun). This land use category is a subcategory of 'wetland'.

<i>Land use category</i>	<i>Area (kha)</i>	<i>Geographically identifiable</i>
<i>Subcategory</i>	<i>main category</i>	<i>Subcategory</i>
Grassland¹	3843.9	Majority
<i>Drained organic soil</i>		446.5 Partly
<i>Grassland on mineral soil</i>		3397.4 Partly
Wetland	877	
<i>Peatland</i>		668 Partly
<i>Reservoirs</i>		25 Identifiable with little effort
<i>Rivers and lakes</i>		184 Identifiable
Settlements	68	Identifiable
Cropland	129	Partly
Forest¹	52,4	Majority
<i>Native birch woodlands³</i>		25 Partly
<i>Afforestation</i>		27.4 Partly
<i>Thereof Area afforested on organic soil</i>		3.53 Partly
Other land	5310	
<i>Glaciers</i>		1324 Identifiable
<i>Other(sparsely or not vegetated)</i>		3986 Identifiable

Table 7.3 Land use classification used in GHG inventory 2005 submitted 2007.

³ In NIR report for the 2004 inventory Area of "Native birch forests" is reported as 44 kha and but in CRF 2004 and 2005 the area reported was 25 kha. Consistent with area reported in CRF 25 kha is reported here, although 44 kha is considered more accurate number.

The unified dataset does provide two subcategories in addition to the main categories, i.e. lakes and rivers as ‘wetland’ subcategories and glaciers as a subcategory of ‘other land’. The resulting estimate of land use categories is listed in Table 7.3.

7.2.5 Land use changes

The only reported land use changes involves afforestation. All afforestation is assumed to be on land classified as grassland. Other land use changes are at present not reported due to lack of data.

7.2.6 Land use definitions and the classification system and their correspondence to the LULUCF categories

Definitions of land use categories have not been elaborated. The present status of land use information in Iceland is as described above very fractional and in many ways inconsistent between data sources. The definitions on land use categories used in this inventory are those presented in Table 7.1 with the modifications described. The elaboration of definitions for land use categories is a vital part of building the geographical land use database described in chapter 7.2.8

7.2.7 Uncertainties QA/QC

The NYTJALAND database, which is the main source of land use information supporting the land use classification used for the inventory, has the scoring accuracy of 85% on its categorization. The modifications and additions of other data needed for the complete coverage of all land has most likely decreased that accuracy to some level. The amount of that increase in uncertainty has not been quantified.

7.2.8 Planned improvements regarding land use identification

Land use registration in Iceland is very fragmentary and not based on long term traditions as in many other European countries. Construction of geographically identifiable land use database covering the whole country will start the year 2007. The Government of Iceland has ensured financing of the project. This work is expected to gradually increase the quality of land use information as well as provide data for estimates of ongoing land use changes and to some extent past changes.

Iceland has agreed to participate in CORINE land cover mapping of Europe. The proposal is at present in the application phase. The proposal involves full participation in CLC 2006 mapping and CLC 2000/2006 changes the latter will be integrated to give CLC 2000.

The Icelandic mapping effort will be based on the NYTJALAND –database and other sources available. The planned construction of Icelandic land use database and CORINE mapping effort will be carried out in close cooperation between the relevant institutions (Icelandic Agricultural University and National Land Survey of Iceland).

These two projects are expected to improve the quality of land use information considerably and provide backward estimates of land use changes.

7.2.9 Completeness and method

Based on the above described accumulation of land use data and emission factors the emission by source and removal by sinks were calculated.

Source/sink	Area kha	Method	EF	CO ₂ Gg Emission/ Removal (-)
Forest remaining forest	52,4			
- Living biomass		T2	CS	-120,56
- Dead organic matter		NE (T1)		
- Soils		NE,IE		
Mineral Soils		NE		
Organic Soils	3,53			2,08
Land converted to forest	1,8			
- Living biomass		T2	CS	-7,92
- Dead organic matter		NE		
- Soils		NE,IE		
Mineral Soils	1,57			
Organic Soils	0,23			0,14
Cropland remaining cropland	129			
- Living biomass		NO		
- Dead organic matter		NE		
- Soils		NE		
Mineral Soils		NE		
Organic Soils		IE		
- Lime application		T2	D,CS	3,46
Land converted to cropland	NE	NE		
Grassland remaining grassland	3843,9			
- Living biomass		NE		
- Dead organic matter		NA		
- Soils				
Mineral Soil		NE		
Organic Soil	446,47	T1	D,CS	1800,74
- Lime application	NO			
Land converted to grassland	NE			
Wetlands remaining wetland				
Lakes and rivers	183,9	NE		
Peatland	667,59	NE		
Reservoirs	25			
- Living biomass		NE		
- Dead organic matter		NE		
- Soils (flooded)		T1	D	141,42
Land converted to wetland	NE			
Settlements remaining settlements	68,45	NE		
Land converted to settlements	NE			
Other land remaining other land	5310,06			
Land converted to other land	NE			
Biomass burning	NE			
Other				
Revegetation	NA	T2	CS	-533,24
Harvested wood		NE		

Table 7.4 Summary of method and emission factors applied on CO₂ emission calculation.

EF = emission factor, D = default (IPCC), CS = country specific, NA = not applicable, NE= not estimated, NO = not occurring, IE=included elsewhere, T1 = Tier 1, T2 = Tier 2 and T3 = Tier 3,

Source/sink	Area kha	Method	CH ₄ EF	Gg Emission/ Removal (-)	CO ₂ equivalent Gg
CH₄ emission from drainage of soil					
<i>Forest remaining forest</i>					
Organic Soil	3,53	NA			
Mineral Soil	NE				
<i>Grassland organic soil</i>					
Organic Soils	NA				
Mineral Soils	NO				
<i>Other</i>					
Revegetation					
Organic Soils	NO				
Mineral Soils	NO				
Biomass burning	NE				
Other CH₄ emission					
Wetland remaining wetland					
Reservoirs	25	T1	D	2,34	49,14
N₂O					
5(I) Direct N₂O emission from N fertilization					
Forest remaining forest	52,4	T1	D	0,00	0,01
Land converted to forest	1,8	T1	D	0,00	0,10
Grassland organic soil		NE			
Revegetation	NA	T1		0,02	4,95
5(II) N₂O emission from drainage of soil					
<i>Forest remaining forest</i>					
Organic Soil	3,53	T1	D	0,00	1,03
Mineral Soil		NE			
<i>Land converted to wetland</i>	NE				
<i>Grassland organic soil</i>					
Organic Soils	446,47	T1	D	1,26	391,49
Mineral Soils	NE				
Revegetation	NO				
Biomass burning	NE				
Other N₂O emission					
Wetland remaining wetland					
Reservoirs	25	T1	D	0,07	21,7

Table 7.5 Summary of method and emission factors applied on CH₄ and N₂O emission calculations.

EF = emission factor, D = default (IPCC), CS = country specific, NA = not applicable, NE= not estimated, NO = not occurring, IE=included elsewhere, T1 = Tier 1, T2 = Tier 2 and T3 = Tier 3,

Summary of method and emission factors used is provided in Table 7.4 and Table 7.5

7.2.10 Key sources/sink and key areas

Of the sources/sinks calculated, four were recognized as LULUCF level key source with regard to CO₂ equivalents (Table 7.6). Non-estimated categories can not be excluded as a potential level key source.

Source/sink	Direct greenhouse gas Gg Emission/ Removal (-)	Absolute value Gg	CO ₂ equivalent		Key source
			Level %	cumulative level %	
Grassland remaining grassland - Organic Soil CO ₂	1800,74	1800,74	58,5	58,5	x
Revegetation CO ₂	-533,24	533,24	17,3	75,8	x
Grassland Organic Soils-N ₂ O emission from drainage of soil	1,26	391,49	12,7	88,5	x
Reservoirs CO ₂	141,42	141,42	4,6	93,1	x
Forest remaining forest- Living biomass CO ₂	-120,56	120,56	3,9	97,1	
Reservoirs CH ₄	2,34	49,14	1,6	98,7	
Reservoirs N ₂ O	0,07	21,7	0,7	99,4	
Land converted to forest - Living biomass CO ₂	-7,92	7,92	0,3	99,6	
Revegetation Direct N ₂ O emission from N fertilization	0,02	4,95	0,2	99,8	
Cropland - Lime application CO ₂	3,46	3,46	0,1	99,9	
Forest remaining forest Organic Soils CO ₂	2,08	2,08	0,1	100,0	
Forest remaining forest-Organic Soil N ₂ O emission from drainage of soil	0,00	1,03	0,0	100,0	
Land converted to forest- Organic Soils CO ₂	0,14	0,14	0,0	100,0	
Land converted to forest Direct N ₂ O emission from N fertilization	0,00	0,10	0,0	100,0	
Forest remaining forest- Direct N ₂ O emission from N fertilization	0,00	0,01	0,0	100,0	
Total		3077,98			

Table 7.6 LULUCF level key source assessment of land use categories, for which emissions/removals were calculated

Trend key source assessment was not done as land use change data is not available for most of the categories. Considering the present status of land use information the key land use categories were assessed. Two assessments were performed on the land use categories as reported (Table 7.7) and on applicable land use categories adding revegetation as land use category and excluding other land as by definition unmanaged and emission/removal calculation not applicable. Considering reported area four land use categories are recognized as key categories in relation to area i.e.; other land, grasslands total area, peatland and grassland on organic soil.

Land use category as reported	Area kha	Area Level %	Cumilative area level	Key land use category
Other land remaining other land	5310,06	49,5	49,5	x
Grassland remaining grassland-total area	3843,90	35,8	85,3	x
Peatland	667,59	6,2	91,5	x
Grassland - Organic Soil	446,47	4,2	95,7	x
Lakes and rivers	183,90	1,7	97,4	
Cropland remaining cropland	129,00	1,2	98,6	
Settlements remaining settlements	68,45	0,6	99,2	
Forest remaining forest total area	52,40	0,5	99,7	
Reservoirs	25,00	0,2	99,9	
Forest remaining forest- Plantations- Organic Soils	3,53	0,0	100,0	
Land converted to forest- total area	1,80	0,0	100,0	
Land converted to forest-Mineral Soils	1,57	0,0	100,0	
Land converted to forest-Organic Soils	0,23	0,0	100,0	
Total *	10733,90			

Table 7.7 LULUCF area level assessment of land use categories where area identified.

*Total area exceeds country total area due to land use sub categories reported.

Applicable land use categories	Area	Area Level %	Cumilative area level	Key land use category
Grassland remaining grassland-total area	3843,9	68,4	68,4	x
Peatland	667,59	11,9	80,3	x
Grassland - Organic Soil	446,47	7,9	88,3	x
Revegetation	193,91	3,5	91,7	x
Lakes and rivers	183,90	3,3	95,0	x
Cropland remaining cropland	129,00	2,3	97,3	
Settlements remaining settlements	68,45	1,2	98,5	
Forest remaining forest total area	52,40	0,9	99,4	
Reservoirs	25,00	0,4	99,9	
Forest remaining forest- Plantations- Organic Soils	3,53	0,1	99,9	
Land converted to forest- total area	1,80	0,0	100,0	
Land converted to forest-Mineral Soils	1,57	0,0	100,0	
Land converted to forest-Organic Soils	0,23	0,0	100,0	
Total	5617,75			

Table 7.8 LULUCF area level assessments of land use categories considered relevant as potential source/ sinks and where area was identified

Considering only applicable categories, (Table 7.8) two land use categories i.e. lakes and rivers and revegetation are assessed as key areas.

7.3 Forest land

Current calculation of C-stock changes is simple and deficient. The whole process of forest inventories and aggregation of forestry data is under total revision as described below. Total recalculations of the forest sector are to be expected in the nearest future.

7.3.1 Carbon stock changes (5A)

7.3.1.1 Carbon stock changes in living biomass

In accordance to the GPG arising from the Kyoto Protocol a country-specific definition of forest has been adopted. The minimal crown cover of forest is 10%, the minimal height 2 m, minimal area 0.5 ha and minimal width 20 m. This definition is also used for the national inventory. Further description of forest definition is to be found in methodological report of carbon accounting of forests (Snorrason and Kjartansson, 2004)). All forests are defined as managed.

The total area of native woodlands in Iceland has been inventoried twice in the 20th century, in the periods 1972-1975 and 1987-1991, resulting in estimates of 125 and 118 kha respectively. Maps and data sampled in the inventories have newly been put to GIS. New SPOT images helped refine the mapping and now the total area of the native birch woodlands is estimated to 120 kha. The Icelandic Forest Research estimated in 2006 that 25 kha of the native birch woodland is higher than 2m. New estimate of Icelandic Forest Research is that 37% or 44 kha of this area is forest meets the forest criteria decided for KP-reporting. In the NIR for 2004 the 44 kha was used but not in the CRF submitted 2006 where 25 kha was reported for native birch forests. This inconsistency was caused by insecure data transfer procedures. In this years submission the native birch forests are reported as 25 kha both in CRF and NIR recognising uncertainty in area and maintaining consistency in CRF.

Total woody C-stock was from these data estimated at 1300 kt C with average of 11 t C ha⁻¹ in 1990. The two inventories are not comparable in methodology and can not be directly compared to show changes in area or woody stock during this period. (Sigurðsson and Snorrason 2000). The C stock of the native birch woodlands is assumed to remain constant with no changes reported.

Afforestation and reforestation started in Iceland 1899. Before 1970 planting of forest was mostly done in natural woodlands. The total area of plantations from 1970 to 1989 has been estimated to be 3 kha and of older plantations 3,6 kha. The annual changes of the woody biomass of these plantations are estimated the same as in plantations from 1990 or later.

Most afforestation areas in Iceland are relatively young and clear cutting has not started. The only exceptions of deforestation are when natural woodland and plantations have to give way for road or house building. A preliminary investigation of deforestation has shown that it is very rare and at a small scale. Neither the clear cuttings nor the thinning of managed forests is presently systematically recorded.

Current C-stock change calculation of living biomass is described in chapter 7.3.3

7.3.1.2 Net carbon stock changes in dead organic matter

No attempt is made to estimate changes in dead organic matter due to lack of data. Tier 1 (GPG for LULUCF) default assumes no changes in dead wood or litter. Changes in dead organic matter are in GPG for LULUCF connected with forest management and due to the young age of most Icelandic plantations, this category is not considered important.

7.3.1.3 Net carbon stock change in soils

The Icelandic Forest Research has estimated that 13% of planted forest since 1990 were planted on wetland or drained wetlands. Assuming this ratio also applies to plantations before 1990, and these areas represent organic soils in forests, the area of organic soils in forest land is reported for all years since 1990. The area estimated for organic soils in forest land was subsequently subtracted from the aggregated estimate for drained organic soils previously reported under Grassland. Net carbon stock changes in soils are only estimated for the organic soils. Net carbon stock changes in mineral soils are not estimated. The native forest and the remaining afforested areas are mostly situated on mineral soils which can be highly variable regarding carbon content. At present the Icelandic Forest Research states that soil carbon stock in forest land on mineral soil is not decreasing (Snorrason 2003b).

7.3.2 Other emissions (5(I), 5 (II), 5(III))

For the first time direct N₂O emission from use of N fertilisers is reported separately for forest remaining forest and land converged to forest land.

N₂O emission from drainage of organic soils is also reported separately for forest land.

Ploughing is sometimes part of the preparation of new plantations and might therefore cause N₂O emissions comparable to land conversion to cropland. Although possible, no data is available on amount of afforested land ploughed or emissions caused by such activity reported under 'other' in Table 5(III).

7.3.3 Land converted to forest land.

At present state of information on land use changes all new plantations are reported to involve conversion of grassland to forest land. The average ratio of plantation area on organic soil estimated from information from the Icelandic Forest Research for all plantations from 1990 is also assumed to apply for land converted the inventory year.

The Icelandic Forest Research records in the new forest inventory project previous land use of planted area involved. Information is thus available on more detailed land use conversion than reported (Table 7.9). Due to high uncertainty in area of other land use categories this information is not considered applicable as only source of land use changes.

Land category	% of afforested area
Sparsely vegetated upland (<30% vegetation cover)	16%
Other upland	69%
Wetland	3%
Drained wetland	10%
Cropland	2%
Total	100%

Table 7.9 Former land categories of afforested areas as estimated by the Icelandic Forestry Research

7.3.4 Methodological issues

The area of new plantations is estimated from number of seedlings delivered from plant breeding stations. The estimation of afforested area is based on the following assumptions: On average planting density was 4000 seedlings ha⁻¹, 25% of afforested area is lost to various reasons (Sigurðsson and Snorrason 2000). Since 1990 the seedling density has decreased and new comparison of number of seedling and area of land did show that the number had dropped down to 2350 seedlings per ha (Snorrason and Kjartansson. 2004). First results of the ongoing new national forest inventory (Snorrason 2006 personal communication) are consistent with the total area of forests estimated from number of seedlings planted.

7.3.5 Emission/removal factors

Tier 2 is used to estimate increment in carbon stock change in living biomass. The annual C removal factor used in the inventory (1.2 t C ha⁻¹) is a precautionary estimate of data from Icelandic Forest Research, (Snorrason 2003a) including both surface biomass and below ground biomass of coarse roots.

Tier 1 and default EF = 0,16 [t C ha⁻¹ yr⁻¹] (LULUCF GPG Table 3.2.3.) is used to estimate net carbon stock change in forest organic soils. For direct N₂O emission from N fertilization and N₂O emission from drained organic soils, tier 1 and default EF=1,25% [kg N₂O-N/kg N input] (GPG2000) and EF=0,6 [kg N₂O-N ha⁻¹yr⁻¹] (LULUCF GPG Table 3.a.2.1.) were used respectively.

7.3.6 Uncertainties QA/QC

Using the average annual C removal factor overestimates the removal of C by young plantations. The C uptake factor is based on measurements where the biomass of forest plantations of known age was measured. These measurements have resulted in highly variable results ranging from 0.9-3.8 t C ha⁻¹ (Jónsson and Óskarsson 1996) (Snorrason 2003a).

How well the used factor represents the actual plantations is thus a source of error which acts both on age of plantation and spatial variability. The area estimate is based on indirect data sources calibrated to field data. (Snorrason and Kjartansson. 2004).

The new forest inventory project, established by the Icelandic Forest Research, yielded its first unpublished results which were consistent with previous estimates of total forest area. The annual removal factor is based on field measurements (Tier 2). The ongoing improvement in forest inventory will improve the control and verification options.

7.3.7 Recalculations

Area of organic soils in forest land is estimated for all years from 1990 and subsequently emission of CO₂ and N₂O from these soils is calculated for each year changing the previously reported removals.

7.3.8 Planned improvements regarding Forest land

The methodology for the new inventory is based on systematic sampling consisting of a total amount of nearly 1.000 permanent plots. One fifth of the plots are measured each year and remeasured at 5 year intervals. The sample will be used to estimate both area classes and C-stock changes over time (Snorrason and Kjartansson. 2004). Preparation of this work started in 2001 and the measurement on field plot started in 2005. One can therefore expect gradually improved estimates of carbon stock and carbon stock changes in both managed and unmanaged woodlands in Iceland. Improvements in forest inventories will also improve uncertainty estimates both on area and stock changes. Soil sampling is included as part of the forestry inventory program and estimates of changes in soil carbon stock is expected in future reporting.

Major recalculations of the forest sector are therefore to be expected in the nearest future.

7.4 Cropland

Cropland in Iceland consists mainly of cultivated hayfields, many of which on drained organic soil. A small but increasing part is used for cultivation of barley. Cultivation of potatoes and vegetables also takes place. No information is available on emission/removal regarding different cultivation types and subdivision of areas is not attempted. Cropland is identified as a key area in applicable land use categories.

7.4.1 Carbon stock changes (5B)

7.4.1.1 Carbon stock changes in living biomass

As no perennial woody crops are cultivated in Iceland, no biomass changes need to be reported.

7.4.1.2 Net carbon stock changes in dead organic matter

The GPG for LULUCF does provide default parameters to estimate carbon stock changes in dead organic matter. No data is available to estimate the possible changes in dead organic matter in remaining cropland. The majority of land classified as cropland in Iceland is hayfields with perennial grasses only ploughed or harrowed at decade intervals. A turf layer is formed and depending on the soil horizon definition it can be considered as dead organic matter. This is therefore recognised as a possible sink/source although no data is available.

7.4.1.3 Net carbon stock change in soils

Net carbon stock changes in mineral cropland soil are not estimated. No data available and no default relative stock change factors recognised as applicable to perennial hayfield.

Emissions from organic cropland soils are reported as an aggregate number along with emission from drained grassland organic soils. Data for partitioning of drained organic soils between cropland and grassland is not available.

7.4.2 Other emissions (5(I), 5 (II), 5(III), 5(IV))

Direct N₂O emission from use of N fertilisers is included under emissions from agricultural soils.

N₂O emissions from drainage of organic soils are reported as an aggregated number under emissions from grassland. Separation of drained organic soil to land use categories is not possible on basis of present land use data.

Carbon emissions from agricultural lime application are estimated. Information on lime application are obtained from distributors. Numbers reported hereto have included lime application in the form of shellsand which contains 90 % CaCO₃, dolomite and limestone. Many of the fertilizers imported include limestone or other calcifying agents as additive. Although the ratio of calcifying materials is low in these fertilizers the amount of fertilizers applied make this source relatively large. This source is reported for the first time and explains the 37% increase in CO₂ emission and 31% increase in applied amount. Numbers only available on national level and all of it is assumed to be applied on cropland.

7.4.3 Land converted to cropland

As no data is available on area of land converted to cropland, its emissions are not estimated

7.4.4 Emission factors

The only cropland emission reported is CO₂ emission due to liming. Emissions are calculated by conversion of carbonated carbon to CO₂.

7.4.5 Uncertainty QA/QC

The only reported emission/removal under 'cropland' is emission due to agricultural liming. No quality control or assurance has been undertaken regarding the submitted amounts. The largest uncertainty in cropland emissions/removals is probably the area estimate. Cropland as reported by Iceland Agricultural Statistics is mostly hayfields with perennial grasses where only a small part is used for annual crops. Elaboration of definitions on cropland could shift a large section of 'cropland' to the 'grassland' category. Due to the lack of subdivision of cropland to cultivation categories and soil type, the emissions/removals of cropland are included in aggregated numbers in other categories resulting in relatively substantial uncertainty within the category. The quantity of uncertainty for cropland emissions/removals is not estimated.

7.4.6 Planned improvements regarding cropland

Mapping of cropland and subdivision with regard to soil types and cultivation will improve along with the quality of other land use information. One of the objectives by constructing the land use database is to identify relevant subcategories regarding soil types and management.

7.5 Grassland

Grassland is the second largest land use category identified by present land use mapping described above. Only the land use category “other land” which includes glaciers is estimated larger. If glaciers were excluded from ‘other land’ the grassland category would be the largest land use category in Iceland. The ‘grassland’ category is very diverse with regards to vegetation, soil type, erosion and management. Included are heathlands with dwarf shrubs, small bushes, grasses and mosses in variable combinations, fertile grasslands, all in highly variable condition regarding erosion. Also included in the category are large areas of drained peatlands. This category is likely to overlap with wetland, cropland and other land depending on definitions of each category. There are large areas suffering from severe erosion where vegetation cover is severely damaged but the Andic soil still has high amounts of carbon. Recent research results in Iceland indicate that the carbon budget of such areas might be positive, resulting in CO₂ emission to the atmosphere. The vegetation cover in many other grassland areas in Iceland is at present increasing both in vigour and continuity (Náttúrufræðistofnun Íslands 2005) (*Icelandic Institute of Natural History Annual Report 2005*). In these areas, the annual carbon budget might be negative at present, meaning C is removed from the atmosphere. The size of these subcategories of grassland is not known at present. Whether these changes in vegetation are due to changes in climate, management or a combination of both is not clear.

Numbers are only available on the total area of drained organic soil and not for divisions between land use categories, with the exception of forest plantations. All other drained organic soil is at present assumed to be included under ‘grassland’ as subcategory.

Both the ‘grassland’ total area and area of organic soil under ‘grasslands’ are identified as key areas considering both reported area and applicable area.

7.5.1 Carbon stock changes (5C)

7.5.1.1 Carbon stock changes in living biomass

No information is available on overall changes in living biomass although it is known that changes are occurring. Division of grassland to subcategories is not possible with the present status of geographical information. Changes in carbon stock in living biomass are therefore not estimated, as is consistent with the Tier 1 methodology for the category.

7.5.1.2 Net carbon stock changes in dead organic matter

Changes in dead organic matter are not requested by the GPG for LULUCF and no information is available for this stock.

7.5.1.3 Net carbon stock change in soils

Changes in carbon stock in mineral grassland soils are not estimated due to lack of data. Tier 1 methodology gives by default no changes if land use, management and input (F_{LU} , F_{MG} , F_I) are unchanged over a period. This subcategory is therefore not estimated.

Extensive drainage of wetland has taken place in Iceland mostly in the period 1940-1985. Information on the areas drained is neither geographically identifiable nor can be subdivided to different soil types at the present time.

Although data separation is not possible at present, Iceland's drained soils include three soil types; Histosol, Histic Andosol and Gleyic Andosol. The two organic soil types are Histic Andosol and Histosol. Although Gleyic Andosol is not classified as organic, it is included here. Organic soils in Iceland generally are with relatively low C/N ratio and are therefore considered nutrient rich.

The area of grassland on organic soils is reported as an aggregate number which includes all drained soils except area of organic soil in forest land. The reported area is the estimate of drained areas as described in chapter 7.2.4 subtracted by the area of afforested organic soils as estimated on base of information from Icelandic Forestry Research. Some of area may eventually be included under cropland or even managed wetland, depending on the definitions of the land use categories still to be elaborated.

This factor is identified as level key source factor of LULUCF and as a key area both regarding reported area and applicable area.

7.5.2 Other emissions (5(IV))

All CO₂ emission due to liming is reported as aggregate number under land use category "cropland".

7.5.3 Emission factors

Emissions of CO₂ from drained organic soil are calculated according to Tier 1 methodology and uses the emission factor $EF = 1,1$ [t C ha⁻¹ yr⁻¹] (GPG Table 3A.3.2) considering high N content and aggregation of areas from different categories. Recent research on upland CO₂ indicate a higher emission factor than used.

7.5.4 Land converted to grassland.

Revegetation in Iceland often involves converting previously unvegetated area ('other land') into grassland. In those instances it could be reported as such. Revegetation is reported in this submission as separate activity not involving land use changes. Reporting revegetation under land converted to grassland would therefore involve double accounting. As no data except for revegetation is available on area of land converted to grassland, its emissions are not estimated

7.5.5 Uncertainty QA/QC

Uncertainty in reported emission from this category is supposed to be large. Several components contribute to this uncertainty. The CO₂ emission from mineral soils, which is not estimated, is potentially a large source considering the severe erosion in large areas. Counteracting these emissions might be removal in areas where vegetation is recovering from previous degradation.

Uncertainty in reported emissions from drained soil is also substantial. The total area drained is based on the estimate of drainage effectiveness and the total length of ditches (Óskarsson 1998). Both numbers have a large uncertainty range. The total amount of ditches includes only ditches dug prior to 1993 for agricultural purposes subsidized by the government. Newer ditches and ditches excavated for other purposes such as road building are not included. Effectiveness estimates range from 7.3 km/km² (Óskarsson 1998) to 20 km/km² (Geirsson 1975). Digitized mapping of ditches in Iceland has started at the Agricultural University of Iceland. Preliminary data from those maps indicate effectiveness around 8.4 km/km² (unpublished data from Agricultural University of Iceland), which is around 15% larger number than effectiveness used.

Elaboration on definitions of land use categories might shift large areas into or out of grassland category.

Emission factors for both CO₂ and N₂O are stated with large uncertainty range in GPG for LULUCF tables.

7.5.6 Planned improvements regarding grassland

Due to the potential importance of emissions, and removal in case of e.g. changed management, from/to mineral grassland soils, it is recognised as high priority issue to move up to a higher tier level regarding estimate of carbon stock changes in soil for the subcategory.

As severely degraded soils are widespread in Iceland as a result of extensive erosion over a long period of time, the changes in carbon stocks of mineral soil are a potentially large source of carbon. The importance of this source is emphasized, since mineral grassland soil in Iceland is almost always Andosol with high C content (Arnalds and Gretarsson 2001).

Emissions of both CO₂ and N₂O from grassland organic soils are identified as key sources for LULUCF and improving the resolution in recording land use, soil types and refinement of emission factors is highly important. Improvements in ascertaining the extent of drained organic soils within different land use categories will be a priority of the new land use database to be established.

How to incorporate revegetation activities in the geographical land use database and ensure consistent accounting to UNFCCC and KP remains to be solved. The recording of land converted to grassland is closely connected to that question. Solving this is a priority task in construction of the new geographical land use database.

7.6 Wetland

7.6.1 Carbon stock changes (5D)

Wetland areas are divided to three subcategories, lakes and rivers, peatlands and reservoirs. Emissions are only estimated for reservoirs as other areas are either considered unmanaged or methodology not been developed. Peat extraction is not occurring in Iceland for fuel combustion. The only peat extraction occurring is due to land converted to settlement. This source is not estimated in the inventory. There is no data available on the amount extracted and emission from that source is not requested either under land converted to settlement. Some of the lowland peatlands in Iceland could fall under managed land due to livestock grazing and should be reported as such, no information is at present available on area of grazed peatlands. Drained peatlands are reported as grassland on organic soils. All lakes and rivers are considered unmanaged. Emissions of CO₂ from reservoirs are identified as level key source of LULUCF.

Area estimates for reservoirs were obtained from the National Power Company (Landsvirkjun).

As for other categories no information is available on land converted to wetlands.

The subcategories ‘lakes and rivers’ and ‘peatland’ are identified as key areas both with regards to reported and applicable land use categories.

7.6.2 Other emissions (5)

Emissions of CH₄ and N₂O from reservoirs are reported. Emission of CH₄ and N₂O from drained peatlands is reported under grassland organic soil.

7.6.3 Emission factors

CO₂ emission from reservoirs is estimated according to the Tier 1 method using eq. 3a.3.8 in GPG for LULUCF and the default emission factor EF=15,5 [kg ha⁻¹d⁻¹] (GPG LULUCF Table 3A.3.5) for boreal, wet climate zone.

Emissions of CH₄ and N₂O from reservoirs is estimated according to Tier 1 using eq. 3a.3.9 and 3a.3.10 and respectively in GPG for LULUCF and the default emission factors E(CH₄)_{diff} = 0,11 [kg CH₄ ha⁻¹d⁻¹] and E(CH₄)_{bubb} = 0,29 [kg CH₄ ha⁻¹d⁻¹] for ice-free period and E(CH₄)_{diff+bubb} = 0,05 [kg CH₄ ha⁻¹d⁻¹] for icecover period and E(N₂O)_{diff} = 0,008 from Table 3A.3.5 under ‘boreal, wet climate zone’. Spillway emission of CH₄ is not included as no information on [CH₄]_{spillway} is available.

7.6.4 Land converted to wetland

Two sources of land converted to wetland are recognized; by flooding due to construction of new hydropower reservoirs and through reclamation of wetland to counteract damaged wetlands due to road building or as recreational area connected to tourism. No information is at present available on area converted.

7.6.5 Uncertainty QA/QC

Main uncertainty is associated with emission factors as presented in GPG for LULUCF.

7.6.6 Planned improvements regarding Wetland

Ongoing research projects on reservoir greenhouse gas emission is expected to provide country specific emission factors enabling the movement to Tier 2 or 3 methods of estimation for all gasses.

The new geographical land use database is expected to improve area data on wetland and its subcategories.

7.7 Settlements

7.7.1 Carbon stock changes (5E)

The GPG for LULUCF focus only on stock changes in living tree biomass for this category. No data is available for neither this stock nor other possible sources so emissions are not estimated.

7.7.2 Other emissions (5)

Burning of biomass in open areas within the category settlement does take place (see chapter 7.10). No other sources of CH₄ or N₂O have been recognized.

7.7.3 Land converted to settlement

At present no official country wise compilation of land converted to settlement is done. Previous land use categories are generally not recorded in municipal area planning.

7.7.4 Planned improvements regarding Settlement

The new geographical land use database is expected to improve area data on previous land use of land converted to settlement. Compilation of municipal area mapping is part of the CORINE mapping effort and will be carried out by National Land Survey of Iceland. Therefore estimate of area of land use category ‘settlement remaining settlement’ and ‘land converted to settlement’ are expected to improve. In the new geographical database data for the land use category ‘settlement’ will be provided by National Land Survey of Iceland.

7.8 Other land (5, 5F)

No emission/removal is reported for “other land remaining other land” in accordance with GPG for LULUCF. As with most other land use categories information on land use conversion is not available at present.

This land is identified as a key area considering reported land use categories but not as part of applicable land use categories due to its definition as unmanaged.

7.8.1 Planned improvements regarding other land

Data on area covered by this category will be collected in connection with construction of the new geographic land use database. Definition of other land as land use category will be elaborated in that contest too.

7.9 Other (5)

Three emission/removal categories are reported under other. Harvested Wood Products, non-CO₂ emission form Grassland organic soil and emission/removal due to Revegetation.

7.9.1 Harvested Wood Products

No data is available on stock changes in harvested wood products and therefore not estimated. There are no planned improvements regarding recording of this stock.

7.9.2 Grassland organic soil

Under this item non-CO₂ emission from grassland are reported. Present structure of Reporter software does not allow reporting this emission under Grassland land use category.

7.9.2.1 Other emissions (5(I), 5(II), 5(III))

Grasslands in Iceland are generally not fertilized. The main exception is as part of revegetation activity. Use of fertilizers in revegetation is reported separately. Direct N₂O emission from eventual use of N fertilisers on grassland is included under emission from agricultural soils.

Emission of N₂O due to drainage of organic soils is reported here as aggregated number under “Other” specified as grassland on drained organic soils consistent with reporting of CO₂ emission from drained soils. This factor is identified as level key source of LULUCF.

7.9.2.2 Emission factors

Emission of N₂O from drained organic soil is calculated according to Tier 1 using emission factor EF = 1,8 [kg N₂O-N ha⁻¹ yr⁻¹] (GPG for LULUCF Table 3A.3.4) fore nutrient rich organic soils considering high N content.

7.9.2.3 Planned improvements regarding grassland organic soil

See chapter 7.5.6

7.9.3 Revegetation

7.9.3.1 Overview

Since settlement of Iceland large areas of the former vegetated areas have been severely eroded and in large areas the entire soil mantle has been swept away. It has recently been estimated that total of 60-250 *10³ kt C has been oxidized and released into the atmosphere in the past millennium (Óskarsson et al. 2004).

The current ongoing loss of SOC due to erosion was in the same study estimated 50-100 kt C yr⁻¹ (Óskarsson et al. 2004). No attempt is made to include that estimate in the CRF.

The revegetation of deserted areas sequesters some of the carbon back into the soil.

The Icelandic Soil Conservation was established in 1907. Its main purpose was and still is the prevention of ongoing erosion and the revegetation of lost areas. Until 1970 recording of soil conservation and revegetation activities was very limited and consisted only of occasional maps and reports. From 1970 to 1990 most of the activities involved spreading of seeds and/or fertilizers by airplanes. These activities are to a large extent recorded. From 1990 the importance of flight has decreased as other methods have taken over and cooperation with farmers and other parties of interest has increased. At the same time, recording of activities has developed.

7.9.3.2 Carbon stock changes

Changes in carbon stock in soil and vegetation due to revegetation are reported under the category 'other'. Preferably, revegetation would be reported as a separate main land use category or subcategory of relevant land use categories, addressing changes in individual pools and other emissions more directly. Due to this reporting format aggregated carbon stock changes (accumulation) is reported as CO₂ removal. In order to increase the transparency of reporting for this category and improve consistency to other land use categories a sectoral background table for revegetation comparable to other CRF prepared by the Reporter software was prepared for this report (Table 7.10). There the reported removal of CO₂ is calculated according to the estimated area revegetated and the emission/removal factor for carbon stock changes in living biomass and soil. The area reported is divided to three periods: pre-1990, 1990 - inventory year, and the area revegetated in the inventory year.

The reported CO₂ removal by revegetation after 1990 is based on area recorded by Icelandic Soil Conservation. The removal by revegetation before 1990 is based on estimate of area provided by the Icelandic Soil Conservation.

The Icelandic Soil Conservation has for this year's submission revised previous area estimates for all years from 1990, which explains the large increase (26%) in removal from 423 Gg CO₂ reported for the year 2004 in 2006 to 533 Gg CO₂ in this year submission. The revised area estimate changes the reported CO₂ removal to 511 Gg CO₂ and the increase from last submission to 4%.

This factor is identified as a key source of LULUCF and as a key area considering only applicable land use categories.

TABLE 5.G SECTORAL BACKGROUND DATA FOR LAND USE, LAND-USE CHANGE AND FORESTRY												Iceland
Revegetated land												2004
(Sheet 1 of 1)												2006
GREENHOUSE GAS SOURCE AND SINK CATEGORIES		ACTIVITY DATA	IMPLIED EMISSION FACTORS					EMISSIONS/REMOVALS				
Land-Use Category	Sub-division ⁽¹⁾	Total area (kha)	Carbon stock change in living biomass per area ^(2,3)			Net carbon stock change in dead organic matter per area ⁽³⁾	Net carbon stock change in soils per area ⁽³⁾	Carbon stock change in living biomass ^(2,3)			Net carbon stock change in dead organic matter ⁽³⁾	Net carbon stock change in soils ⁽³⁾
			Increase	Decrease	Net change			Increase	Decrease	Net change		
			(Mg C/ha)					(Gg C)				
H. Total Revegetated Land		193,91	0,07	0,00	0,07	0,00	0,68	14,54	0,00	14,54	0,00	130,89
1. Revegetated Land remaining Revegetated Land		185,92	0,08	0,00	0,08	0,00	0,68	13,94	0,00	13,94	0,00	125,50
	after 1990	80,92	0,08	0,00	0,08	0,00	0,67	6,07		6,07		54,62
	before 1990	105,00	0,08	0,00	0,08	0,00	0,68	7,88		7,88		70,88
2. Land converted to Revegetated Land ⁽⁴⁾		7,99	0,07	0,00	0,07	0,00	0,68	0,60	0,00	0,60	0,00	5,39
2.1 Forest converted to Revegetated Land		0,00						0,00	0,00	0,00	0,00	0,00
2.2 Cropland converted to Revegetated Land		0,00						0,00	0,00	0,00	0,00	0,00
2.3 Grassland converted to Revegetated Land		0,00						0,00	0,00	0,00	0,00	0,00
2.4 Wetlands converted to Revegetated Land		0,00						0,00	0,00	0,00	0,00	0,00
2.5 Settlements converted to Revegetated Land		0,00						0,00	0,00	0,00	0,00	0,00
2.6 Other Land converted to Revegetated Land		7,99	0,07	0,00	0,07	0,00	0,68	0,60	0,00	0,60	0,00	5,39
		7,99	0,07	0,00	0,07	0,00	0,68	0,60		0,60		5,39

(1) Land categories may be further divided according to climate zone, management system, soil type, vegetation type, tree species, ecological zones or national land classification.

(2) CO₂ emissions and removals (carbon stock increase and decrease) should be listed separately except where, due to the methods used, it is technically impossible to separate information on increases and decreases.

(3) The signs for estimates of increases in carbon stocks are positive (+) and of decreases in carbon stocks are negative (-).

(4) A Party may report aggregate estimates for all conversions of land to forest land when data are not available to report them separately. A Party should specify in the documentation box which types of land conversion are included. Separate estimates for grassland conversion should be provided in table 5 as an information item.

Documentation box:
Parties should provide detailed explanations on the Land Use, Land-Use Change and Forestry sector in Chapter 7: Land Use, Land-Use Change and Forestry (CRF sector 5) of the NIR. Use this documentation box to provide references to relevant sections of the NIR if any additional information and/or further details are needed to understand the content of this table.

Table 7.10 Sectoral background table for revegetation not included in CRF prepared by the UNFCCC Reporter software

7.9.3.3 Other emissions (5(I), 5(II), 5(III))

The direct emissions of N₂O from the use of N-fertilizers on revegetated land are reported separately for revegetation for the first time.

Drainage of revegetated area is reported as not occurring and subsequently no emission reported. Non- CO₂ emission from disturbance associated with conversion of land to cropland is not estimated due to lack of data.

7.9.3.4 Emission/removal factors

The Icelandic Soil Conservation records the revegetation efforts conducted. In 1998-2000 a special governmental effort to sequester carbon with revegetation and afforestation was carried out. Along with that effort a research effort to document carbon sequestration and estimate its rate was carried out (Arnalds et al. 2000).

No Tier 1 default emission/removal factors are available for revegetation effort. The emission factor used for calculating emission/removal due to revegetation efforts are estimated as -0,75 kt C/kha/yr based on pre-cautious estimate of results from the research effort conducted 1998-2000. Also based on the same research effort the contribution of changes in carbon stock of living biomass and soil were estimated 10% and 90% respectively. All revegetated areas are assumed to accumulate carbon stock at same rate.

7.9.3.5 Uncertainties QA/QC

Calculation of removal/emission of carbon due to revegetation depends on the size of the area and the chosen emission/removal factor. The approach is 'Tier 1 (2)' approach based on simple removal rate factor based on measurement of chronosequential accumulation of carbon on revegetated areas of a known age. Both numbers have a large uncertainty.

The areas where revegetation is carried out are very variable with regards to soil climate condition and methods used. Success of revegetation efforts is also very variable and consequently, as is the rate of sequestration. Although some of the sources of this variability have been identified, it is far from being totally explained (Arnalds et al. 1999; Arnalds et al. 2000; Arnalds et al. 2002).

The mapping method and registration of the revegetation on the first year of reporting (1998) was based on records of the site name and estimate of hectares within that location where the activity took place. The estimated number of hectares is partly based on amount of seeds and fertilizers used. This method may have introduced a relatively large error into the area estimates and may bring a risk of either double counting or not counting some areas. Since 1998 the reported size of area subjected to revegetation has been increasingly based on GPS recordings. The carbon stock changes, reported this year, are based on the Icelandic Soil Conservation GPS-mapping of area revegetated and revised estimate of those areas not yet mapped.

The size of the land revegetated prior to 1990 is much more uncertain than after that time. It is possible that today some of this land should today be identified under a different land use category.

Generally it is a necessary part of the revegetation effort to protect the area from grazing by erecting permanent fences. In some cases the whole area within such

fences is reported as revegetated although only a part of it has been directly subject to the field of activities such as fertilization or seed spreading. It is important to bear in mind that the registration was designed to serve other purposes than the needs of greenhouse gas inventories.

7.9.3.6 Planned improvements regarding revegetation

An effort in GIS mapping of the revegetation areas and improvements of the precision of size estimate of the areas has been ongoing since 1998. This effort has resulted in increased accuracy of area estimates reflected in this year's submission. In 2005 and 2006 some of the areas, sampled during 1998-2000 research effort, were resampled. Results from this research are not yet available. Improvements in both the sequestration rate estimates and area recording aim at establishing a transparent, verifiable inventory for revegetation efforts accountable according to the Kyoto Protocol.

Two main improvements are planned and partly being carried out. First there is the improvement in recording of activities both, in location and description of activities and management. The second improvement planned is pre-activity sampling to establish a baseline for future comparisons of SOC.

The Icelandic Soil Conservation has introduced sample based approach combined with GIS mapping to identify land subjected to revegetation and to improve emission/removal factors and quality control on different activity practices. The systematic sampling approach to identify land subjected to revegetation and changes in carbon stocks will provide more accurate estimates of land use and stock changes and enable the quantification of uncertainty for these estimates. The systematic sampling plots are selected on the same grid as used by the Icelandic Forestry Research.

7.9.3.7 Recalculation

Emission of N₂O and removal of CO₂ has been recalculated for all years from 1990. The separation of fertilizers used in revegetation and new revised area estimate of revegetated areas for all years from 1990 has changed the previous reported emission/removals.

Table 7.11 shows the new area estimate of the Icelandic Soil Conservation for 1990-2005 and calculated C stock changes and CO₂ removal for that area.

Table 7.12 shows the amount of synthetic fertilizers used on revegetated areas in the years 1990-2005 and calculated direct N₂O emission due to that usage.

<i>Background table for CO₂ emission/removal by revegetation</i>									<i>Iceland 2005 submission 2007</i>
<i>year of initiation</i>	<i>Area [kha]</i>	<i>Emission /removal factor [Gg C/kha]</i>		<i>Net carbon stock change Gg C</i>			<i>CO₂ emission/removal Gg CO₂</i>		
		<i>soil</i>	<i>living biomass</i>	<i>soil</i>	<i>living biomass</i>	<i>total</i>	<i>new area</i>	<i>all years 1990 to the year prior to</i>	<i>total reported CO₂ that year</i>
before 1990	105	0,675	0,075	70,875	7,875	78,750	-288,75		
1990	3,038	0,675	0,075	2,051	0,228	2,279	-8,355	0,000	-297,105
1991	3,291	0,675	0,075	2,221	0,247	2,468	-9,050	-8,355	-306,155
1992	4,393	0,675	0,075	2,965	0,329	3,294	-12,080	-17,405	-318,234
1993	2,785	0,675	0,075	1,880	0,209	2,089	-7,659	-29,484	-325,893
1994	4,192	0,675	0,075	2,830	0,314	3,144	-11,528	-37,143	-337,421
1995	5,508	0,675	0,075	3,718	0,413	4,131	-15,147	-48,671	-352,568
1996	4,742	0,675	0,075	3,201	0,356	3,557	-13,041	-63,818	-365,609
1997	5,529	0,675	0,075	3,732	0,415	4,147	-15,205	-76,859	-380,813
1998	7,177	0,675	0,075	4,844	0,538	5,383	-19,737	-92,063	-400,550
1999	7,823	0,675	0,075	5,281	0,587	5,867	-21,513	-111,800	-422,063
2000	6,606	0,675	0,075	4,459	0,495	4,955	-18,167	-133,313	-440,230
2001	4,031	0,675	0,075	2,721	0,302	3,023	-11,085	-151,480	-451,315
2002	4,791	0,675	0,075	3,234	0,359	3,593	-13,175	-162,565	-464,490
2003	7,133	0,675	0,075	4,815	0,535	5,350	-19,616	-175,740	-484,106
2004	9,877	0,675	0,075	6,667	0,741	7,408	-27,162	-195,356	-511,268
2005	7,991	0,675	0,075	5,394	0,599	5,993	-21,975	-222,518	-533,243

Table 7.11 Revised area estimate for revegetated area and calculated C stock changes and CO₂ removal.

Year	Tons fertilizers used in revegetation	Tons N applied	Gg N	Gg N ₂ O-N	Gg N ₂ O
1990	1390,1	333,6	0,334	0,004	0,007
1991	1668,6	400,5	0,400	0,005	0,008
1992	1521,4	365,1	0,365	0,005	0,007
1993	870,3	208,9	0,209	0,003	0,004
1994	787,2	188,9	0,189	0,002	0,004
1995	1260,9	302,6	0,303	0,004	0,006
1996	1116,9	268,1	0,268	0,003	0,005
1997	1475,6	354,1	0,354	0,004	0,007
1998	1884,3	452,2	0,452	0,006	0,009
1999	1704,6	409,1	0,409	0,005	0,008
2000	1508,2	362,0	0,362	0,005	0,007
2001	1617,1	388,1	0,388	0,005	0,008
2002	1763,4	423,2	0,423	0,005	0,008
2003	2748,9	659,7	0,660	0,008	0,013
2004	3093,2	742,4	0,742	0,009	0,015
2005	3385,0	812,4	0,812	0,010	0,016

Table 7.12 Amount of synthetic fertilizers used on revegetated land and calculated direct N₂O emission.

7.10 Biomass burning (5V)

Accounting of biomass burning for all land use categories is addressed commonly in this section.

No emissions due to biomass burning are reported. It is considered not occurring for controlled burning of forest land, land converted to forest land, land converted to cropland, forest land converted to grassland, forest land converted to wetland and wildfires on forest land converted to: cropland, grassland or wetland. For other categories it is not estimated due to lack of information.

Burning the biomass on grazing land near the farm was for a long time common practice in sheep farming. This management regime of grasslands and wetlands is becoming less common. The recording of the activity is minimal although formal approval of local polity authority is needed for safety and birdlife protection purposes. A large wildfire broke out in the year 2006 and research project was initiated aimed at recording the ecological effects of biomass burning. This project is expected to provide data for Tier 2 assessment of amount of biomass burned per area burned. As land use identification is expected to improve in general due to the planned increased effort on establishing a georeferenced land use database, information on area burned annually are also expected to improve.

7.11 Planned improvements of emission/removal data for LULUCF

Improvements which apply specifically to one of the land use categories or one of their pools are listed above in relevant chapter.

Simultaneously with gathering of land use information for the purpose of the new georeferenced land use database, data will be collected regarding the carbon stocks of the land use category used in the classification. These efforts are aimed at gradually improving the reliability of reported emission/removal of the LULUCF sector and

enable the movement from Tier1, which is presently used to calculate emission/removal in most categories, to higher tier levels.

The results of ongoing and recent research activity on emissions/removal and stocks in several ecosystems will be implemented in emissions calculations.

8 WASTE

8.1 Overview

Practices of waste disposal and wastewater treatment have undergone a radical change in Iceland since 1990. Open pit burning that used to be the most common means of waste disposal outside the capital area, has gradually decreased since 1990, as landfills have become the main option. Recycling of waste has also increased due to efforts made by local municipalities. A twofold increase was noted between 1994 and 2005. Municipalities have also increasingly cooperated to run waste collection schemes and operate common landfill sites. This has resulted in larger landfills and enabled closedown of a number of small sites. Currently about 76% of municipal waste is landfilled, 20% recycled or recovered, 3% incinerated with energy recovery and 0,5% are incinerated without energy recovery.

In 1990, 6% of the Icelandic nation was connected to wastewater treatment plants, but in 2005 the ratio was 60,7%.

8.1.1 Methodology

The calculation of greenhouse gas emissions from waste is based on the methodologies suggested by the IPCC Guidelines and the Good Practice Guidance.

As indicated in Table 1.1, the key source analysis performed for 2005 has revealed that in terms of total level and/or trend uncertainty the only key source in the waste sector is the following:

- Emissions from Solid Waste Disposal Sites – CH₄ (6A)

8.1.2 Completeness

Table 8.1 gives an overview of the IPCC source categories included in this chapter and presents the status of emission estimates from all sub-sources in the industry sector.

Table 8.1 Waste - completeness

Sector	Direct GHG			Indirect GHG			
	CO ₂	CH ₄	N ₂ O	NO _x	CO	NMVOC	SO ₂
Solid waste disposal on land							
Managed waste disposal on land	NE	X	NA	NA	NA	NE	NA
Unmanaged waste disposal on land	NE	X	NA	NA	NA	NE	NA
Wastewater treatment							
Industrial wastewater	NE	NE	NE	NE	NE	NE	NE
Domestic and commercial wastewater	NE	X	X	NE	NE	NE	NE
Waste incineration							
Other	X	NE	X	X	X	X	X

8.2 Solid waste disposal sites

Methane from solid waste disposal sites is emitted during the biological decomposition of waste. This transformation of organic matter takes place in several

steps. During the first weeks or months, decomposition is aerobic and the main decomposition product is CO₂. When there is no oxygen left, the decomposition becomes anaerobic and methane levels starts to increase. After about one year CH₄ emissions peak and levels decrease over some decades.

In Iceland, solid waste disposal is divided between managed landfill sites and unmanaged landfill sites. Managed sites are typically deeper than 5 meters and have thorough registration on waste sorts and amounts disposed. Unmanaged landfill sites are shallow with less than 5 meters of waste. Total waste going to these landfills is disaggregated into two major waste streams, municipal solid waste (MSW) and industrial waste (IW).

The methodology for calculating methane from solid waste disposal on land is in accordance with the IPCC First Order of Decay method. The total amount of methane gas generated by the disposal of MSW and IW on landfill sites is calculated with the following equations.

$$\text{CH}_4 \text{ emissions (Gg/yr)} = [(Q_{\text{MSW}} + Q_{\text{IW}}) - R] \cdot (1 - \text{OX})$$

Q_{MSW} = Methane generated by MSW disposal (Gg/yr)

Q_{IW} = Methane generated by IW disposal (Gg/yr)

R = Methane recovery (Gg/yr)

OX = Oxidation factor

$$Q_{\text{MSW}} \text{ (Gg/yr)} = \sum_i Q_i$$

Q_i = Methane generated by landfilled waste type i (Gg/yr)

i = type of waste (see par. 8.2.4.1)

$$Q_i \text{ (Gg/yr)} = E \cdot 16/12 \cdot F$$

E = Calculates the total mass of DDOC decomposed to methane each year.

F = is the fraction of methane in developed landfill gas

$$E = C + H_{\text{last year}} \cdot (1 - \text{exp1})$$

C = DDOC_m Decomposable Degradable Organic Carbon

H = DDOC_m accumulated in SWDS end of year

$$H = B + (H_{\text{last year}} \cdot \text{exp1})$$

B = DDOC_m not reacted

$$C = D \cdot (1 - \text{exp2})$$

D = Decomposable DOC (DDOC_m) deposited

$$B = D \cdot \text{exp2}$$

D = Decomposable DOC (DDOC_m) deposited

$$D = W \cdot \text{DOC} \cdot \text{DOC}_f \cdot \text{MCF}$$

W = Amount of waste landfilled

DOC = Degradable Organic Carbon

DOC_f = Fraction of DOC dissimilated

MCF = Methane Correction factor

8.2.1 Activity data

Activity data on waste in Iceland has proven to have been insufficient in the past. There is little information about actual amounts of generated waste as well as on its composition and characteristics, before 1990. Reporting of waste amounts received by managed landfill sites started after 1980 and is done by the landfill operators. Consistent and relatively reliable data-sets on total waste generation and treatment are available from 1995.

Using the calculation based on the Icelandic GDP (Gross Domestic Product) the total amount of generated waste can be extrapolated from 1994 back to 1950. This GDP based calculation uses 2004 as reference year. It calculates the amount of generated MSW per person per year and the total generated for both MSW and IW. Data quality for the reference year is ensured because of accurate waste reporting for that year.

GDP is strongly correlated with a country's waste production and is a reliable estimation method. Icelandic GDP figures date back before 1950 and are considered reliable. With regard to total waste amounts generated, it has been decided to use available reported waste figures above the on GDP-based waste figure calculations.

The activity data was mostly collated by the EFA. Secondary data sources are the municipalities and the larger waste companies in Iceland. The total amounts of MSW and IW generated and treated in Iceland between 1950 and 2005 are reported in Table 8.2.

Table 8.2 Waste generation and treatment from 1950 to 2005

Year	Generated (A+B+C)*			A. Landfilled		B. Recycled		C. Incinerated	
	Total	MSW	IW	MSW	IW	MSW	IW	MSW	IW
1950	42	8	34	8	34	-	-	-	-
1960	69	16	53	16	53	-	-	-	-
1970	112	29	83	29	83	-	-	-	-
1980	214	60	154	60	154	-	-	-	-
1990	288	87	201	58	196	-	-	29	5
1991	289	88	201	60	196	-	-	29	5
1992	282	87	195	59	190	-	-	28	5
1993	287	89	198	62	193	-	-	27	5
1994	299	93	206	68	201	-	-	25	4
1995	360	114	246	90	242	-	-	24	4
1996	370	117	253	93	249	1	-	23	4
1997	378	120	258	96	254	2	-	22	4
1998	386	123	263	97	260	6	-	20	3
1999	394	126	268	98	265	11	-	17	3
2000	403	130	273	97	270	17	-	16	3
2001	413	133	280	94	267	23	10	16	3
2002	420	137	283	92	237	30	43	15	3
2003	431	140	291	94	242	32	47	14	2
2004	443	147	296	101	243	31	50	15	3
2005	462	152	310	99	253	38	54	15	3

* Amounts is x 1.000 tons

8.2.2 Emission factors

Municipal solid waste

Municipal Solid Waste corresponds to waste from households and similar waste from the commerce and trade industry. MSW can be disaggregated into a mix of waste categories that contain significant fractions of biodegradable carbon, which are: food, garden, paper, wood, textile, nappies and sludge (from wastewater handling (Sector Waste 6B)).

The composition of MSW going to landfill has been surveyed starting from 1999 and is done by SORPA, the biggest waste treatment facility in Iceland. SORPA serves the Reykjavik capital area and thus covers around 63% of the Icelandic population. The composition of MSW over the last 7 years has shown to be relatively consistent. Because very little is known about the MSW composition before 1999 the average composition from 1999 to 2004 has been used in the IPCC model for each year between 1950 and 2004.

It is understood that different MSW compositions are likely to have existed over the last 60 years. For example, the fraction of garden waste in 1950 might have been higher than in 2000. Also the fraction of plastic (packaging) waste in MSW is expected to have increased significantly since 1950.

A sensitivity analysis, however, showed very little variation in total methane emission in Sector 6A when applying different (estimated) waste compositions between 1950 and 2004. The difference calculated did not exceed 2%. Because it is impossible to estimate the exact composition of waste each year and has very little effect on the final outcome, the composition of MSW has not been further investigated. The waste composition surveys results for 1999 to 2004 and their averages are reported in Table 8.3.

Table 8.3: Municipal Solid waste composition survey results 1999 – 2004

Type of waste	1999	2000	2001	2002	2003	2004	Average
Food waste	33%	28%	31%	26%	24%	26%	28,1%
Garden waste	4%	0%	1%	0%	2%	1%	1,4%
Paper and Cardboard	24%	29%	21%	22%	26%	27%	24,8%
Wood waste	0%	1%	1%	1%	1%	0%	0,6%
Textile waste	4%	4%	3%	3%	3%	4%	3,4%
Diapers/nappies	5%	4%	6%	7%	5%	6%	5,6%
Sludge	4%	4%	4%	4%	4%	4%	3,9%
Plastics, other inert	26%	30%	33%	37%	35%	32%	32,2%

The emission factors and parameters for IPCC Category 6A Municipal Solid Waste are reported in Table 8.4

Table 8.4: Emission factors and parameters for Municipal Solid Waste

Parameters	Food	Garden	Paper	Wood	Textile	Nappies	Sludge
MSW composition (average 1999 -2004)	28,1%	1,4%	24,8 %	0,6%	3,4%	5,6%	3,9%
Methane Correction Factor (MCF)*							
- Unmanaged-shallow				0,4			
- Managed				1,0			
- Uncategorized				0,6			
Fraction of degradable organic carbon dissimilated (DOC _F)*				0,5			
Degradable organic carbon (DOC)*	0,15	0,17	0,4	0,43	0,24	0,24	0,05
Methane generation constant (k)*	0,185	0,1	0,06	0,03	0,06	0,1	0,185
Half-life time (h) (years) (h = Ln(2)/k)	4	7	12	23	12	7	4
Delay time (month)*				6			
Number of considered years				55			
Fraction of CH ₄ in landfill gas (F) *				0,5			
Oxidation factor (OX) *				0,05			
Conversion factor (C to CH ₄)				1,33			

* IPCC default value

Industrial waste

Industrial waste (IW) comes from agriculture, fisheries and other industrial activities. The amounts of IW used in the IPCC model are excluding separated waste fractions such as scrap metal, tyres and construction and demolition waste. It is expected that significant fractions of MSW-related waste can be found in IW and will be further explained under paragraph 8.2.3.

The emission factors and parameters for IPCC Category 6A Industrial Waste are reported in Table 8.5.

Table 8.5: Emission factors and parameters for Industrial Waste

Parameters	
Methane Correction Factor (MCF)*	
- Unmanaged-shallow	0,4
- Managed	1,0
- Uncategorized	0,6
Fraction of degradable organic carbon dissimilated (DOC _F)*	0,5
Degradable organic carbon (DOC)*	0,15
Methane generation constant (k)*	0,09
Half-life time (h) (years) (h = Ln(2)/k)	8
Delay time (month)*	6
Number of considered years	55
Fraction of methane in landfill gas*	0,5
Oxidation factor (OX) *	0,05
Conversion factor (C to CH ₄)	1,33

* IPCC default value

Landfill gas recovery

The recovery of landfill gas (CH₄) is done at only one landfill site (Álfsnes) in Iceland, which receives the waste from the Reykjavík capital area. The recovery of CH₄ from landfill gas started in 1997 and amounts are reported in Table 8.6.

Table 8.6: Landfill gas recovery in Iceland, 1997 - 2005

Gg CH ₄	1997	1998	1999	2000	2001	2003	2004	2005
Methane recovery from Solid Waste disposal sites (SWDS)	0,105	0,240	0,349	0,430	0,407	1,108	1,143	1,550

Uncertainties

The preliminary estimate of quantitative uncertainty has revealed that the uncertainty of CH₄ emissions from solid waste disposal sites is 52%. The quality of the activity data for Iceland may be considered sufficient, but needs further improvement to ensure its accuracy and quality. The uncertainties in the IPCC model for Sector 6A are regarding:

- **Landfilled waste between 1950 to 1980**

The exact amounts of waste going to managed or unmanaged landfill sites between 1950 and 1980 are unknown. Therefore the Methane Correction factor (MCF) in the IPCC model has been set to uncategorised for this period (MCF = 0,6 – see Table 8.4 and 8.5).

- **Amount and composition of Industrial waste**

The total amount of Industrial Waste has shown to be very significant over the last 10 years (1995-2005). Because of these large amounts of waste going to landfill, the emissions that are calculated using the IPCC model are significant as well.

Although separated waste sorts such as scrap metal, tyres and construction and demolition waste are excluded from IW it is expected to include waste amounts that should be allocated to MSW. This is because large amounts of company related waste, thus with a similar composition as MSW, is included in mixed fraction of Industrial Waste.

The exact composition of mixed IW and thus the fraction of biodegradable waste remain unknown. Therefore, both the total amount of mixed IW and its composition need further investigation. This has shown to be a barrier to accurate emission calculations as well as general waste statistics.

The methane emission from landfilled IW might be overestimated. However, if a fraction from IW is allocated to MSW, it will result in an increase of methane emission from landfilled MSW. The resulting absolute difference has not been estimated.

- **MSW composition between 1950 and 1998**

The composition of MSW for the years between 1950 and 1998 is rather difficult to estimate. The sensitivity analysis, using different estimated waste compositions showed very little change in total methane emissions. The

calculated differences in total methane emission in Sector 6A did not exceed 2%.

Recalculations

In 2006 Iceland applied for the first time the IPCC First Order of Decay (FOD) method. A preliminary version of the IPCC Model was used. A flaw in this preliminary version of the model resulted in wrong results for the years 2000 to 2004. This was corrected for this submission.

Table 8.7 Comparison between results of the 2006 and 2007 submissions, in Gg CH₄

Year	2006 submission	2007 submission
2000	7,392	7,315
2001	7,630	7,652
2002	8,056	7,390
2003	7,595	7,562
2004	7,740	7,496

8.3 Emission from Wastewater Handling (6B)

Very few wastewater treatment plants are operational in Iceland and most of them are located in the greater Reykjavík capital area and a few other larger municipalities. The wastewater treating systems are mostly settling tanks or septic tanks and filters (e.g. sandfilters or biological filters).

The sludge from wastewater handling is disposed on (managed and unmanaged) landfill sites. Amounts of sludge are included in total generated waste. Emissions from sludge disposal on landfills are included under Solid Waste Disposal (6A). No methane gas is recovered from wastewater handling in Iceland.

8.3.1 Methodological issues

The Icelandic methodology for calculation of methane emission from wastewater handling is consistent with the IPCC default methodology. To estimate the CH₄ emissions from domestic wastewater the '*Decision Tree for CH₄ emission*' presented in the IPCC Good Practice Guidance was used. Because very little data on wastewater treatment are available and its quality is poor, Iceland used the 'Check method' in accordance with IPCC guidelines.

$$\mathbf{WM = P \cdot D \cdot SBF \cdot EF \cdot FTA \cdot 365 \cdot 10^{-12}}$$

WM = Annual CH₄ emission from domestic wastewater (Tg)

P = Population (person)

D = Organic load in biochemical oxygen demand per person (g BOD/person/day)

SBF = Fraction of BOD that readily settles

EF = Emission factor (g CH₄/g BOD)

FTA = Fraction of BOD in sludge that degrades anaerobically

The 'Check method' equation can be used to roughly estimate the CH₄ emissions from domestic wastewater. The variable P (population) in this equation is country specific and only includes the population that is connected to the various wastewater treatment facilities. For the other emission factors, either IPCC default values or estimated values are used.

Nitrous oxide emissions from human sewage were calculated according to the IPCC default method, which is based on the annual per capita protein intake.

$$\text{N}_2\text{O} = \text{Protein} \cdot \text{Frac(NPR)} \cdot \text{population} \cdot \text{EF} \cdot 44/28$$

Activity data

The activity data used for the 'Check method' is represented by the population portion that is connected to wastewater handling facilities and is reported in Table 8.8. The total number of population is obtained from the Statistics Iceland.

The percentage of population that is connected to wastewater handling facilities has been reported in the EFTA report to the Surveillance Authority regarding the implementation of Directive 91/271/EU on the treatment of wastewater from built-up areas.

Table 8.8: Total population and population connected to wastewater handling facilities in Iceland

	Population	Connected to wastewater facilities		
		Total	Septic tanks	Sand filters, settling tanks, etc.
1990	255.708	6,0%	0%	6,0%
1991	259.577	6,0%	0%	6,0%
1992	262.193	6,0%	0%	6,0%
1993	264.919	6,0%	0%	6,0%
1994	266.783	10,0%	4,0%	6,0%
1995	267.806	10,0%	4,0%	6,0%
1996	269.727	10,0%	4,0%	6,0%
1997	272.069	10,0%	4,0%	6,0%
1998	275.264	14,4%	8,4%	6,0%
1999	279.049	22,4%	16,4%	6,0%
2000	282.849	39,0%	33,0%	6,0%
2001	286.250	39,0%	33,0%	6,0%
2002	288.201	60,7%	48,7%	12,0%
2003	290.490	60,7%	48,7%	12,0%
2004	293.291	60,7%	48,7%	12,0%
2005	299.404	60,7%	48,7%	12,0%

Emission factors

Of the total population connected to wastewater handling facilities, some are connected to handling facilities such as septic (and settling) tanks and some are connected to sand-filters. Different FTA factors apply to the two handling methods. It is expected that emissions from wastewater handling in Iceland based on the 'Check method' are still overestimated. The annual per capita protein intake is based on Dietary Surveys of the Icelandic Nutrition Council and the Dietary Survey Unit for Nutrition Research performed in 2002 to 2003. The emission factors and parameters for IPCC Category 6B Wastewater Handling are reported in Table 8.9.

Table 8.9: Emission factors and parameters for Wastewater

Parameters	
D *	60 g/person/day
SBF _{septic}	0,5
SBF _{sand filter, etc.}	0,33
EF *	0,6 g CH₄/g BOD
FTA _{septic}	0,6
FTA _{sand filter, etc.}	0,2
Protein	31,76 kg/person/year

* IPCC default value

Uncertainties

The uncertainties in the IPCC model for Sector 6B are regarding:

- **SBF parameter**

Around 50% of the BOD in domestic water is associated with non-dissolved solids, much of which rapidly settles. A conventional settling tank typically removes 33% of suspended solids, whereas 50% is more appropriate to many long-term processes as lagoons or septic-tanks. Since both conventional and long term processes are used in Iceland, both SBF values are used in the Check method.

- **FTA parameter**

For countries or areas that are extensively sewerred, use only aerobic processes and whose sludge is treated by non-CH₄ producing procedures or by anaerobic digestion with combustion of CH₄, the FTA will be significantly lower or zero.

Wastewater handling in Iceland is done using both septic-tanks and (sand) filters. In septic-tanks there is possible anaerobic digestion and therefore a higher FTA (0,6) applies than for (sand)filters (0,2). Similar to the SBF parameter, both FTA values are used in the 'Check method'.

The calculation of emissions from wastewater handling confirms the expectation done in the NIR submission of 2005 that very little emission is generated from wastewater handling in Iceland.

8.4 Waste incineration

Emissions from waste incineration with energy recovery are reported in sector 1A1a (public electricity and heat production). Emissions from waste incineration have decreased by 99,9% from 1990 to 2005. This is because the total amount of waste being incinerated in Iceland has decreased while increasing levels have been incinerated with energy recovery and thus reported under 1A1a. Waste incineration without energy recovery is virtually non-existent.

The methodology for calculating emissions from waste incineration is in accordance with the IPCC Guidelines. The activity data are the waste inputs into the incinerator, and the emission factor is based on the carbon content of the waste that is of fossil origin only. The burn out efficiency of the combustion is also included in the calculation. Although the most accurate way to estimate CO₂ emissions is by disaggregating the activity data into different waste types (e.g. municipal solid waste,

clinical waste, hazardous waste) this could not be done for this submission. The following equation is used for calculating CO₂ emissions from waste incineration:

$$\text{CO}_2 \text{ emissions (Gg/yr)} = \text{IMSW} \cdot \text{CCW} \cdot \text{FCF} \cdot \text{BEF} \cdot 44/12$$

IMSW: Amount of incinerated waste (Gg/yr)

CCW: Fraction of carbon content in waste

FCF: Fraction of fossil carbon in waste

BEF: Burn out efficiency of incinerator

44/12: Conversion from C to CO₂

Activity data

Activity data on incinerated waste from major incineration plants have been collected by the EFA since 2000. Historic data as well as data on open pit burning not reported to EFA, was estimated with the assumptions that 500 kg of wastes have been incinerated per inhabitant in the communities where waste is known to have been incinerated (both in primitive incineration plants as well as open pit burning) in 1990, 1995 and 2000 and interpolated in the years between. These communities were mapped by EFA in the respective years. The data after the year 2000 is considered rather reliable, but pre-2000 data very unreliable.

Emission factors

Data for estimation of CO₂ from waste incineration are default values for municipal solid waste (MSW) taken from the IPCC Good Practice Guidance. They are presented in Table 8.10.

Table 8.10 Parameters for waste incineration

Waste Stream	MSW
C content of waste	40%
Fossil Carbon as % of Total Carbon	40%
Efficiency of Combustion	95%

9 OTHER – GEOTHERMAL ENERGY

9.1 Overview

Iceland relies heavily on geothermal energy for space heating and to some extent for electricity production. Geothermal energy is generally considered to have relatively low environmental impact. Emissions of CO₂ are commonly considered to be among the negative environmental effects of geothermal power production, even though they have been shown to be considerably less extensive than from fossil fuel power plants.

As indicated in Table 1.1, the key source analysis performed for 2005 has revealed that geothermal energy is a key source in terms of both level and trend.

9.2 Methodology

Three major geothermal power plants in operation in Iceland are at Krafla, Svartsengi and Nesjavellir. The Svartsengi and Nesjavellir plants produce both electricity and hot water for space heating, whereas the Krafla plant generates electricity only. The total installed capacity of these three power plants is 195 MW and they produce about 17% of the total electricity used in Iceland.

Geothermal systems can be considered as geochemical reservoirs of CO₂. Degassing of mantle-derived magma is the sole source of CO₂ in these systems in Iceland. CO₂ sinks include calcite precipitation, CO₂ discharge to the atmosphere and release of CO₂ to enveloping groundwater systems. The CO₂ concentration in the geothermal steam is site and time-specific, and varies greatly between areas, wells within an area as well as the time of extraction.

Iceland reported greenhouse gas emissions from geothermal energy utilization prior to 2000 when it was decided to not include these emissions in the national emissions, as research in Iceland and some other countries indicated great uncertainty in the estimation of such emissions. At the time it was considered likely that no net emissions were taking place from geothermal power plants, only a relocation of the natural emissions within the wider geothermal area. Recent research, however, indicates that emissions associated with the utilisation of geothermal energy in Iceland do constitute a net increase in emissions.

The total emissions estimate is based on direct measurements. The enthalpy and flow of each well are measured and the CO₂ concentration of the steam fraction determined at the wellhead pressure. The steam fraction of the fluid and its CO₂ concentration at the wellhead pressure and the geothermal plant inlet pressure are calculated for each well. Information about the period each well discharged in each year is then used to calculate the annual CO₂ discharge from each well and finally the total CO₂ is found by adding up the CO₂ discharge from individual wells.

Recalculations

In the 2007 submissions minor correction were done for the years 1995 to 1998 because better data was obtained from Iceland Energy Research and for the years 2002 to 2003 typos were corrected.

REFERENCES

- Ármansson, H., Friðriksson, Þ., Kristjánsson, B. (2005). "CO₂ emissions from geothermal power plants and natural geothermal activity in Iceland". *Geothermics*.
- Arnalds, O., A. L. Aradóttir and G. Gudbergsson (2002). Organic Carbon Sequestration by Restoration of Severely Degraded Areas in Iceland. *Agricultural Practices and Policies for Carbon Sequestration in Soil*. R. L. J.M. Kimble, R.E. Follett, Lewis Publisher: 267-280.
- Arnalds, O. and E. Gretarsson (2001). Soil Map of Iceland. 2nd edition. Reykjavík, Agricultural Research Institute.
- Arnalds, Ó., Á. L. Aradóttir, A. Snorrason, G. Gudbergsson, T. H. Jónsson and A. M. Ágústsdóttir (1999). Organic carbon sequestration by restoration of severely degraded areas in Iceland. Iceland, Agricultural Research Institute: 1-19.
- Arnalds, Ó., E.F. Thorarinsdóttir, S. Metúsalemsson, Á. Jónsson, E. Gretarsson and A. Árnason. (2001). *Soil erosion in Iceland*. Reykjavík, Soil Conservation Service, Agricultural Research Institute.
- Arnalds, Ó., G. Guðbergsson and J. Guðmundsson (2000). "Carbon sequestration and reclamation of severely degraded soils in Iceland." *Búvísindi* 13: 87-97.
- EFA (2002). Starfsreglur um góða búskaparhætti (Codes of good agricultural practice). Starfshópur um meðferð úrgangs frá landbúnaði: 17. <http://www.ust.is/media/fraedsluefni/buskaparhaettir.pdf>
- EFA (2004). Landsáætlun um meðhöndlun úrgangs 2004 - 2016 (National waste treatment plan 2004 - 2016). Environment and Food Agency of Iceland: 46.
- EFA (2004). Report to the EFTA Surveillance Authority regarding the implementation of Directive 92/271/EU on the treatment of wastewater from built-up areas. Environmental and Food Agency of Iceland.
- Energy in Iceland: Historic Perspective, Present Status, Future Outlook (2004). National Energy Authority and Ministries of Industry and Commerce.
- External trade by HS-numbers (2005). Statistics Iceland. (available on the website: <http://hagstofan.is/?PageID=1053>)
- Friðriksson et al (2005). "CO₂ emissions and heat flow through soil, fumaroles, and steam heated mud pools at the Reykjanes geothermal area, SW Iceland". In preparation for Applied Geochemistry.
- Geirsson, Ó. (1975). Framræsla ("Drainage"). *Votlendi*. A. Garðarsson. Reykjavík, Landvernd. 4: 143-154.
- Icelandic Agricultural Statistics 2005. Icelandic Agriculture.

057 National Inventory Report 2007 - ICELAND
Icelandic Association of Farmers, Agricultural Genetic Resources Committee, and Nordic Gene Bank for Domestic Animals (2004). Icelandic Livestock Breeds, ISBN 9979-885-02

Icelandic Nutrition Council (2002). The Diet of Icelanders. Dietary Survey of the Icelandic Nutrition Council 2002. Main findings.

Inga Þórsdóttir, Ingibjörg Gunnarsdóttir (2006). The Diet of Icelandic 9- and 15-year-old children and adolescents. Dietary Survey of Unit for Nutrition Research 2002-2003.

IPCC (1997). Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories: Volume 1 – 3. Intergovernmental Panel on Climate Change.

IPCC (2000). Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories. Intergovernmental Panel on Climate Change.

IPCC, Ed. (2003). Good Practice Guidance for Land Use, Land-Use Change and Forestry, IGES.

Jónsson, T. H. and Ú. Óskarsson (1996). "Skógrækt og landgræðsla til að nema koltvísýring úr andrúmsloftinu. (Afforestation and soil reclamation as tool to remove carbon dioxide from atmosphere)." Ársrit Skógræktarfélagss Íslands 1996: 65-87.

Kamsma & Meyles (2003). Landfill Gas Formation in Iceland. Environmental and Food Agency of Iceland: 37

National Energy Forecast Committee (2005). Eldsneytisnotkun Íslendinga eftir notkunarflokkum, innlend notkun. Fuel use in Iceland per type, domestic use. (Available on the committee's website: <http://orkuspa.is/eldsneyti/Innlend.PDF>)

Náttúrufræðistofnun Íslands (2005). Ársskýrsla 2005 (annual report). Reykjavík, Náttúrufræðistofnun Íslands: 42.

Óskarsson, H. (1998). Peatland draining in Western Iceland. Icelandic Wetlands: exploitation and conservation. J. S. Ólafsson. Reykjavík, University of Iceland Press.: 121-129.

Óskarsson, H., O. Arnalds, J. Gudmundsson and G. Gudbergsson (2004). "Organic carbon in Icelandic Andosols: geographical variation and impact of erosion." CATENA 56(1-3): 225-238.

Óskarsson, M. and Eggertsson, M. (1991). Áburðarfræði (Fertilisers). Búnaðarfélag Íslands: 135.

SFT, SN (2004). National Inventory Report 2004 – Norway. Norwegian Pollution Control Authority & Statistics Norway: 176.

Sigurðsson, B. D. and A. Snorrason (2000). "Carbon sequestration by afforestation and revegetation as a means of limiting net-CO₂ emissions in Iceland." Biotechnol. Agron. Soc. Environ. 4(4): 303-307.

Snorrason, A. (2003a). Binding koldíoxíðs samfara nýskógrækt á Íslandi á árunum 1990-2000. Reykjavík, Rannsóknastöð Skógræktar ríkisins Mógilsá: 8.

Snorrason, A., B.D. Sigurðsson, G. Guðbergsson, K.Svavarsdóttir, and Þ.J.Jónsson (2003b). "Carbon sequestration in forest plantations in Iceland." Búvísindi (Icel. Agr. Sci.) 15(02): 81-93.

Snorrason, A. and B. Kjartansson. (2004). "Íslensk skógarúttekt. Verkefni um landssúttekt á skóglendum á Íslandi. Kynning og fyrstu niðurstöður. (Icelandic National Inventory. Project on inventory of forests in Iceland. Presentation and First Results)." Skógræktarritið(2): 101-108 (In Icelandic).

Statistical Yearbook of Iceland, 2005 (Statistics Iceland).

UNFCCC secretariat (2004). "Report of the individual review of the greenhouse gas inventory of Iceland submitted in the year 2004 (in-country review). FCCC/WEB/IRI/2004/ISL (available on the secretariat web site http://unfccc.int/files/national_reports/annex_i_ghg_inventories/inventory_review_reports/application/pdf/iceland_final_report_to_web.pdf)

UNFCCC secretariat. "Guidelines for the preparation of national communications by Parties included in Annex I to the Convention, Part I: UNFCCC reporting guidelines on annual inventories" (available on the secretariat web site <http://unfccc.int/resource/docs/cop5/07.pdf>)

ANNEX I: KEY SOURCES

According to the IPCC definition, key sources are those that add up to 90% of the total uncertainty in level and/or in trend. In the Icelandic Emission Inventory key source categories are identified by means of Tier 1 method.

A key source analysis was prepared for this round of reporting. Table 1.1 in Chapter 1 lists identified key sources. Table A1 shows the level assessment of the key source analysis and Table A2 the trend assessment of the key source analysis.

Table A1. Key source analysis – level assessment

		1990	2005	Level assessment	Cumulative total
Mobile Combustion: Fishing	CO ₂	655,5	702,9	0,19	0,19
Mobile combustion: Road vehicles	CO ₂	509,0	673,2	0,19	0,38
CO ₂ emissions from aluminium production	CO ₂	136,5	408,7	0,11	0,49
CO ₂ emissions from ferroalloys production	CO ₂	203,5	371,3	0,10	0,60
Mobile combustion: Construction	CO ₂	120,7	254,2	0,07	0,67
CH ₄ emissions from enteric fermentation in domestic livestock	CH ₄	269,9	231,5	0,06	0,73
CO ₂ emissions from stationary combustion, oil	CO ₂	237,0	168,0	0,05	0,78
CH ₄ emissions from solid waste disposal sites	CH ₄	113,6	154,0	0,04	0,82
CO ₂ emissions from geothermal energy	CO ₂	66,6	123,4	0,03	0,86
Direct N ₂ O emissions from agricultural soils	N ₂ O	142,1	116,6	0,03	0,89
Indirect N ₂ O emissions from Nitrogen used in agriculture	N ₂ O	99,8	80,9	0,02	0,91
Emissions from Substitutes for Ozone Depleting Substances	HFC	NO	76,7	0,02	0,93
CO ₂ emissions from Cement Production	CO ₂	51,6	53,9	0,02	0,95

Table A1. Key source analysis – trend assessment

		1990	2005	Level assessment	Trend assessment	Contribution to trend	Cumulative total
PFC emissions from aluminium production	PFC	419,6	26,1	0,01	0,110	0,275	0,28
CO ₂ emissions from aluminium production	CO ₂	136,5	408,7	0,11	0,063	0,158	0,43
CO ₂ emissions from ferroalloys production	CO ₂	203,5	371,3	0,10	0,035	0,088	0,52
Mobile combustion: Construction	CO ₂	120,7	254,2	0,07	0,029	0,073	0,60
Mobile combustion: Road vehicles	CO ₂	509,0	673,2	0,19	0,025	0,063	0,66
CO ₂ emissions from stationary combustion, oil	CO ₂	237,0	168,0	0,05	0,024	0,061	0,72
Emissions from Substitutes for Ozone Depleting Substances	HFC	NO	76,7	0,02	0,019	0,047	0,77
CH ₄ emissions from enteric fermentation in domestic livestock	CH ₄	269,9	231,5	0,06	0,018	0,044	0,81
CO ₂ emissions from geothermal energy	CO ₂	66,6	123,4	0,03	0,012	0,030	0,84
Direct N ₂ O emissions from agricultural soils	N ₂ O	142,1	116,6	0,03	0,011	0,027	0,87
Mobile combustion: Fishing	CO ₂	655,5	702,9	0,19	0,008	0,021	0,89
Indirect N ₂ O emissions from Nitrogen used in agriculture	N ₂ O	99,8	80,9	0,02	0,008	0,019	0,91
Mobile combustion: Road vehicles	N ₂ O	4,4	35,6	0,01	0,008	0,019	0,93
CO ₂ emissions from stationary combustion, coal	CO ₂	48,3	25,7	0,01	0,007	0,018	0,94
CH ₄ emissions from solid waste disposal sites	CH ₄	113,6	154,0	0,04	0,007	0,016	0,96

National Inventory Report 2007 - ICELAND

ANNEX II: QUANTITATIVE UNCERTAINTY

TIER 1 UNCERTAINTY CALCULATION AND REPORTING OF SOURCES IN ICELAND

Input Data						Uncertainty of Emissions		Uncertainty of Trend				
IPCC Source Category	Gas	Base year emissions (1990)	Year t emissions (2005)	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combine uncertainty as % of total national emissions in year 2005	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by EF unc.	Uncertainty in trend in national emissions introduced by a.d.	Uncertainty introduced into the trend in total national emissions
		Gg CO ₂ equivalents		%	%	%	%	%	%	%	%	%
1.A.3.b Transport - Road Transportation	CO ₂	509,02	673,19	10,0	5,0	11,18	2,03	0,033	0,201	0,16	2,84	2,84
Mobile Combustion - Construction industry	CO ₂	120,67	254,18	10,0	5,0	11,18	0,77	0,036	0,076	0,18	1,07	1,09
Mobile Combustion - Fishing	CO ₂	655,49	702,87	2,0	5,0	5,39	1,02	-0,006	0,210	-0,03	0,59	0,59
Stationary Combustion - Oil	CO ₂	237,00	168,00	5,0	5,0	7,07	0,32	-0,028	0,050	-0,14	0,35	0,38
Stationary Combustion - Coal	CO ₂	48,27	25,73	5,0	10,0	11,18	0,08	-0,008	0,008	-0,08	0,05	0,10
2.A.1 Cement Production	CO ₂	51,56	53,95		6,5	6,50	0,09	-0,001	0,016	-0,01	0,00	0,01
2.C.2 Ferroalloys Production	CO ₂	203,47	371,31	5,0	10,0	11,18	1,12	0,044	0,111	0,44	0,78	0,90
2.C.3 Aluminium Production	CO ₂	136,49	408,69	5,0	10,0	11,18	1,23	0,077	0,122	0,77	0,86	1,16
4.A Enteric Fermentation	CH ₄	269,92	231,49	20,0	50,0	53,85	3,36	-0,020	0,069	-1,00	1,95	2,19
6.A Solid Waste Disposal on Land	CH ₄	113,57	157,03	15,0	50,0	52,20	2,21	0,009	0,047	0,47	0,99	1,10
1.A.3.b Transport - Road Transportation	N ₂ O	4,39	35,58	50,0	200,0	206,16	1,98	0,009	0,011	1,83	0,75	1,98
Direct emissions from agricultural soil	N ₂ O	142,15	116,63	20,0	100,0	101,98	3,21	-0,012	0,035	-1,21	0,98	1,56
Indirect emissions from Nitrogen used in agriculture	N ₂ O	99,79	80,94	20,0	100,0	101,98	2,23	-0,009	0,024	-0,88	0,68	1,11
2.C.3 Aluminium Production												
CF ₄	PFC	355,02	7,09	5,0	7,0	8,60	0,02	-0,115	0,002	-0,80	0,01	0,80
C ₂ F ₆	PFC	64,61	1,29	5,0	22,0	22,56	0,01	-0,021	0,000	-0,46	0,00	0,46
Substitutes for Ozone Depleting Substances	HFC	0,00	76,74		100,0	100,00	2,07	0,023	0,023	2,29	0,00	2,29
Other non-key source emissions	All	340,79	340,77		30,0	30,00	2,76	-0,011	0,102	-0,32	0,00	0,32
Total emissions (all sources):		3.352,23	3.705,47				Total H :	7,5	Level Uncertainty		Total M :	5,6

National Inventory Report 2007 - ICELAND

Annex III: CRF TABLES FOR YEAR 2005

TABLE 1 SECTORAL REPORT FOR ENERGY
(Sheet 1 of 2)

Inventory 2005
Submission 2007 v1.1
ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	CO ₂	CH ₄	N ₂ O	NO _x	CO	NM VOC	SO ₂
	(Gg)						
Total Energy	1,913,62	0,16	0,24	26,65	19,35	4,06	2,48
A. Fuel Combustion Activities (Sectoral Approach)	1,913,62	0,16	0,24	26,65	19,35	4,06	2,48
1. Energy Industries	22,60	0,00	0,00	0,24	0,07	0,00	0,05
a. Public Electricity and Heat Production	22,60	0,00	0,00	0,24	0,07	0,00	0,05
b. Petroleum Refining	NO	NO	NO	NO	NO	NO	NO
c. Manufacture of Solid Fuels and Other Energy Industries	NO	NO	NO	NO	NO	NO	NO
2. Manufacturing Industries and Construction	442,28	0,02	0,11	4,52	1,35	0,59	1,23
a. Iron and Steel	1,45	0,00	0,00	0,00	0,00	0,00	0,00
b. Non-Ferrous Metals	19,00	0,00	0,00	0,05	0,00	0,00	0,02
c. Chemicals	NO	NO	NO	NO	NO	NO	NO
d. Pulp, Paper and Print	NO	NO	NO	NO	NO	NO	NO
e. Food Processing, Beverages and Tobacco	99,09	0,00	0,00	0,22	0,02	0,01	1,16
f. Other (as specified in table 1.A(a) sheet 2)	322,74	0,01	0,10	4,25	1,33	0,58	0,05
Mineral industry	55,73	0,00	0,00	0,30	0,05	0,01	0,00
Construction	254,18	0,01	0,10	3,92	1,28	0,57	0,01
Other non-specified	12,83	0,00	0,00	0,03	0,00	0,00	0,04
3. Transport	716,38	0,08	0,12	4,66	16,21	2,97	0,10
a. Civil Aviation	24,23	0,00	0,00	0,10	0,03	0,02	0,03
b. Road Transportation	673,19	0,07	0,11	4,09	16,13	2,94	0,02
c. Railways	NA,NO	NA,NO	NA,NO	NO	NO	NO	NO
d. Navigation	18,96	0,00	0,00	0,46	0,05	0,01	0,06
e. Other Transportation (as specified in table 1.A(a) sheet 3)	NA,NO	NA,NO	NA,NO				
Other non-specified	NA,NO	NA,NO	NA,NO				

National Inventory Report 2007 - ICELAND

TABLE 1 SECTORAL REPORT FOR ENERGY
(Sheet 2 of 2)

Inventory 2005
Submission 2007 v1.1
ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	CO ₂	CH ₄	N ₂ O	NO _x	CO	NM VOC	SO ₂
	(Gg)						
4. Other Sectors	716,26	0,07	0,02	17,18	1,72	0,50	0,93
a. Commercial/Institutional	1,54	0,00	NA,NO	0,00	0,00		0,00
b. Residential	11,86	0,00	0,00	0,01	0,00	0,00	0,01
c. Agriculture/Forestry/Fisheries	702,87	0,07	0,02	17,17	1,72	0,50	0,92
5. Other (as specified in table 1.A(a) sheet 4)	16,11	0,00	0,00	0,04	0,00	0,00	0,16
a. Stationary	16,11	0,00	0,00	0,04	0,00	0,00	0,16
Other non-specified	16,11	0,00	0,00	0,04	0,00	0,00	0,16
b. Mobile	NA	NA	NA	NA	NA	NA	NA
B. Fugitive Emissions from Fuels	NA,NE,NO	NA,NE,NO	NA,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
1. Solid Fuels	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO
a. Coal Mining and Handling	NO	NO	NO	NO	NO	NO	NO
b. Solid Fuel Transformation	NO	NO	NO	NO	NO	NO	NO
c. Other (as specified in table 1.B.1)	NA	NA	NA	NA	NA	NA	NA
2. Oil and Natural Gas	NA,NE,NO	NA,NE,NO	NA,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
a. Oil	NE,NO	NE,NO		NE	NE	NE	NE
b. Natural Gas	NO	NO				NO	NO
c. Venting and Flaring	NO	NO	NO	NO	NO	NO	NO
Venting	NO	NO				NO	NO
Flaring	NO	NO	NO	NO	NO	NO	NO
d. Other (as specified in table 1.B.2)	NA	NA	NA	NA	NA	NA	NA
Memo Items: ⁽¹⁾							
International Bunkers	407,16	0,00	0,01				
Aviation	407,16	0,00	0,01				
Marine	NA,NE,NO	NA,NE,NO	NA,NE,NO				
Multilateral Operations	NO	NO	NO	NO	NO	NO	NO
CO₂ Emissions from Biomass	NA,NO						

National Inventory Report 2007 - ICELAND

TABLE 1.A(a) SECTORAL BACKGROUND DATA FOR ENERGY
Fuel Combustion Activities - Sectoral Approach
(Sheet 1 of 4)

Inventory 2005
 Submission 2007 v1.1
 ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	AGGREGATE ACTIVITY DATA		IMPLIED EMISSION FACTORS ⁽²⁾			EMISSIONS		
	Consumption		CO ₂	CH ₄	N ₂ O	CO ₂	CH ₄	N ₂ O
	(TJ)	NCV/GCV ⁽¹⁾	(t/TJ)	(kg/TJ)		(Gg)		
I.A. Fuel Combustion	26.326,05	NCV				1.913,62	0,16	0,24
Liquid Fuels	25.866,40	NCV	72,59	6,21	9,30	1.877,74	0,16	0,24
Solid Fuels	277,56	NCV	92,71	1,00	1,40	25,73	0,00	0,00
Gaseous Fuels	NA,NO	NCV	NA,NO	NA,NO	NA,NO	0,11	NA,NO	NA,NO
Biomass	NA,NO	NCV	NA,NO	NA,NO	NA,NO ⁽³⁾		NA,NO	NA,NO
Other Fuels	182,09	NCV	55,15	NA,NE,NO	9,35	10,04	NA,NE,NO	0,00
I.A.1. Energy Industries	352,97	NCV				22,60	0,00	0,00
Liquid Fuels	170,88	NCV	73,47	3,86	0,59	12,55	0,00	0,00
Solid Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Gaseous Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Biomass	NO	NCV	NO	NO	NO ⁽³⁾	NO	NO	NO
Other Fuels	182,09	NCV	55,15	NE,NO	9,35	10,04	NE,NO	0,00
a. Public Electricity and Heat Production	352,97	NCV				22,60	0,00	0,00
Liquid Fuels	170,88	NCV	73,47	3,86	0,59	12,55	0,00	0,00
Solid Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Gaseous Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Biomass	NO	NCV	NO	NO	NO ⁽³⁾	NO	NO	NO
Other Fuels	182,09	NCV	55,15	NE	9,35	10,04	NE	0,00
b. Petroleum Refining	NO	NCV				NO	NO	NO
Liquid Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Solid Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Gaseous Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Biomass	NO	NCV	NO	NO	NO ⁽³⁾	NO	NO	NO
Other Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
c. Manufacture of Solid Fuels and Other Energy Industries	NO	NCV				NO	NO	NO
Liquid Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Solid Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Gaseous Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Biomass	NO	NCV	NO	NO	NO ⁽³⁾	NO	NO	NO
Other Fuels	NO	NCV	NO	NO	NO	NO	NO	NO

Note: All footnotes for this table are given at the end of the table on sheet 4.

National Inventory Report 2007 - ICELAND

TABLE 1.A(a) SECTORAL BACKGROUND DATA FOR ENERGY
Fuel Combustion Activities - Sectoral Approach
 (Sheet 2 of 4)

Inventory 2005
 Submission 2007 v1.1
 ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	AGGREGATE ACTIVITY DATA		IMPLIED EMISSION FACTORS ⁽³⁾			EMISSIONS		
	Consumption		CO ₂	CH ₄	N ₂ O	CO ₂	CH ₄	N ₂ O
	(TJ)	NCV/GCV ⁽¹⁾	(t/TJ)	(kg/TJ)		(Gg)		
I.A.2 Manufacturing Industries and Construction	5 793,56	NCV				442,28	0,02	0,11
Liquid Fuels	5 515,99	NCV	75,50	3,16	19,07	416,44	0,02	0,11
Solid Fuels	277,56	NCV	92,71	1,00	1,40	25,73	0,00	0,00
Gaseous Fuels		NCV				0,11		
Biomass		NCV			(3)			
Other Fuels		NCV						
a. Iron and Steel	19,72	NCV				1,45	0,00	0,00
Liquid Fuels	19,72	NCV	73,33	2,00	0,60	1,45	0,00	0,00
Solid Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Gaseous Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Biomass	NO	NCV	NO	NO	NO ⁽³⁾	NO	NO	NO
Other Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
b. Non-Ferrous Metals	42,82	NCV				19,00	0,00	0,00
Liquid Fuels	42,82	NCV	443,81	10,87	3,26	19,00	0,00	0,00
Solid Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Gaseous Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Biomass	NO	NCV	NO	NO	NO ⁽³⁾	NO	NO	NO
Other Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
c. Chemicals	NO	NCV				NO	NO	NO
Liquid Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Solid Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Gaseous Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Biomass	NO	NCV	NO	NO	NO ⁽³⁾	NO	NO	NO
Other Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
d. Pulp, Paper and Print	NO	NCV				NO	NO	NO
Liquid Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Solid Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Gaseous Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Biomass	NO	NCV	NO	NO	NO ⁽³⁾	NO	NO	NO
Other Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
e. Food Processing, Beverages and Tobacco	1 293,76	NCV				99,09	0,00	0,00
Liquid Fuels	1 293,76	NCV	76,59	2,10	0,44	99,09	0,00	0,00
Solid Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Gaseous Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Biomass	NE	NCV	NE	NE	NE ⁽³⁾	NE	NE	NE
Other Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
f. Other (please specify)⁽⁴⁾	4 437,26	NCV				322,74	0,01	0,10
Mineral industry								
Liquid Fuels	298,72	NCV	100,43	1,02	1,28	30,00	0,00	0,00
Solid Fuels	277,56	NCV	92,71	1,00	1,40	25,73	0,00	0,00
Gaseous Fuels		NCV						
Biomass		NCV			(3)			
Other Fuels		NCV						
Construction								
Liquid Fuels	3 466,40	NCV	73,33	3,92	30,00	254,18	0,01	0,10
Solid Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Gaseous Fuels		NCV						
Biomass		NCV			(3)			
Other Fuels		NCV						
Other non-specified								
Liquid Fuels	394,58	NCV	32,24	0,82	0,25	12,72	0,00	0,00
Solid Fuels		NCV						
Gaseous Fuels		NCV				0,11		
Biomass	NO	NCV	NO	NO	NO ⁽³⁾	NO	NO	NO
Other Fuels	NO	NCV	NO	NO	NO	NO	NO	NO

Note: All footnotes for this table are given at the end of the table on sheet 4.

National Inventory Report 2007 - ICELAND

TABLE 1.A(a) SECTORAL BACKGROUND DATA FOR ENERGY
Fuel Combustion Activities - Sectoral Approach
 (Sheet 3 of 4)

Inventory 2005
 Submission 2007 v1.1
 ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	AGGREGATE ACTIVITY DATA		IMPLIED EMISSION FACTORS ⁽²⁾			EMISSIONS		
	Consumption		CO ₂	CH ₄	N ₂ O	CO ₂	CH ₄	N ₂ O
	(TJ)	NCV/GCV ⁽¹⁾	(t/TJ)	(kg/TJ)		(Gg)		
I.A.3 Transport	10,233,73	NCV				716,38	0,08	0,12
Liquid Fuels	10,233,73	NCV	70,00	7,35	11,33	716,38	0,08	0,12
Solid Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Gaseous Fuels	NA,NO	NCV	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO
Biomass	NA,NO	NCV	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO
Other Fuels	NA	NCV	NA	NA	NA ⁽³⁾	NA	NA	NA
a. Civil Aviation	343,49	NCV				24,23	0,00	0,00
Aviation Gasoline	40,28	NCV	68,61	0,50	2,00	2,76	0,00	0,00
Jet Kerosene	303,21	NCV	70,79	0,50	2,00	21,46	0,00	0,00
b. Road Transportation	9,633,40	NCV				673,19	0,07	0,11
Gasoline	7,033,60	NCV	68,61	8,92	15,30	482,55	0,06	0,11
Diesel Oil	2,599,80	NCV	73,33	4,06	2,74	190,63	0,01	0,01
Liquefied Petroleum Gases (LPG)	NO	NCV	NO	NO	NO	NO	NO	NO
Other Liquid Fuels (please specify)	NA	NCV				NA	NA	NA
Gaseous Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Biomass	NO	NCV	NO	NO	NO ⁽³⁾	NO	NO	NO
Other Fuels (please specify)	NA	NCV				NA	NA	NA
c. Railways	NA,NO	NCV				NA,NO	NA,NO	NA,NO
Liquid Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Solid Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Gaseous Fuels	NA	NCV	NA	NA	NA	NA	NA	NA
Other Fuels (please specify)	NA	NCV				NA	NA	NA
d. Navigation	256,84	NCV				18,96	0,00	0,00
Residual Oil (Residual Fuel Oil)	40,19	NCV	76,59	7,00	2,00	3,08	0,00	0,00
Gas/Diesel Oil	216,65	NCV	73,33	7,00	2,00	15,89	0,00	0,00
Gasoline	NO	NCV	NO	NO	NO	NO	NO	NO
Other Liquid Fuels (please specify)	NA	NCV				NA	NA	NA
Solid Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Gaseous Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Other Fuels (please specify)	NA	NCV				NA	NA	NA
e. Other Transportation (please specify)⁽⁵⁾	NA,NO	NCV				NA,NO	NA,NO	NA,NO
Other non-specified	NA,NO	NCV				NA,NO	NA,NO	NA,NO
Liquid Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Solid Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Gaseous Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Biomass	NA	NCV	NA	NA	NA ⁽³⁾	NA	NA	NA
Other Fuels	NA	NCV	NA	NA	NA	NA	NA	NA

Note: All footnotes for this table are given at the end of the table on sheet 4.

National Inventory Report 2007 - ICELAND

TABLE 1.A(a) SECTORAL BACKGROUND DATA FOR ENERGY
Fuel Combustion Activities - Sectoral Approach
 (Sheet 4 of 4)

Inventory 2005
 Submission 2007 v1.1
 ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	AGGREGATE ACTIVITY DATA		IMPLIED EMISSION FACTORS ⁽²⁾			EMISSIONS		
	Consumption		CO ₂	CH ₄	N ₂ O	CO ₂	CH ₄	N ₂ O
	(TJ)	NCV/GCV ⁽¹⁾	(t/TJ)	(kg/TJ)		(Gg)		
I.A.4 Other Sectors	9,733.67	NCV				716.26	0.07	0.02
Liquid Fuels	9,733.67	NCV	73.59	6.88	1.97	716.26	0.07	0.02
Solid Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Gaseous Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Biomass	NO	NCV	NO	NO	NO ⁽³⁾	NO	NO	NO
Other Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
a. Commercial/Institutional	24.60	NCV				1.54	0.00	NA,NO
Liquid Fuels	24.60	NCV	62.44	1.10	NA	1.54	0.00	NA
Solid Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Gaseous Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Biomass	NO	NCV	NO	NO	NO ⁽³⁾	NO	NO	NO
Other Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
b. Residential	168.31	NCV				11.86	0.00	0.00
Liquid Fuels	168.31	NCV	70.44	0.80	0.45	11.86	0.00	0.00
Solid Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Gaseous Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Biomass	NO	NCV	NO	NO	NO ⁽³⁾	NO	NO	NO
Other Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
c. Agriculture/Forestry/Fisheries	9,540.76	NCV				702.87	0.07	0.02
Liquid Fuels	9,540.76	NCV	73.67	7.00	2.00	702.87	0.07	0.02
Solid Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Gaseous Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Biomass	NO	NCV	NO	NO	NO ⁽³⁾	NO	NO	NO
Other Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
I.A.5 Other (Not specified elsewhere)⁽⁶⁾	212.13	NCV				16.11	0.00	0.00
a. Stationary (please specify)⁽⁷⁾	212.13	NCV				16.11	0.00	0.00
Other non-specified								
Liquid Fuels	212.13	NCV	75.93	2.00	0.60	16.11	0.00	0.00
Solid Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Gaseous Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
Biomass	NO	NCV	NO	NO	NO ⁽³⁾	NO	NO	NO
Other Fuels	NO	NCV	NO	NO	NO	NO	NO	NO
b. Mobile (please specify)⁽⁸⁾	NA	NCV				NA	NA	NA

- (1) If activity data are calculated using net calorific values (NCV) as specified by the IPCC Guidelines, write NCV in this column. If gross calorific values (GCV) are used, write GCV in this column.
- (2) Accurate estimation of CH₄ and N₂O emissions depends on combustion conditions, technology and emission control policy, as well as on fuel characteristics. Therefore, caution should be used when comparing the implied emission factors across countries.
- (3) Although carbon dioxide emissions from biomass are reported in this table, they will not be included in the total CO₂ emissions from fuel combustion. The value for total CO₂ from biomass is recorded in Table1 sheet 2 under the Memo Items.
- (4) Use the cell below to list all activities covered under "f. Other".
- (5) Use the cell below to list all activities covered under "e. Other transportation".
- (6) Include military fuel use under this category.
- (7) Use the cell below to list all activities covered under "I.A.5.a Other - stationary".
- (8) Use the cell below to list all activities covered under "I.A.5.b Other - mobile".

Documentation Box:

- Parties should provide detailed explanations on the fuel combustion sub-sector in the corresponding part of Chapter 3: Energy (CRF sub-sector 1.A) of the NIR. Use this documentation box to provide references to relevant sections of the NIR if any additional information and/or further details are needed to understand the content of this table.
- If estimates are based on GCV, use this documentation box to provide reference to the relevant section of the NIR where the information necessary to allow the calculation of the activity data based on NCV can be found.
- If some derived gases (e.g. gas works gas, coke oven gas, blast furnace gas) are considered, use this documentation box to provide a reference to the relevant section of the NIR containing the information on the allocation of these derived gases under the above fuel categories (liquid, solid, gaseous, biomass and other fuels).

I.A.A Fuel Combustion - Sectoral Approach: I.A.2f Other manufacturing industries & construction includes: mineral industry, construction and other industries not included above.

National Inventory Report 2007 - ICELAND

TABLE 1.A(b) SECTORAL BACKGROUND DATA FOR ENERGY
CO₂ from Fuel Combustion Activities - Reference Approach (IPCC Worksheet 1-1)
 (Sheet 1 of 1)

Inventory 2005
 Submission 2007 v1.1
 ICELAND

FUEL TYPES			Unit	Production	Imports	Exports	International bunkers	Stock change	Apparent consumption	Conversion factor (TJ/Unit)	NCV/GCV ⁽¹⁾	Apparent consumption (TJ)	Carbon emission factor (t C/TJ)	Carbon content (Gg C)	Carbon stored (Gg C)	Net carbon emissions (Gg C)	Fraction of carbon oxidized	Actual CO ₂ emissions (Gg CO ₂)	
Liquid Fossil	Primary Fuels	Crude Oil		NA	NA	NA		NA	NA	NA	NCV	NA	NA	NA	NA	NA	NA	NA	
		Orimulsion		NA	NA	NA		NA	NA	NA	NCV	NA	NA	NA	NA	NA	NA	NA	NA
		Natural Gas Liquids		NA	NA	NA		NA	NA	NA	NCV	NA	NA	NA	NA	NA	NA	NA	NA
	Secondary Fuels	Gasoline	Gg		164,17	NA	NA	6,32	157,85	44,80	NCV	7,071,68	18,90	133,65	NA	133,65	0,99	485,17	
		Jet Kerosene			139,36	NA	129,00	2,86	7,50	44,59	NCV	334,43	19,50	6,52	NA	6,52	0,99	23,67	
		Other Kerosene	Gg		0,01	NA	NA	-0,14	0,15	44,75	NCV	6,71	19,60	0,13	NA	0,13	0,99	0,48	
		Shale Oil			NA	NA	NA	NA	NA	NA	NCV	NA	NA	NA	NA	NA	NA	NA	NA
		Gas / Diesel Oil	Gg		418,23	NA	55,40	4,33	358,50	43,33	NCV	15,533,81	20,20	313,78	NA	313,78	0,99	1,139,03	
		Residual Fuel Oil	Gg		62,90	NA	1,56	-0,65	61,99	40,19	NCV	2,491,38	21,10	52,57	NA	52,57	0,99	190,82	
		Liquefied Petroleum Gas (LPG)	Gg		2,46	NA	NA	0,14	2,32	47,31	NCV	109,66	17,20	1,89	NA	1,89	0,99	6,85	
		Ethane			NA	NA	NA	NA	NA	NA	NCV	NA	NA	NA	NA	NA	NA	NA	NA
		Naphtha			NA	NA	NA	NA	NA	NA	NCV	NA	NA	NA	NA	NA	NA	NA	NA
		Bitumen	Gg		41,39	NA	NA	NA	41,39	40,19	NCV	1,663,50	22,00	36,60	36,60	0,00	0,99	0,00	
		Lubricants	Gg		6,12	NA	NA	NA	6,12	40,19	NCV	246,03	20,00	4,92	2,44	2,49	0,99	9,02	
		Petroleum Coke	Gg		143,25	21,75	NA	NA	121,50	31,00	NCV	3,766,38	27,50	103,58	103,58	0,00	0,99	0,00	
		Refinery Feedstocks			NA	NA	NA	NA	NA	NA	NCV	NA	NA	NA	NA	NA	NA	NA	NA
Other Oil			NA	NA	NA	NA	NA	NA	NCV	NA	NA	NA	NA	NA	NA	NA	NA		
Other Liquid Fossil												NA	NA	NA	NA	NA	NA	NA	
Liquid Fossil Totals												31,223,57	653,64	142,61	511,03			1,855,04	
Solid Fossil	Primary Fuels	Anthracite ⁽²⁾		NA	NA	NA	NA	NA	NA	NCV	NA	NA	NA	NA	NA	NA	NA	NA	
		Coking Coal	Gg	NA	13,56	NA	NA	3,64	9,91	28,00	NCV	277,56	25,80	7,16	NA	7,16	0,98	25,73	
		Other Bituminous Coal	Gg	NA	103,40	NA	NA	16,53	86,87	28,00	NCV	2,432,30	25,80	62,75	62,75	0,00	0,98	0,00	
		Sub-bituminous Coal			NA	NA	NA	NA	NA	NA	NCV	NA	NA	NA	NA	NA	NA	NA	NA
		Lignite			NA	NA	NA	NA	NA	NA	NCV	NA	NA	NA	NA	NA	NA	NA	NA
		Oil Shale			NA	NA	NA	NA	NA	NA	NCV	NA	NA	NA	NA	NA	NA	NA	NA
		Peat			NA	NA	NA	NA	NA	NA	NCV	NA	NA	NA	NA	NA	NA	NA	NA
	Secondary Fuels	BKB ⁽³⁾ and Patent Fuel			NA	NA	NA	NA	NA	NA	NCV	NA	NA	NA	NA	NA	NA	NA	NA
		Coke Oven/Gas Coke	Gg		33,36	NA	NA	-9,23	42,59	28,00	NCV	1,192,63	29,50	35,18	35,18		0,98		
		Other Solid Fossil											NA	NA	NA	NA	NA	NA	NA
Solid Fossil Totals												3,902,50	105,10	97,94	7,16			25,73	
Gaseous Fossil												NA	NA	NA	NA	NA	NA	NA	
Gaseous Fossil (Dry)												NA	NA	NA	NA	NA	NA	NA	
Other Gaseous Fossil												NA	NA	NA	NA	NA	NA	NA	
Gaseous Fossil Totals												NA	NA	NA	NA	NA	NA	NA	
Total												35,126,07	758,73	240,54	518,19			1,880,77	
Biomass total												NA	NA	NA	NA	NA	NA	NA	
	Solid Biomass	Solid Biomass	Gg	NA	NA	NA	NA	NA	NA	16,72	NCV	NA	20,93	NA	NA	NA	0,98	NA	
		Liquid Biomass			NA	NA	NA	NA	NA	NA	NCV	NA	NA	NA	NA	NA	NA	NA	
		Gas Biomass			NA	NA	NA	NA	NA	NA	NCV	NA	NA	NA	NA	NA	NA	NA	

⁽¹⁾ To convert quantities in previous columns to energy units, use net calorific values (NCV) and write NCV in this column. If gross calorific values (GCV) are used, write GCV in this column.

⁽²⁾ If data for Anthracite are not available separately, include with Other Bituminous Coal.

⁽³⁾ BKB: Brown coal/peat briquettes.

Documentation Box:

Parties should provide detailed explanations on the fuel combustion sub-sector, including information relating to CO₂ from the Reference approach, in the corresponding part of Chapter 3: Energy (CRF sub-sector 1.A) of the NIR. Use this documentation box to provide references to relevant sections of the NIR if any additional information and/or further details are needed to understand the content of this table.

National Inventory Report 2007 - ICELAND

Inventory 2005
Submission 2007 v1.1
ICELAND

TABLE 1.A(c) COMPARISON OF CO₂ EMISSIONS FROM FUEL COMBUSTION
(Sheet 1 of 1)

FUEL TYPES	REFERENCE APPROACH			SECTORAL APPROACH ⁽¹⁾		DIFFERENCE ⁽²⁾	
	Apparent energy consumption ⁽³⁾ (PJ)	Apparent energy consumption (excluding non-energy use and feedstocks) ⁽⁴⁾ (PJ)	CO ₂ emissions (Gg)	Energy consumption (PJ)	CO ₂ emissions (Gg)	Energy consumption (%)	CO ₂ emissions (%)
Liquid Fuels (excluding international bunkers)	31,22	27,41	1.855,04	25,87	1.877,74	5,99	-1,21
Solid Fuels (excluding international bunkers) ⁽⁵⁾	3,90	0,28	25,73	0,28	25,73	0,00	0,00
Gaseous Fuels	NA	NA	NA	NA,NO	0,11		-100,00
Other ⁽⁵⁾	NA	NA	NA	0,18	10,04	-100,00	-100,00
Total ⁽⁵⁾	35,13	27,69	1.880,77	26,33	1.913,62	5,19	-1,72

⁽¹⁾ "Sectoral approach" is used to indicate the approach (if different from the Reference approach) used by the Party to estimate CO₂ emissions from fuel combustion as reported in table 1.A(a), sheets 1-4.

⁽²⁾ Difference in CO₂ emissions estimated by the Reference approach (RA) and the Sectoral approach (SA) (difference = 100% x ((RA-SA)/SA)). For calculating the difference in energy consumption between the two approaches, data as reported in the column "Apparent energy consumption (excluding non-energy use and feedstocks)" are used for the Reference approach.

⁽³⁾ Apparent energy consumption data shown in this column are as in table 1.A(b).

⁽⁴⁾ For the purposes of comparing apparent energy consumption from the Reference approach with energy consumption from the Sectoral approach, Parties should, in this column, subtract from the apparent energy consumption (Reference approach) the energy content corresponding to the fuel quantities used as feedstocks and/or for non-energy purposes, in accordance with the accounting of energy use in the Sectoral approach

⁽⁵⁾ Emissions from biomass are not included.

Note: The Reporting Instructions of the Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories require that estimates of CO₂ emissions from fuel combustion, derived using a detailed Sectoral approach, be compared to those from the Reference approach (Worksheet 1-1 of the IPCC Guidelines, Volume 2, Workbook). This comparison is to assist in verifying the Sectoral data.

Documentation Box:

Parties should provide detailed explanations on the fuel combustion sub-sector, including information related to the comparison of CO₂ emissions calculated using the Sectoral approach with those calculated using the Reference approach, in the corresponding part of Chapter 3: Energy (CRF sub-sector 1.A) of the NIR. Use this documentation box to provide references to relevant sections of the NIR if any additional information and/or further details are needed to understand the content of this table.

If the CO₂ emission estimates from the two approaches differ by more than 2 per cent, Parties should briefly explain the cause of this difference in this documentation box and provide a reference to relevant section of the NIR where this difference is explained in more detail.

1.AA Fuel Combustion - Sectoral Approach:1A2f Other manufacturing industries & construction includes: mineral industry, construction and other industries not included above.

National Inventory Report 2007 - ICELAND

Inventory 2005
Submission 2007 v1.1
ICELAND

**TABLE 1.A(d) SECTORAL BACKGROUND DATA FOR ENERGY
Feedstocks and Non-Energy Use of Fuels
(Sheet 1 of 1)**

FUEL TYPE	ACTIVITY DATA AND RELATED INFORMATION		IMPLIED EMISSION FACTOR	ESTIMATE
	Fuel quantity (TJ)	Fraction of carbon stored	Carbon emission factor (t C/TJ)	Carbon stored in non-energy use of fuels (Gg C)
Naphtha ⁽¹⁾		0,00	NA	NA
Lubricants	243,52	0,50	20,00	2,44
Bitumen	1.663,50	1,00	22,00	36,60
Coal Oils and Tars (from Coking Coal)		0,00	NA	NA
Natural Gas ⁽¹⁾		0,00	NA	NA
Gas/Diesel Oil ⁽¹⁾		0,00	NA	NA
LPG ⁽¹⁾		0,00	NA	NA
Ethane ⁽¹⁾		0,00	NA	NA
Other (please specify)				201,52
Coke oven/gas coke	1.192,63	1,00	29,50	35,18
Other Bituminous Coal	2.432,30	1,00	25,80	62,75
Petroleum Coke	3.766,60	1,00	27,50	103,58
Total				240,58
Total amount of C and CO₂ from feedstocks and non-energy use of fuels that is included as emitted CO₂ in the Reference approach				2,44

⁽¹⁾ Enter data for those fuels that are used as feedstocks (fuel used as raw materials for manufacture of products such as plastics or fertilizers) or for other non-energy use (fuels not used as fuel or transformed into another fuel (e.g. bitumen for road construction, lubricants)).

Documentation box:

- Parties should provide detailed explanations on the fuel combustion sub-sector, including information related to feedstocks, in the corresponding part of Chapter 3: Energy (CRF sub-sector 1.A) of the NIR. Use this documentation box to provide references to relevant sections of the NIR if any additional information and/or further details are needed to understand the content of this table.
- The above table is consistent with the IPCC Guidelines. Parties that take into account the emissions associated with the use and disposal of these feedstocks could continue to use their methodology, but should indicate this in this documentation box and provide a reference to the relevant section of the NIR where further explanation can be found.

Additional information ^(a)

CO ₂ not emitted (Gg CO ₂)	Subtracted from energy sector (specify source category)	Associated CO ₂ emissions (Gg)	Allocated under (Specify source category, e.g. Waste Incineration)
NA			
8,93	NE		
134,19	NE		
NA			
129,00			
230,10			
379,80			
882,02			
8,93			

^(a) The fuel lines continue from the table to the left.

A fraction of energy carriers is stored in such products as plastics or asphalt. The non-stored fraction of the carbon in the energy carrier or product is oxidized, resulting in carbon dioxide emissions, either during use of the energy carriers in the industrial production (e.g. fertilizer production), or during use of the products (e.g. solvents, lubricants), or in both (e.g. monomers). To report associated emissions, use the above table.

National Inventory Report 2007 - ICELAND

Inventory 2005
Submission 2007 v1.1
ICELAND

TABLE 1.B.1 SECTORAL BACKGROUND DATA FOR ENERGY
Fugitive Emissions from Solid Fuels
(Sheet 1 of 1)

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	ACTIVITY DATA		IMPLIED EMISSION FACTORS		EMISSIONS		
	Amount of fuel produced	(Mt)	CH ₄ ⁽¹⁾	CO ₂	CH ₄		CO ₂
					Recovery/Flaring ⁽²⁾	Emissions ⁽³⁾	
		(kg/t)		(Gg)			
1. B. 1. a. Coal Mining and Handling	NO					NO	NO
i. Underground Mines ⁽⁴⁾	NO		NO	NO		NO	NO
Mining Activities			NO	NO		NO	NO
Post-Mining Activities			NO	NO		NO	NO
ii. Surface Mines ⁽⁴⁾	NO		NO	NO		NO	NO
Mining Activities			NO	NO		NO	NO
Post-Mining Activities			NO	NO		NO	NO
1. B. 1. b. Solid Fuel Transformation	NO		NO	NO		NO	NO
1. B. 1. c. Other (please specify)⁽⁵⁾					NA	NA	NA

⁽¹⁾ The IEFs for CH₄ are estimated on the basis of gross emissions as follows: (CH₄ emissions + amounts of CH₄ flared/recovered) / activity data.

⁽²⁾ Amounts of CH₄ drained (recovered), utilized or flared.

⁽³⁾ Final CH₄ emissions after subtracting the amounts of CH₄ utilized or recovered.

⁽⁴⁾ In accordance with the IPCC Guidelines, emissions from Mining Activities and Post-Mining Activities are calculated using the activity data of the amount of fuel produced for Underground Mines and Surface Mines.

⁽⁵⁾ This category is to be used for reporting any other solid-fuel-related activities resulting in fugitive emissions, such as emissions from abandoned mines and waste piles.

Note: There are no clear references to the coverage of 1.B.1.b. and 1.B.1.c. in the IPCC Guidelines. Make sure that the emissions entered here are not reported elsewhere. If they are reported under another source category, indicate this by using notation key IE and making the necessary reference in Table 9 (completeness).

Documentation box:

- Parties should provide detailed explanations on the fugitive emissions from source category 1.B.1 Solid Fuels, in the corresponding part of Chapter 3: Energy (CRF source category 1.B.1) of the NIR. Use this documentation box to provide
- Regarding data on the amount of fuel produced entered in the above table, specify in this documentation box whether the fuel amount is based on the run-of-mine (ROM) production or on the saleable production.
- If entries are made for "Recovery/Flaring", indicate in this documentation box whether CH₄ is flared or recovered and provide a reference to the section in the NIR where further details on recovery/flaring can be found.
- If estimates are reported under 1.B.1.b. and 1.B.1.c., use this documentation box to provide information regarding activities covered under these categories and to provide a reference to the section in the NIR where the background information can be found.

National Inventory Report 2007 - ICELAND

TABLE 1.B.2 SECTORAL BACKGROUND DATA FOR ENERGY
Fugitive Emissions from Oil, Natural Gas and Other Sources
 (Sheet 1 of 1)

Inventory 2005
 Submission 2007 v1.1
 ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	ACTIVITY DATA ⁽¹⁾			IMPLIED EMISSION FACTORS			EMISSIONS		
	Description ⁽¹⁾	Unit ⁽¹⁾	Value	CO ₂	CH ₄	N ₂ O	CO ₂	CH ₄	N ₂ O
				(kg/unit) ⁽²⁾			(Gg)		
I. B. 2. a. Oil ⁽³⁾							NE,NO	NE,NO	
i. Exploration	<i>(specify)</i>		NO	NO	NO	NO	NO	NO	
ii. Production ⁽⁴⁾	<i>(specify)</i>		NO	NO	NO		NO	NO	
iii. Transport	<i>(specify)</i>		NO	NO	NO		NO	NO	
iv. Refining / Storage	<i>(specify)</i>		NO	NO	NO	NO	NO	NO	
v. Distribution of Oil Products	<i>(specify)</i>		NE	NE	NE		NE	NE	
vi. Other	<i>(specify)</i>		NO	NO	NO		NO	NO	
I. B. 2. b. Natural Gas							NO	NO	
i. Exploration	<i>(specify)</i>		NO	NO	NO		NO	NO	
ii. Production ⁽⁴⁾ / Processing	<i>(specify)</i>		NO	NO	NO		NO	NO	
iii. Transmission	<i>(specify)</i>		NO	NO	NO		NO	NO	
iv. Distribution	<i>(specify)</i>		NO	NO	NO		NO	NO	
v. Other Leakage	<i>(specify)</i>		NO	NO	NO		NO	NO	
<i>at industrial plants and power stations</i>	<i>(specify)</i>		NO	NO	NO		NO	NO	
<i>in residential and commercial sectors</i>	<i>(specify)</i>		NO	NO	NO		NO	NO	
I. B. 2. c. Venting ⁽⁵⁾							NO	NO	
i. Oil	<i>(specify)</i>		NO	NO	NO		NO	NO	
ii. Gas	<i>(specify)</i>		NO	NO	NO		NO	NO	
iii. Combined	<i>(specify)</i>		NO	NO	NO		NO	NO	
Flaring							NO	NO	NO
i. Oil	<i>(specify)</i>		NO	NO	NO	NO	NO	NO	NO
ii. Gas	<i>(specify)</i>		NO	NO	NO	NO	NO	NO	NO
iii. Combined	<i>(specify)</i>		NO	NO	NO	NO	NO	NO	NO
I.B.2.d. Other <i>(please specify)</i> ⁽⁶⁾							NA	NA	NA

⁽¹⁾ Specify the activity data used in the Description column (see examples). Specify the unit of the activity data in the Unit column using one of the following units: PJ, Tg, 10⁶ m³, 10⁶ bbl/yr, km, number of sources (e.g. wells).

National Inventory Report 2007 - ICELAND

Inventory 2005
Submission 2007 v1.1
ICELAND

TABLE 1.C SECTORAL BACKGROUND DATA FOR ENERGY
International Bunkers and Multilateral Operations
(Sheet 1 of 1)

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	ACTIVITY DATA Consumption (TJ)	IMPLIED EMISSION FACTORS			EMISSIONS		
		CO ₂	CH ₄	N ₂ O	CO ₂	CH ₄	N ₂ O
		(t/TJ)			(Gg)		
Aviation Bunkers	5,752,11				407,16	0,00	0,01
Jet Kerosene	5,752,11	70,79	0,00	0,00	407,16	0,00	0,01
Gasoline	NO	NO	NO	NO	NO	NO	NO
Marine Bunkers	NA,NE,NO				NA,NE,NO	NA,NE,NO	NA,NE,NO
Gasoline	NO	NO	NO	NO	NO	NO	NO
Gas/Diesel Oil							
Residual Fuel Oil							
Lubricants	NE	NE	NE	NE	NE	NE	NE
Coal	NO	NO	NO	NO	NO	NO	NO
Other (<i>please specify</i>)	NA				NA	NA	NA
Multilateral Operations ⁽¹⁾	NO	NO	NO	NO	NO	NO	NO

⁽¹⁾ Parties may choose to report or not report the activity data and implied emission factors for multilateral operations consistent with the principle of confidentiality stated in the UNFCCC. In any case, Parties should report the emissions from multilateral operations, where available, under the Memo Items section of the Summary tables and in the Sectoral report table for energy.

Note: In accordance with the IPCC Guidelines, international aviation and

Documentation box:

- Parties should provide detailed explanations on the fuel combustion sub-sector, including international bunker fuels, in the corresponding part of Chapter 3: Energy (CRF sub-sector 1.A) of the NIR. Use this documentation box to provide references to relevant sections of the NIR if any additional information and/or further details are needed to understand the content of this table.
- Provide in this documentation box a brief explanation on how the consumption of international marine and aviation bunker fuels was estimated and separated from domestic consumption, and include a reference to the section of the NIR where the explanation is provided in more detail.

Additional information

Fuel consumption	Distribution ^(a) (per cent)	
	Domestic	International
Aviation	5,64	94,36
Marine	100,00	NA,NE,NO

^(a) For calculating the allocation of fuel consumption, the sums of fuel consumption for domestic navigation and aviation (table 1.A(a)) and for international bunkers (table 1.C) are used.

National Inventory Report 2007 - ICELAND

 TABLE 2(I) SECTORAL REPORT FOR INDUSTRIAL PROCESSES
 (Sheet 1 of 2)

 Inventory 2005
 Submission 2007 v1.1
 ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	CO ₂	CH ₄	N ₂ O	HFCs ⁽¹⁾		PFCs ⁽¹⁾		SF ₆		NO _x	CO	NMVOC	SO ₂
				P	A	P	A	P	A				
	(Gg)			CO ₂ equivalent (Gg)						(Gg)			
Total Industrial Processes	835,11	0,05	NA,NE,NO	76,74	NA,NE,NO	NA,NE,NO	26,09	0,00	NA,NE,NO	1,18	0,23	0,10	5,68
A. Mineral Products	55,10	NA,NE	NA,NE							0,02	0,05	0,00	0,03
1. Cement Production	53,95												IE
2. Lime Production	NO												
3. Limestone and Dolomite Use	NO												
4. Soda Ash Production and Use	NO												
5. Asphalt Roofing	NO										NO	NO	
6. Road Paving with Asphalt	NE									0,02	0,02	0,00	0,02
7. Other (as specified in table 2(I).A-G)	1,15	NA,NE	NA,NE							NA,NE	0,02	NA,NE	0,00
Glass Production	NO	NA	NA							NA	NA	NA	NA
Mineral wool production	1,15	NE	NE							NE	0,02	NE	0,00
B. Chemical Industry		NE,NO	NE,NO	NA	NA	NA	NA	NA	NA		NE,NO	NE,NO	NE,NO
1. Ammonia Production	NO	NO	NO							NO	NO	NO	NO
2. Nitric Acid Production			NO							NO			
3. Adipic Acid Production			NO							NO	NO	NO	
4. Carbide Production	NO	NO									NO	NO	NO
5. Other (as specified in table 2(I).A-G)		NE,NO	NE,NO	NA	NA	NA	NA	NA	NA		NE,NO	NE,NO	NE,NO
Carbon Black		NO											
Ethylene	NO	NO	NO										
Dichloroethylene		NO											
Styrene		NO											
Methanol		NO											
Fertilizer production	NO	NO	NO	NA	NA	NA	NA	NA	NA	NO	NO	NO	NO
Silicon Production		NE	NE	NA	NA	NA	NA	NA	NA		NE	NE	NE
C. Metal Production	780,00	0,05	NA	NA	NA	NA	26,09	NA	NA,NO	1,17	0,18	0,09	5,65
1. Iron and Steel Production	NA,NO	NA,NO								NO	NO	NO	NO
2. Ferroalloys Production	371,31	0,05								1,17	0,18	0,09	2,22
3. Aluminium Production	408,69	NE					26,09			NE	NE	NE	3,43
4. SF ₆ Used in Aluminium and Magnesium Foundries									NO				
5. Other (as specified in table 2(I).A-G)	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

Note: P = Potential emissions based on Tier 1 approach of the IPCC Guidelines. A = Actual emissions based on Tier 2 approach of the IPCC Guidelines. This applies only to source categories where methods exist for both tiers.

National Inventory Report 2007 - ICELAND

TABLE 2(I) SECTORAL REPORT FOR INDUSTRIAL PROCESSES
(Sheet 2 of 2)

Inventory 2005
Submission 2007 v1.1
ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	CO ₂	CH ₄	N ₂ O	HFCs ⁽¹⁾		PFCs ⁽¹⁾		SF ₆		NO _x	CO	NMVOC	SO ₂
				P	A	P	A	P	A				
	(Gg)			CO ₂ equivalent (Gg)						(Gg)			
D. Other Production	NE									NO	NO	NE,NO	NO
1. Pulp and Paper										NO	NO	NO	NO
2. Food and Drink ⁽²⁾	NE											NE	
E. Production of Halocarbons and SF₆					NA,NO		NA,NO		NA,NO				
1. By-product Emissions					NA,NO		NA,NO		NO				
Production of HCFC-22					NO								
Other					NA,NO		NA,NO		NO				
2. Fugitive Emissions					NA,NO		NA,NO		NO				
3. Other (as specified in table 2(II))					NA		NA		NA				
F. Consumption of Halocarbons and SF₆				76,74	NA,NE,NO	NE,NO	NA,NE,NO	0,00	NA,NE,NO				
1. Refrigeration and Air Conditioning Equipment				76,74	NE,NO	NO	NE,NO	NO	NE				
2. Foam Blowing				NO	NO	NO	NO	NO	NO				
3. Fire Extinguishers				NO	NO	NO	NO	NO	NO				
4. Aerosols/ Metered Dose Inhalers				NE	NE	NO	NE	NO	NE				
5. Solvents				NO	NO	NO	NO	NO	NO				
6. Other applications using ODS ⁽³⁾ substitutes													
7. Semiconductor Manufacture				NO	NO	NO	NO	NO	NO				
8. Electrical Equipment				NO		NO		0,00	NE				
9. Other (as specified in table 2(II))				NA	NA	NA	NA	NA	NA				
G. Other (as specified in tables 2(I),A-G and 2(II))	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

Note: P = Potential emissions based on Tier 1 approach of the IPCC Guidelines. A = Actual emissions based on Tier 2 approach of the IPCC Guidelines. This applies only to source categories where methods exist for both tiers.

⁽¹⁾ The emissions of HFCs and PFCs are to be expressed as CO₂ equivalent emissions. Data on disaggregated emissions of HFCs and PFCs are to be provided in Table 2(II).

⁽²⁾ CO₂ from Food and Drink Production (e.g. gasification of water) can be of biogenic or non-biogenic origin. Only information on CO₂ emissions of non-biogenic origin should be reported.

⁽³⁾ ODS: ozone-depleting substances.

Documentation box:

Parties should provide detailed explanations on the industrial processes sector in Chapter 4: Industrial processes (CRF sector 2) of the NIR. Use this documentation box to provide references to relevant sections of the NIR if any additional information and/or further details are needed to understand the content of this table.

National Inventory Report 2007 - ICELAND

TABLE 2(I).A-G SECTORAL BACKGROUND DATA FOR INDUSTRIAL PROCESSES
Emissions of CO₂, CH₄ and N₂O
 (Sheet 1 of 2)

Inventory 2005
 Submission 2007 v1.1
 ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	ACTIVITY DATA		IMPLIED EMISSION FACTORS ⁽²⁾			EMISSIONS					
	Production/Consumption quantity		CO ₂	CH ₄	N ₂ O	CO ₂		CH ₄		N ₂ O	
	Description ⁽¹⁾	(kt)				(t/t)	Emissions ⁽³⁾	Recovery ⁽⁴⁾	Emissions ⁽³⁾	Recovery ⁽⁴⁾	Emissions ⁽³⁾
			(Gg)								
A. Mineral Products						55,10	NA	NA,NE	NA	NA,NE	NA
1. Cement Production	clinker production	99,17	0,54			53,95					
2. Lime Production	(specify)	NO	NO			NO					
3. Limestone and Dolomite Use	(specify)	NO	NO			NO					
4. Soda Ash						NO					
Soda Ash Production	(specify)	NO	NO			NO					
Soda Ash Use	(specify)	NO	NO			NO					
5. Asphalt Roofing	(specify)	NO	NO			NO					
6. Road Paving with Asphalt	asphalt production	201,60	NE			NE					
7. Other (please specify)						1,15	NA	NA,NE	NA	NA,NE	NA
Glass Production	(specify)	NO	NO	NA	NA	NO	NA	NA	NA	NA	NA
Mineral wool production	mineral wool production	9,30	0,12	NE	NE	1,15	NA	NE	NA	NE	NA
B. Chemical Industry							NA	NE,NO	NA	NE,NO	NA
1. Ammonia Production ⁽⁵⁾	(specify)	NO	NO	NO	NO	NO		NO		NO	
2. Nitric Acid Production	(specify)	NO			NO					NO	
3. Adipic Acid Production	(specify)	NO	NO		NO					NO	
4. Carbide Production	(specify)	NO	NO	NO		NO		NO			
Silicon Carbide	(specify)	NO	NO	NO		NO		NO			
Calcium Carbide	(specify)	NO	NO	NO		NO		NO			
5. Other (please specify)							NA	NE,NO	NA	NE,NO	NA
Carbon Black	(specify)	NO		NO				NO	NA		
Ethylene	(specify)	NO	NO	NO	NO	NO	NA	NO	NA	NO	NA
Dichloroethylene	(specify)	NO		NO				NO	NA		
Styrene	(specify)	NO		NO				NO	NA		
Methanol	(specify)	NO		NO				NO	NA		
Fertilizer production	fertilizer production	NO	NO	NO	NO	NO	NA	NO	NA	NO	NA
Silicon Production	Silicon Production			NE	NE		NA	NE	NA	NE	NA

⁽¹⁾ Where the IPCC Guidelines provide options for activity data, e.g. cement production or clinker production for estimating the emissions from Cement Production, specify the activity data used (as shown in the example in parentheses) in order to make the choice of emission factor more transparent and to facilitate comparisons of implied emission factors.

⁽²⁾ The implied emission factors (IEF) are estimated on the basis of gross emissions as follows: IEF = (emissions plus amounts recovered, oxidized, destroyed or transformed) / activity data.

⁽³⁾ Final emissions are to be reported (after subtracting the amounts of emission recovery, oxidation, destruction or transformation).

⁽⁴⁾ Amounts of emission recovery, oxidation, destruction or transformation.

⁽⁵⁾ To avoid double counting, make offsetting deductions for fuel consumption (e.g. natural gas) in Ammonia Production, first for feedstock use of the fuel, and then for a sequestering use of the feedstock.

National Inventory Report 2007 - ICELAND

TABLE 2(I).A-G SECTORAL BACKGROUND DATA FOR INDUSTRIAL PROCESSES
Emissions of CO₂, CH₄ and N₂O
 (Sheet 2 of 2)

Inventory 2005
 Submission 2007 v1.1
 ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	ACTIVITY DATA		IMPLIED EMISSION FACTORS ⁽²⁾			EMISSIONS					
	Production/Consumption quantity		CO ₂	CH ₄	N ₂ O	CO ₂		CH ₄		N ₂ O	
	Description ⁽¹⁾	(kt)				(t/t)	Emissions ⁽³⁾	Recovery ⁽⁴⁾	Emissions ⁽³⁾	Recovery ⁽⁴⁾	Emissions ⁽³⁾
			(Gg)								
C. Metal Production						780,00	NA	0,05	NA	NA	NA
1. Iron and Steel Production			NA,NO	NA,NO		NA,NO	NA	NA,NO	NA		
Steel	(specify)	NO	NO	NA		NO	NA	NA	NA		
Pig Iron	(specify)	NO	NO	NO		NO	NA	NO	NA		
Sinter	(specify)	NO	NO	NO		NO	NA	NO	NA		
Coke	(specify)	NO	NO	NO		NO	NA	NO	NA		
Other (please specify)						NA	NA	NA	NA		
2. Ferroalloys Production	Ferrosilicon - 75% Si	110,96	3,35	0,00		371,31		0,05			
3. Aluminium Production	Aluminium production	272,49	1,50	NE		408,69		NE			
4. SF ₆ Used in Aluminium and Magnesium Foundries											
5. Other (please specify)						NA	NA	NA	NA	NA	NA
D. Other Production						NE					
1. Pulp and Paper											
2. Food and Drink	(specify)	NE	NE			NE					
G. Other (please specify)						NA	NA	NA	NA	NA	NA

⁽¹⁾ Where the IPCC Guidelines provide options for activity data, e.g. cement production or clinker production for estimating the emissions from Cement Production, specify the activity data used (as shown in the example in parentheses) in order to make the choice of emission factor more transparent and to facilitate comparisons of implied emission factors.

⁽²⁾ The implied emission factors (IEF) are estimated on the basis of gross emissions as follows: IEF = (emissions + amounts recovered, oxidized, destroyed or transformed) / activity data.

⁽³⁾ Final emissions are to be reported (after subtracting the amounts of emission recovery, oxidation, destruction or transformation).

⁽⁴⁾ Amounts of emission recovery, oxidation, destruction or transformation.

Documentation box:

- Parties should provide detailed explanations on the industrial processes sector in Chapter 4: Industrial processes (CRF sector 2) of the NIR. Use this documentation box to provide references to relevant sections of the NIR if any additional information and/or further details are needed to understand the content of this table.
- In relation to metal production, more specific information (e.g. data on virgin and recycled steel production) could be provided in this documentation box, or in the NIR, together with a reference to the relevant section.
- Confidentiality: Where only aggregate figures for activity data are provided, e.g. due to reasons of confidentiality, a note indicating this should be provided in this documentation box.

National Inventory Report 2007 - ICELAND

TABLE 2(H) SECTORAL REPORT FOR INDUSTRIAL PROCESSES - EMISSIONS OF HFCs, PFCs AND SF₆
(Sheet 1 of 2)

Inventory 2005
Submission 2007 v.1.1
ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	HFC-23	HFC-32	HFC-41	HFC-43-10mixe	HFC-125	HFC-134	HFC-134a	HFC-152a	HFC-143	HFC-143a	HFC-227ea	HFC-236fa	HFC-245ea	Unspecified mix of listed HFCs ^(b)	Total HFCs	CF ₄	C ₂ F ₆	C ₃ F ₈	C ₄ F ₁₀	e-C ₆ F ₆	C ₆ F ₁₂	C ₆ F ₁₄	Unspecified mix of listed PFCs ^(b)	Total PFCs	SF ₆	
	(t) ^(a)														CO ₂ equivalent (Gg)	CO ₂ equivalent (Gg)	(t) ^(a)						CO ₂ equivalent (Gg)	CO ₂ equivalent (Gg)	(t) ^(a)	
Total Actual Emissions of Halocarbons (by chemical) and SF₆	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NO		3,40	0,44	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NO		NA,NE,NO
C. Metal Production																	3,40	0,44	NE	NE	NE	NE	NE			NA,NO
Aluminium Production																	3,40	0,44	NE	NE	NE	NE	NE			
SF ₆ Used in Aluminium Foundries																										NO
SF ₆ Used in Magnesium Foundries																										NO
E. Production of Halocarbons and SF₆	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA		NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA		NA,NO
1. By-product Emissions	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO			NO	NO	NO	NO	NO	NO	NO			NO
Production of HCFC-22	NO																									NO
Other	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO			NO	NO	NO	NO	NO	NO	NO			NO
2. Fugitive Emissions	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO			NO	NO	NO	NO	NO	NO	NO			NO
3. Other (as specified in table 2(H), C,E)	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA		NA	NA	NA	NA	NA	NA	NA	NA			NA
F(a). Consumption of Halocarbons and SF₆ (actual)	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NO		NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NO		NA,NE,NO
1. Refrigeration and Air Conditioning Equipment	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NO		NE	NE	NE	NE	NE	NE	NE	NO		NE
2. Foam Blowing	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO			NO	NO	NO	NO	NO	NO	NO			NO
3. Fire Extinguishers	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO			NO	NO	NO	NO	NO	NO	NO			NO
4. Aerosols/Metered Dose Inhalers	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE			NE	NE	NE	NE	NE	NE	NE			NE
5. Solvents	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO			NO	NO	NO	NO	NO	NO	NO			NO
6. Other applications using ODS ^(b) substitutes																										
7. Semiconductor Manufacture	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO			NO	NO	NO	NO	NO	NO	NO			NO
8. Electrical Equipment																										NE
9. Other (as specified in table 2(H),F)	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA		NA	NA	NA	NA	NA	NA	NA	NA			NA
G. Other (please specify)	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA		NA	NA	NA	NA	NA	NA	NA	NA			NA

Note: All footnotes for this table are given at the end of the table on sheet 2.

Note: Gases with global warming potential (GWP) values not yet agreed upon by the Conference of the Parties should be reported in table 9(b).

National Inventory Report 2007 - ICELAND

TABLE 2(II) SECTORAL REPORT FOR INDUSTRIAL PROCESSES - EMISSIONS OF HFCs, PFCs AND SF₆
(Sheet 2 of 2)

Inventory 2005
Submission 2007 v1.1
ICELAND

GREENHOUSE GAS SOURCE AND CATEGORIES	SINK														Unspecified mix of HFCs ⁽¹⁾	Total HFCs	CF ₄	C ₂ F ₆	C ₃ F ₈	C ₄ F ₁₀	C ₆ F ₁₄	C ₈ F ₁₈	C ₁₀ F ₂₂	C ₁₂ F ₂₆	C ₁₄ F ₃₀	Unspecified mix of HFCs ⁽¹⁾	Total PFCs	SF ₆
	HFC-23	HFC-32	HFC-41	HFC-43-1b-misc	HFC-125	HFC-134a	HFC-134b	HFC-152a	HFC-143a	HFC-143a	HFC-227ea	HFC-280fa	HFC-345ca	(t) ⁽²⁾														
Fip). Total Potential Emissions of Halocarbons (by chemical) and SF₆ ⁽⁴⁾	NE,NO	0,32	NE,NO	NE,NO	10,76	NE,NO	8,39	0,35	NE,NO	9,33	NE,NO	NE,NO	NE,NO			NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO			NE,NO			
Production ⁽⁵⁾	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO			NO	NO	NO	NO	NO	NO	NO			NO			
Import	NE,NO	0,32	NE,NO	NE,NO	10,76	NE,NO	8,39	0,35	NE,NO	9,33	NE,NO	NE,NO	NE,NO			NE,NO	NE	NE	NE,NO	NE,NO	NE,NO	NE,NO			NE,NO			
In bulk	NO	0,32	NO	NO	10,76	NO	8,39	0,35	NO	9,33	NO	NO	NO			NO	NE	NE	NO	NO	NO	NO			NO			
In products ⁽⁶⁾	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE			NE	NE	NE	NE	NE	NE	NE			NE			
Export	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE			NE	NE	NE	NE	NE	NE	NE			NE			
In bulk	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE			NE	NE	NE	NE	NE	NE	NE			NE			
In products ⁽⁶⁾	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE			NE	NE	NE	NE	NE	NE	NE			NE			
Destroyed amount	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO			NO	NO	NO	NO	NO	NO	NO			NO			
GWP values used	11700	650	150	1300	2800	1000	1300	140	300	3800	2900	6300	560			6500	9200	7000	7000	8700	7500	7400			23900			
Total Actual Emissions⁽⁷⁾ (CO₂ equivalent (Gg))	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NO	NA,NE,NO	22,08	4,02	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NO	26,09	NA,NE,NO			
C. Metal Production																22,08	4,02	NE	NE	NE	NE	NE			NA,NO			
E. Production of Halocarbons and SF₆	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA	NA,NO	NA,NO			
F(a). Consumption of Halocarbons and SF₆	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NO	NA,NO	NA,NE,NO			
G. Other	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA			
Ratio of Potential/Actual Emissions from Consumption of Halocarbons and SF₆																												
Actual emissions - F(a) (Gg CO₂ eq.)	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NO	NA,NE,NO	NA,NE,NO			
Potential emissions - F(p)⁽⁸⁾ (Gg CO₂ eq.)	NE,NO	0,21	NE,NO	NE,NO	30,12	NE,NO	10,90	0,05	NE,NO	35,46	NE,NO	NE,NO	NE,NO			76,74	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO			NE,NO			
Potential/Actual emissions ratio	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NO	NA,NE,NO	NA,NE,NO			

⁽¹⁾ In accordance with the UNFCCC reporting guidelines, HFC and PFC emissions should be reported for each relevant chemical. However, if it is not possible to report values for each chemical (i.e. mixtures, confidential data, lack of disaggregation), these columns could be used for reporting aggregate figures for HFCs and PFCs, respectively. Note that the unit used for these columns is Gg of CO₂ equivalent.

⁽²⁾ Note that the units used in this table differ from those used in the rest of the Sectoral report tables, i.e. t instead of Gg.

⁽³⁾ ODS: ozone-depleting substances

⁽⁴⁾ Potential emissions of each chemical of halocarbons and SF₆ estimated using Tier 1a or Tier 1b of the IPCC Guidelines (Volume 3. Reference Manual, pp. 2.47-2.50). Where potential emission estimates are available in a disaggregated manner for the source categories F.1 to F.9, these should be reported in the NIR and a reference should be provided in the documentation box. Use table Summary 3 to indicate whether Tier 1a or Tier 1b was used.

⁽⁵⁾ Production refers to production of new chemicals. Recycled substances could be included here, but avoid double counting of emissions. An indication as to whether recycled substances are included should be provided in the documentation box to this table.

⁽⁶⁾ Relevant only for Tier 1b.

⁽⁷⁾ Total actual emissions equal the sum of the actual emissions of each halocarbon and SF₆ from the source categories 2.C, 2.E, 2.F and 2.G as reported in sheet 1 of this table multiplied by the corresponding GWP values.

⁽⁸⁾ Potential emissions of each halocarbon and SF₆ taken from row Fip) multiplied by the corresponding GWP values.

Note: As stated in the UNFCCC reporting guidelines, Parties should report actual emissions of HFCs, PFCs and SF₆ where data are available, providing disaggregated data by chemical and source category in units of mass and in CO₂ equivalent. Parties reporting actual emissions should also report potential emissions for the sources where the concept of potential emissions applies, for reasons of transparency and comparability. Gases with GWP values not yet agreed upon by the COP should be reported in Table 9 (b).

Documentation box:

Parties should provide detailed explanations on the industrial processes sector in Chapter 4: Industrial processes (CRF sector 2) of the NIR. Use this documentation box to provide references to relevant sections of the NIR if any additional information and/or further details are needed to understand the content of this table.

If estimates are reported under "2.G Other", use this documentation box to provide information regarding activities covered under this category and to provide reference to the section in the NIR where background information can be found.

National Inventory Report 2007 - ICELAND

TABLE 2(II).C SECTORAL BACKGROUND DATA FOR INDUSTRIAL PROCESSES

Metal Production
(Sheet 1 of 1)

Inventory 2005
Submission 2007 v1.1
ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	ACTIVITY DATA		IMPLIED EMISSION FACTORS ⁽²⁾			EMISSIONS											
			CF ₄	C ₂ F ₆	SF ₆	CF ₄		C ₂ F ₆		SF ₆							
	Description ⁽¹⁾					(t)		(kg/t)			Emissions ⁽³⁾	Recovery ⁽⁴⁾	Emissions ⁽³⁾	Recovery ⁽⁴⁾	Emissions ⁽³⁾	Recovery ⁽⁴⁾	
C. PFCs and SF₆ from Metal Production																	
PFCs from Aluminium Production	Aluminium production	272,488,00	0,01	0,00		3,40		0,44								NA,NO	
SF ₆ used in Aluminium and Magnesium Foundries						3,40		0,44								NO	
Aluminium Foundries	(specify)				NO											NO	
Magnesium Foundries	(specify)				NO											NO	

- (1) Specify the activity data used as shown in the examples in parentheses.
- (2) The implied emission factors (IEFs) are estimated on the basis of gross emissions as follows: IEF = (emissions + amounts recovered, oxidized, destroyed or transformed) / activity data.
- (3) Final emissions (after subtracting the amounts of emission recovery, oxidation, destruction or transformation).
- (4) Amounts of emission recovery, oxidation, destruction or transformation.

Documentation box:

- Parties should provide detailed explanations on the industrial processes sector in Chapter 4: Industrial processes (CRF sector 2) of the NIR. Use this documentation box to provide references to relevant sections of the NIR if any additional information and/or further details are needed to understand the content of this table.
- Where only aggregate figures for activity data are provided, e.g. due to reasons of confidentiality (see footnote 1 to table 2(II)), a note indicating this should be provided in this documentation box.
- Where applying Tier 1b and country-specific methods, specify any other relevant activity data used in this documentation box, including a reference to the section of the NIR where more detailed information can be found.
- Use this documentation box for providing clarification on emission recovery, oxidation, destruction and/or transformation, and provide a reference to the section of the NIR where more detailed information can be found.

TABLE 2(II).E SECTORAL BACKGROUND DATA FOR INDUSTRIAL PROCESSES
Production of Halocarbons and SF₆
 (Sheet 1 of 1)

Inventory 2005
 Submission 2007 v1.1
 ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	ACTIVITY DATA		IMPLIED EMISSION FACTORS ⁽²⁾	EMISSIONS	
	Description ⁽¹⁾	(t)		(kg/t)	Emissions ⁽³⁾
E. Production of Halocarbons and SF₆					
1. By-product Emissions					
Production of HCFC-22					
HFC-23	Production of HCFC-22		NO	NO	
Other (specify activity and chemical)					
Other non-specified					
2. Fugitive Emissions (specify activity and chemical)					
HFCs				NA,NO	
HFC-23				NO	
HFC-32				NO	
HFC-41				NO	
HFC-43-10-mee				NO	
HFC-125				NO	
HFC-134				NO	
HFC-134a				NO	
HFC-152a				NO	
HFC-143				NO	
HFC-143a				NO	
HFC-227ea				NO	
HFC-236fa				NO	
HFC-245ca				NO	
Unspecified mix of HFCs				NO	
PFCs				NA,NO	
CF ₄				NO	
C ₂ F ₆				NO	
C ₃ F ₈				NO	
C ₄ F ₁₀				NO	
c-C ₄ F ₈				NO	
CSF ₁₂				NO	
C ₆ F ₁₄				NO	
Unspecified mix of PFCs				NO	
SF ₆				NO	
Other non-specified					
HFCs				NO	
HFC-23				NO	
HFC-32				NO	
HFC-41				NO	
HFC-43-10-mee				NO	
HFC-125				NO	
HFC-134				NO	
HFC-134a				NO	
HFC-152a				NO	
HFC-143				NO	
HFC-143a				NO	
HFC-227ea				NO	
HFC-236fa				NO	
HFC-245ca				NO	
Unspecified mix of HFCs				NO	
PFCs				NO	
CF ₄				NO	
C ₂ F ₆				NO	
C ₃ F ₈				NO	
C ₄ F ₁₀				NO	
c-C ₄ F ₈				NO	
CSF ₁₂				NO	
C ₆ F ₁₄				NO	
Unspecified mix of PFCs				NO	
SF ₆				NO	
3. Other (specify activity and chemical)					
HFCs				NA	
HFC-23				NA	
HFC-32				NA	
HFC-41				NA	
HFC-43-10-mee				NA	
HFC-125				NA	
HFC-134				NA	
HFC-134a				NA	
HFC-152a				NA	
HFC-143				NA	
HFC-143a				NA	
HFC-227ea				NA	
HFC-236fa				NA	
HFC-245ca				NA	
Unspecified mix of HFCs				NA	
PFCs				NA	
CF ₄				NA	
C ₂ F ₆				NA	
C ₃ F ₈				NA	
C ₄ F ₁₀				NA	
c-C ₄ F ₈				NA	
CSF ₁₂				NA	
C ₆ F ₁₄				NA	
Unspecified mix of PFCs				NA	
SF ₆				NA	

⁽¹⁾ Specify the activity data used as shown in the examples within parentheses.

⁽²⁾ The implied emission factors (IEFs) are estimated on the basis of gross emissions as follows: IEF = (emissions + amounts recovered, oxidized, destroyed or transformed) / activity data.

⁽³⁾ Final emissions are to be reported (after subtracting the amounts of emission recovery, oxidation, destruction or transformation).

⁽⁴⁾ Amounts of emission recovery, oxidation, destruction or transformation.

Documentation box:

Parties should provide detailed explanations on the industrial processes sector in Chapter 4: Industrial processes (CRF sector 2) of the NIR. Use this documentation box to provide references to relevant sections of the NIR if any additional information and/or further information is provided. Where only aggregate figures for activity data are provided, e.g. due to reasons of confidentiality (see footnote 1 to table 2(II)), a note indicating this should be provided in this documentation box.

Where applying Tier 2 and country-specific methods, specify any other relevant activity data used in this documentation box, including a reference to the section of the NIR where more detailed information can be found.

Use this documentation box for providing clarification on emission recovery, oxidation, destruction and/or transformation, and provide a reference to the section of the NIR where more detailed information can be found.

National Inventory Report 2007 - ICELAND

Inventory 2005
Submission 2007 v1.1
ICELAND

TABLE 2(II).F SECTORAL BACKGROUND DATA FOR INDUSTRIAL PROCESSES
Consumption of Halocarbons and SF₆
(Sheet 1 of 2)

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	ACTIVITY DATA <i>Amount of fluid</i>			IMPLIED EMISSION FACTORS			EMISSIONS		
	Filled into new manufactured products	In operating systems (average annual stocks)	Remaining in products at decommissioning	Product manufacturing factor	Product life factor	Disposal loss factor	From manufacturing	From stocks	From disposal
	(t)			(% per annum)			(t)		
1. Refrigeration⁽¹⁾									
Air Conditioning Equipment									
Domestic Refrigeration <i>(please specify chemical)⁽¹⁾</i>									
Commercial Refrigeration									
Transport Refrigeration									
Industrial Refrigeration									
Stationary Air-Conditioning									
Mobile Air-Conditioning									
2. Foam Blowing⁽¹⁾									
Hard Foam									
Soft Foam									

⁽¹⁾ Under each of the listed source categories, specify the chemical consumed (e.g. HFC-32) as indicated under category Domestic Refrigeration; use one row per chemical.

Note: This table provides for reporting of the activity data and emission factors used to calculate actual emissions from consumption of halocarbons and SF₆ using the "bottom-up approach" (based on the total stock of equipment and estimated emission rates from this equipment). Some Parties may prefer to estimate actual emissions following the alternative "top-down approach" (based on annual sales of equipment and/or gas). Those Parties should indicate the activity data used and provide any other information needed to understand the content of the table in the documentation box at the end of sheet 2 to this table, including a reference to the section of the NIR where further details can be found. Those Parties should provide the following data in the NIR:

1. the amount of fluid used to fill new products,
2. the amount of fluid used to service existing products,
3. the amount of fluid originally used to fill retiring products (the total nameplate capacity of retiring products),
4. the product lifetime, and
5. the growth rate of product sales, if this has been used to calculate the amount of fluid originally used to fill retiring products.

In the NIR, Parties may provide alternative formats for reporting equivalent information with a similar level of detail.

National Inventory Report 2007 - ICELAND

TABLE 2(II).F. SECTORAL BACKGROUND DATA FOR INDUSTRIAL PROCESSES
Consumption of Halocarbons and SF₆
 (Sheet 2 of 2)

Inventory 2005
 Submission 2007 v1.1
 ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	ACTIVITY DATA <i>Amount of fluid</i>			IMPLIED EMISSION FACTORS			EMISSIONS		
	Filled into new manufactured products	In operating systems (average annual stocks)	Remaining in products at decommissioning	Product manufacturing factor	Product life factor	Disposal loss factor	From manufacturing	From stocks	From disposal
	(t)			(% per annum)			(t)		
3. Fire Extinguishers <i>(please specify chemical)</i> ⁽¹⁾									
4. Aerosols ⁽¹⁾									
Metered Dose Inhalers									
Other									
5. Solvents ⁽¹⁾									
6. Other applications using ODS⁽²⁾ substitutes ⁽¹⁾									
7. Semiconductor Manufacture ⁽¹⁾									
8. Electrical Equipment ⁽¹⁾									
9. Other <i>(please specify)</i> ⁽¹⁾									

⁽¹⁾ Under each of the listed source categories, specify the chemical consumed (e.g. HFC-32) as indicated under category Fire Extinguishers; use one row per chemical.

⁽²⁾ ODS: ozone-depleting substances.

Documentation box:

- Parties should provide detailed explanations on the industrial processes sector in Chapter 4: Industrial processes (CRF sector 2) of the NIR. Use this documentation box to provide references to relevant sections of the NIR if any additional information and/or further details are needed to understand the content of this table.
- Where only aggregate figures for activity data are provided, e.g. due to reasons of confidentiality (see footnote 1 to table 2(II)), a note indicating this should be provided in this documentation box.
- With regard to data on the amounts of fluid that remained in retired products at decommissioning, use this documentation box to provide a reference to the section of the NIR where information on the amount of the chemical recovered (recovery efficiency) and other relevant information used in the emission estimation can be found.
- Parties that estimate their actual emissions following the alternative top-down approach might not be able to report emissions using this table. As indicated in the note to sheet 1 of this table, Parties should in these cases provide, in the NIR, alternative formats for reporting equivalent

National Inventory Report 2007 - ICELAND

Inventory 2005
Submission 2007 v1.1
ICELAND

TABLE 3 SECTORAL REPORT FOR SOLVENT AND OTHER PRODUCT USE
(Sheet 1 of 1)

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	CO ₂	N ₂ O	NM VOC
	(Gg)		
Total Solvent and Other Product Use	NE	0,01	4,14
A. Paint Application	NE		1,29
B. Degreasing and Dry Cleaning	NE	NA	0,16
C. Chemical Products, Manufacture and Processing			1E
D. Other	NE	0,01	2,68
1. Use of N ₂ O for Anaesthesia		0,01	
2. N ₂ O from Fire Extinguishers		NE	
3. N ₂ O from Aerosol Cans		NE	
4. Other Use of N ₂ O		0,00	
5. Other (<i>as specified in table 3.A-D</i>)	NE	NA	2,68
Other non-specified	NE	NA	2,68

Note: The quantity of carbon released in the form of NMVOCs should be accounted for in both the NMVOC and the CO₂ columns. The quantities of NMVOCs should be converted into CO₂ equivalent emissions before being added to the CO₂ amounts in the CO₂ column.

Documentation box:

- Parties should provide detailed explanations about the Solvent and Other Product Use sector in Chapter 5: Solvent and Other Product Use (CRF sector 3) of the NIR. Use this documentation box to provide references to relevant sections of the NIR if any additional information and/or further details are needed to understand the content of this table.
- The IPCC Guidelines do not provide methodologies for the calculation of emissions of N₂O from Solvent and Other Product Use. If reporting such data, Parties should provide in the NIR additional information (activity data and emission factors) used to derive these estimates, and provide in this documentation box a reference to the section of the NIR where this information can be found.

National Inventory Report 2007 - ICELAND

TABLE 3.A-D SECTORAL BACKGROUND DATA FOR SOLVENT AND OTHER PRODUCT USE
(Sheet 1 of 1)

Inventory 2005
Submission 2007 v1.1
ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	ACTIVITY DATA		IMPLIED EMISSION FACTORS ⁽¹⁾	
	Description	(kt)	CO ₂ (t/t)	N ₂ O (t/t)
A. Paint Application	Paint use	2,58	NE	
B. Degreasing and Dry Cleaning	Imports	0,16	NE	NA
C. Chemical Products, Manufacture and Processing	(specify)			
D. Other				
1. Use of N ₂ O for Anaesthesia	Use of N2O	0,01		1,00
2. N ₂ O from Fire Extinguishers	(specify)	NE		NE
3. N ₂ O from Aerosol Cans	(specify)	NE		NE
4. Other Use of N ₂ O	Use of N2O	0,00		1,00
5. Other <i>(please specify)</i> ⁽²⁾				
Other non-specified	(specify)	NA	NE	NA

⁽¹⁾ The implied emission factors will not be calculated until the corresponding emission estimates are entered directly into table 3.

⁽²⁾ Some probable sources to be reported under 3.D Other are listed in this table. Complement the list with other relevant sources, as appropriate.

Documentation box:

Parties should provide detailed explanations on the Solvent and Other Product Use sector in Chapter 5: Solvent and Other Product Use (CRF sector 3) of the NIR. Use this documentation box to provide references to relevant sections of the NIR if any additional information and/or further details are needed to understand the content of this table.

National Inventory Report 2007 - ICELAND

Inventory 2005
Submission 2007 v1.1
ICELAND

TABLE 4 SECTORAL REPORT FOR AGRICULTURE
(Sheet 1 of 2)

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	CH ₄	N ₂ O	NO _x (Gg)	CO	NMVOC
Total Agriculture	12,00	0,72	NA,NO	NA,NO	NA,NE,NO
A. Enteric Fermentation	11,02				
1. Cattle ⁽¹⁾	4,49				
<i>Option A:</i>					
Dairy Cattle	2,45				
Non-Dairy Cattle	2,04				
<i>Option B:</i>					
Mature Dairy Cattle					
Mature Non-Dairy Cattle					
Young Cattle					
2. Buffalo	NO				
3. Sheep	5,09				
4. Goats	0,00				
5. Camels and Llamas	NO				
6. Horses	1,39				
7. Mules and Asses	NO				
8. Swine	0,05				
9. Poultry	NE				
10. Other (as specified in table 4.A)					
Goats					
Horses					
B. Manure Management	0,97	0,08			NE,NO
1. Cattle ⁽¹⁾	0,60				
<i>Option A:</i>					
Dairy Cattle	0,34				
Non-Dairy Cattle	0,25				
<i>Option B:</i>					
Mature Dairy Cattle					
Mature Non-Dairy Cattle					
Young Cattle					
2. Buffalo	NO				
3. Sheep	0,12				
4. Goats	0,00				
5. Camels and Llamas	NO				
6. Horses	0,11				
7. Mules and Asses	NO				
8. Swine	0,10				
9. Poultry	0,05				
10. Other livestock (as specified in table 4.B(a))					
Goats					
Horses					

Note: All footnotes for this table are given at the end of the table on sheet 2.

National Inventory Report 2007 - ICELAND

TABLE 4 SECTORAL REPORT FOR AGRICULTURE
(Sheet 2 of 2)

Inventory 2005
Submission 2007 v1.1
ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	CH ₄	N ₂ O	NO _x	CO	NMVOG
	(Gg)				
B. Manure Management (continued)					
11. Anaerobic Lagoons		NO			NO
12. Liquid Systems		0,00			NE
13. Solid Storage and Dry Lot		0,08			NE
14. Other AWMS					
C. Rice Cultivation	NA,NO				NA,NO
1. Irrigated	NO				NO
2. Rainfed	NO				NO
3. Deep Water	NO				NO
4. Other (as specified in table 4.C)	NA				NA
D. Agricultural Soils⁽²⁾	NA,NE	0,64			NA,NE
1. Direct Soil Emissions	NE	0,24			NE
2. Pasture, Range and Paddock Manure ⁽³⁾		0,14			NE
3. Indirect Emissions	NE	0,26			NE
4. Other (as specified in table 4.D)	NA	NA			NA
E. Prescribed Burning of Savannas	NA	NA	NO	NO	NO
F. Field Burning of Agricultural Residues	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO
1. Cereals	NA,NO	NA,NO	NO	NO	NO
2. Pulses	NA,NO	NA,NO	NO	NO	NO
3. Tubers and Roots	NA,NO	NA,NO	NO	NO	NO
4. Sugar Cane	NO	NO	NO	NO	NO
5. Other (as specified in table 4.F)	NA	NA	NA	NA	NA
G. Other (please specify)	NA	NA	NA	NA	NA

⁽¹⁾ The sum for cattle would be calculated on the basis of entries made under either option A (dairy and non-dairy cattle) or option B (mature dairy cattle, mature non-dairy cattle and young cattle).

⁽²⁾ See footnote 4 to Summary 1.A of this common reporting format. Parties which choose to report CO₂ emissions and removals from agricultural soils under 4.D Agricultural Soils of the sector Agriculture should report the amount (in Gg) of these emissions or removals in table Summary 1.A of the CRF. References to additional information (activity data, emissions factors) reported in the NIR should be provided in the documentation box to table 4.D. In line with the corresponding table in the IPCC Guidelines (i.e. IPCC Sectoral Report for Agriculture), this table does not include provisions for reporting CO₂ estimates.

⁽³⁾ Direct N₂O emissions from pasture, range and paddock manure are to be reported in the "4.D Agricultural Soils" category. All other N₂O emissions from animal manure are to be reported in the "4.B Manure Management" category. See also chapter 4.4 of the IPCC good practice guidance report.

Note: The IPCC Guidelines do not provide methodologies for the calculation of CH₄ emissions and CH₄ and N₂O removals from agricultural soils, or CO₂ emissions from prescribed burning of savannas and field burning of agricultural residues. Parties that have estimated such emissions should provide, in the NIR, additional information (activity data and emission factors) used to derive these estimates and include a reference to the section of the NIR in the documentation box of the corresponding Sectoral background data tables.

National Inventory Report 2007 - ICELAND

Inventory 2005
Submission 2007 v1.1
ICELAND

TABLE 4.A SECTORAL BACKGROUND DATA FOR AGRICULTURE
Enteric Fermentation
(Sheet 1 of 1)

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	ACTIVITY DATA AND OTHER RELATED INFORMATION			IMPLIED EMISSION FACTORS ⁽¹⁾
	Population size ⁽¹⁾ (1000s)	Average gross energy intake (GE) (MJ/head/day)	Average CH ₄ conversion rate (Y _m) ⁽²⁾ (%)	CH ₄ (kg CH ₄ /head/yr)
1. Cattle	66,96			67,06
Option A:				
Dairy Cattle ⁽⁴⁾	24,54			100,00
Non-Dairy Cattle	42,42			48,00
Option B:				
Mature Dairy Cattle				
Mature Non-Dairy Cattle				
Young Cattle				
2. Buffalo	NO			NO
3. Sheep	636,09			8,00
4. Goats	0,66			5,00
5. Camels and Llamas	NO			NO
6. Horses	77,30			18,00
7. Mules and Asses	NO			NO
8. Swine	33,27			1,50
9. Poultry	596,23			NE
10. Other (please specify)				
Goats	0,66			
Horses	77,30			

⁽¹⁾ Parties are encouraged to provide detailed livestock population data by animal type and region, if available, in the NIR, and provide in the documentation box below a reference to the relevant section. Parties should use the same animal population statistics to estimate CH₄ emissions from enteric fermentation, CH₄ and N₂O from manure management, N₂O direct emissions from soil and N₂O emissions associated with manure production, as well as emissions from the use of manure as fuel, and sewage-related emissions reported in the Waste sector.

Additional information (only for those livestock types for which Tier 2 was used)⁽⁶⁾

Disaggregated list of animals ⁽⁴⁾	Dairy Cattle	Non-Dairy Cattle	Mature Dairy Cattle	Mature Non-Dairy Cattle	Young Cattle	Buffalo	Sheep	Goats	Camels and Llamas	Horses	Mules and Asses	Swine	Poultry	Other (specify)	Goats	Horses
	Indicators:															
Weight	(kg)															
Feeding situation ⁽⁵⁾																
Milk yield	(kg/day)															
Work	(h/day)															
Resistant	(%)	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Digestibility of feed	(%)	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00

⁽⁶⁾ See also Tables A-1 and A-2 of the IPCC Guidelines (Volume 3, Reference Manual, pp. 4.31-4.34). These data are relevant if Parties do not have data on average feed intake.

⁽⁴⁾ Disaggregate to the split actually used. Add columns to the table if necessary.

⁽⁵⁾ Specify feeding situation as pasture, stall fed, confined, open range, etc.

National Inventory Report 2007 - ICELAND

TABLE 4.B(a) SECTORAL BACKGROUND DATA FOR AGRICULTURE
CH₄ Emissions from Manure Management
(Sheet 1 of 2)

Inventory 2005
 Submission 2007 v1.1
 ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	ACTIVITY DATA AND OTHER RELATED INFORMATION						IMPLIED EMISSION FACTORS ⁽⁴⁾ CH ₄ (kg CH ₄ /head/yr)	
	Population size (1000s)	Allocation by climate region ⁽¹⁾			Typical animal mass (average) (kg)	VS ⁽²⁾ daily excretion (average) (kg dm/head/day)		CH ₄ producing potential (Bo) ⁽²⁾ (average) (m ³ CH ₄ /kg VS)
		Cool	Temperate	Warm				
			(%)					
1. Cattle	66,96						8,93	
<i>Option A:</i>								
Dairy Cattle ⁽³⁾	24,54	1,00	0,00	0,00			14,00	
Non-Dairy Cattle	42,42	1,00	0,00	0,00			6,00	
<i>Option B:</i>								
Mature Dairy Cattle		0,00	0,00	0,00				
Mature Non-Dairy Cattle		0,00	0,00	0,00				
Young Cattle		0,00	0,00	0,00				
2. Buffalo	NO	0,00	0,00	0,00			NO	
3. Sheep	636,09	1,00	0,00	0,00			0,19	
4. Goats	0,66	1,00	0,00	0,00			0,12	
5. Camels and Llamas	NO	0,00	0,00	0,00			NO	
6. Horses	77,30	1,00	0,00	0,00			1,40	
7. Mules and Asses	NO	0,00	0,00	0,00			NO	
8. Swine	33,27	1,00	0,00	0,00			3,00	
9. Poultry	596,23	1,00	0,00	0,00			0,08	
10. Other livestock (<i>please specify</i>)								
Goats	0,66	1,00	0,00	0,00				
Horses	77,30	1,00	0,00	0,00				

⁽¹⁾ Climate regions are defined in terms of annual average temperature as follows: Cool = less than 15°C; Temperate = 15 - 25°C inclusive; and Warm = greater than 25°C (see table 4.2 of the IPCC Guidelines (Volume 3, Reference Manual, p. 4.8)).

⁽²⁾ VS = Volatile Solids; Bo = maximum methane producing capacity for manure IPCC Guidelines (Volume 3, Reference Manual, p.4.23 and p.4.15); dm = dry matter. Provide average values for VS and Bo where original calculations were made at a more disaggregated level of these livestock categories.

⁽³⁾ Including data on dairy heifers, if available.

⁽⁴⁾ The implied emission factors will not be calculated until the corresponding emission estimates are entered directly into table 4.

Documentation box:

- Parties should provide detailed explanations on the Agriculture sector in Chapter 6: Agriculture (CRF sector 4) of the NIR. Use this documentation box to provide references to relevant sections of the NIR if any additional information and further details are needed to understand the content of this table.
- Indicate in this documentation box whether the activity data used are one-year estimates or three-year averages.
- Provide a reference to the relevant section in the NIR, in particular with regard to:
 - (a) disaggregation of livestock population (e.g. according to the classification recommended in the IPCC good practice guidance), including information on whether these data are one-year estimates or three-year averages.
 - (b) parameters relevant to the application of IPCC good practice guidance;
 - (c) information on how the MCFs are derived, if relevant data could not be provided in the additional information box.

TABLE 4.B(a) SECTORAL BACKGROUND DATA FOR AGRICULTURE
 CH₄ Emissions from Manure Management
 (Sheet 2 of 2)

Inventory 2005
 Submission 2007 v1.1
 ICELAND

Additional information (for Tier 2)^(a)

Animal category	Indicator	Climate region	Animal waste management system						
			Anaerobic lagoon	Liquid system	Daily spread	Solid storage	Dry lot	Pasture range paddock	Other
Dairy Cattle	Allocation (%)	Cool		0,53	NO	0,13		0,34	NO
		Temperate							
		Warm							
	MCF ^(b)	Cool							
		Temperate							
		Warm							
Non-Dairy Cattle	Allocation (%)	Cool		0,53	NO	0,13		0,34	NO
		Temperate							
		Warm							
	MCF ^(b)	Cool							
		Temperate							
		Warm							
Mature Dairy Cattle	Allocation (%)	Cool							
		Temperate							
		Warm							
	MCF ^(b)	Cool							
		Temperate							
		Warm							
Mature Non-Dairy Cattle	Allocation (%)	Cool							
		Temperate							
		Warm							
	MCF ^(b)	Cool							
		Temperate							
		Warm							
Young Cattle	Allocation (%)	Cool							
		Temperate							
		Warm							
	MCF ^(b)	Cool							
		Temperate							
		Warm							
Buffalo	Allocation (%)	Cool							
		Temperate							
		Warm							
	MCF ^(b)	Cool							
		Temperate							
		Warm							
Sheep	Allocation (%)	Cool							
		Temperate							
		Warm							
	MCF ^(b)	Cool							
		Temperate							
		Warm							
Goats	Allocation (%)	Cool							
		Temperate							
		Warm							
	MCF ^(b)	Cool							
		Temperate							
		Warm							
Camels and Llamas	Allocation (%)	Cool							
		Temperate							
		Warm							
	MCF ^(b)	Cool							
		Temperate							
		Warm							
Horses	Allocation (%)	Cool							
		Temperate							
		Warm							
	MCF ^(b)	Cool							
		Temperate							
		Warm							
Mules and Asses	Allocation (%)	Cool							
		Temperate							
		Warm							
	MCF ^(b)	Cool							
		Temperate							
		Warm							
Swine	Allocation (%)	Cool		1,00	NO	0,00		0,00	NO
		Temperate							
		Warm							
	MCF ^(b)	Cool							
		Temperate							
		Warm							
Poultry	Allocation (%)	Cool							
		Temperate							
		Warm							
	MCF ^(b)	Cool							
		Temperate							
		Warm							
Other livestock (please specify)	Allocation (%)	Cool							
		Temperate							
		Warm							
	MCF ^(b)	Cool							
		Temperate							
		Warm							

^(a) The information required in this table may not be directly applicable to country-specific methods developed for MCF calculations. In such cases, information on MCF derivation should be described in the NIR and references to the relevant sections of the NIR should be provided in the documentation box.

^(b) MCF = Methane Conversion Factor (IPCC Guidelines, (Volume 3, Reference Manual, p. 4.9)). If another climate region categorization is used, replace the entries in the cells with the climate regions for which the MCFs are specified.

National Inventory Report 2007 - ICELAND

TABLE 4.B(b) SECTORAL BACKGROUND DATA FOR AGRICULTURE
N₂O Emissions from Manure Management
(Sheet 1 of 1)

Inventory 2005
 Submission 2007 v1.1
 ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	ACTIVITY DATA AND OTHER RELATED INFORMATION									IMPLIED EMISSION FACTORS ⁽¹⁾	
	Population size (1000s)	Nitrogen excretion (kg N/head/yr)	Nitrogen excretion per animal waste management system (AWMS) (kg N/yr)						Emission factor per animal waste management system (kg N ₂ O-N/kg N)		
			Anaerobic lagoon	Liquid system	Daily spread	Solid storage and dry lot	Pasture range and paddock	Other			
Cattle	66.96		NO	1 535 770,40	NO	376 698,40	985 211,20	NO	Anaerobic lagoon	NO	
<i>Option A:</i>									Liquid system	0,00	
Dairy Cattle	24,54	60,00	NO	780 308,40	NO	191 396,40	500 575,20	NO	Solid storage and dry lot	0,02	
Non-Dairy Cattle	42,42	33,60	NO	755 462,00	NO	185 302,00	484 636,00	NO	Other AWMS	NO	
<i>Option B:</i>											
Mature Dairy Cattle											
Mature Non-Dairy Cattle											
Young Cattle											
Sheep	636,09	5,76	NO	622 863,46	NO	1 502 200,11	1 538 839,13	NO			
Swine	33,27	13,30	NO	442 480,72	NO			NO			
Poultry	596,23	0,42	NO		NO	250 417,37		NO			
Buffalo	NO										
Goats	0,66										
Camels and Llamas	NO										
Horses	77,30										
Mules and Asses	NO										
Other livestock (<i>please specify</i>)											
Goats	0,66	5,76	NO	644,80	NO	1 555,11	1 593,04	NO			
Horses	77,30	28,80	NO		NO	378 473,55	1 847 841,47	NO			
Total per AWMS			NO	2 601 759,37	NO	2 509 344,54	4 373 484,85	NO			

⁽¹⁾ The implied emission factor will not be calculated until the emissions are entered directly into table 4.

Documentation box:

- Parties should provide detailed explanations on the Agriculture sector in Chapter 6: Agriculture (CRF sector 4) of the NIR. Use this documentation box to provide references to relevant sections of the NIR if any additional information and/or further details are needed to understand the content of this table.
- Indicate in this documentation box whether the activity data used are one-year estimates or three-year averages.
- Provide a reference to the relevant section in the NIR, in particular with regard to:
 - (a) disaggregation of livestock population (e.g. according to the classification recommended in the IPCC good practice guidance), including information on whether these data are one-year estimates or three-year averages.
 - (b) information on other AWMS, if reported.

National Inventory Report 2007 - ICELAND

Inventory 2005
Submission 2007 v1.1
ICELAND

TABLE 4.C SECTORAL BACKGROUND DATA FOR AGRICULTURE
Rice Cultivation
(Sheet 1 of 1)

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	ACTIVITY DATA AND OTHER RELATED INFORMATION			IMPLIED EMISSION FACTOR ⁽¹⁾ CH ₄ (g/m ²)	EMISSIONS CH ₄ (Gg)
	Harvested area ⁽²⁾ (10 ⁹ m ² /yr)	Organic amendments added ⁽³⁾			
		type	(t/ha)		
1. Irrigated					NO
Continuously Flooded	NO	(specify type)		NO	NO
Intermittently Flooded	NO	(specify type)		NO	NO
	NO	(specify type)		NO	NO
2. Rainfed					NO
Flood Prone	NO	(specify type)		NO	NO
Drought Prone	NO	(specify type)		NO	NO
3. Deep Water					NO
Water Depth 50-100 cm	NO	(specify type)		NO	NO
Water Depth > 100 cm	NO	(specify type)		NO	NO
4. Other (please specify)	NA				NA
Upland Rice ⁽⁴⁾					
Total ⁽⁴⁾	NA,NO				

⁽¹⁾ The implied emission factor implicitly takes account of all relevant corrections for continuously flooded fields without organic amendment, the correction for the organic amendments and the effect of different soil characteristics, if considered in the calculation of methane emissions.

⁽²⁾ Harvested area is the cultivated area multiplied by the number of cropping seasons per year.

⁽³⁾ Specify dry weight or wet weight for organic amendments in the documentation box.

⁽⁴⁾ These rows are included to allow comparison with international statistics. Methane emissions from upland rice are assumed to be zero.

Documentation box:

- Parties should provide detailed explanations on the Agriculture sector in Chapter 6: Agriculture (CRF sector 4) of the NIR. Use this documentation box to provide references to relevant sections of the NIR if any additional information and/or further details are needed to understand the content of this table.
- When disaggregating by more than one region within a country, and/or by growing season, provide additional information on disaggregation and related data in the NIR and provide a reference to the relevant section in the NIR.
- Where available, provide activity data and scaling factors by soil type and rice cultivar in the NIR.

National Inventory Report 2007 - ICELAND

TABLE 4.D SECTORAL BACKGROUND DATA FOR AGRICULTURE

Agricultural Soils
(Sheet 1 of 2)

Inventory 2005
Submission 2007 v1.1
ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	ACTIVITY DATA AND OTHER RELATED INFORMATION		IMPLIED EMISSION FACTORS kg N ₂ O-N/kg N ⁽²⁾	EMISSIONS N ₂ O (Gg)
	Description	Value kg N/yr		
1. Direct Soil Emissions	N input to soils			0,24
1. Synthetic Fertilizers	Nitrogen input from application of synthetic fertilizers	8.051.155,83	0,01	0,16
2. Animal Manure Applied to Soils	Nitrogen input from manure applied to soils	4.088.883,13	0,01	0,08
3. N-fixing Crops	Nitrogen fixed by N-fixing crops	NO	NO	NO
4. Crop Residue	Nitrogen in crop residues returned to soils	14.952,88	0,01	0,00
5. Cultivation of Histosols ⁽²⁾	Area of cultivated organic soils (ha/yr)	NE	NE	NE
6. Other direct emissions (<i>please specify</i>)				NA
2. Pasture, Range and Paddock Manure	N excretion on pasture range and paddock	4.373.484,85	0,02	0,14
3. Indirect Emissions				0,26
1. Atmospheric Deposition	Volatized N from fertilizers, animal manures and other	2.791.490,62	0,01	0,04
2. Nitrogen Leaching and Run-off	N from fertilizers, animal manures and other that is lost through leaching and run-off	5.529.095,24	0,02	0,22
4. Other (<i>please specify</i>)				NA

⁽¹⁾ To convert from N₂O-N to N₂O emissions, multiply by 44/28. Note that for cultivation of Histosols the unit of the IEF is kg N₂O-N/ha.

Documentation box:

- Parties should provide detailed explanations on the Agriculture sector in Chapter 6: Agriculture (CRF sector 4) of the NIR. Use this documentation box to provide references to relevant sections of the NIR if any additional information and/or further details are needed to understand the content of this table.
- Provide a reference to the relevant section in the NIR, in particular with regard to:
 - Background information on CH₄ emissions from agricultural soils, if accounted for under the Agriculture sector;
 - Disaggregated values for Frac_{GRAZ} according to animal type, and for Frac_{BURN} according to crop types;
 - Full list of assumptions and fractions used.

TABLE 4.D SECTORAL BACKGROUND DATA FOR AGRICULTURE

Inventory 2005

Agricultural Soils⁽¹⁾

Submission 2007 v1.1

(Sheet 2 of 2)

ICELAND

Additional information

Fraction ^(a)	Description	Value
Frac _{BURN}	Fraction of crop residue burned	NO
Frac _{FUEL}	Fraction of livestock N excretion in excrements burned for fuel	NO
Frac _{GASF}	Fraction of synthetic fertilizer N applied to soils that volatilizes as NH ₃ and NO _x	0,00
Frac _{GASM}	Fraction of livestock N excretion that volatilizes as NH ₃ and NO _x	0,00
Frac _{GRAZ}	Fraction of livestock N excreted and deposited onto soil during grazing	0,00
Frac _{LEACH}	Fraction of N input to soils that is lost through leaching and run-off	0,00
Frac _{NCRBF}	Fraction of total above-ground biomass of N-fixing crop that is N	0,00
Frac _{NCRO}	Fraction of residue dry biomass that is N	NO
Frac _R	Fraction of total above-ground crop biomass that is removed from the field as a crop product	0,00
Other fractions <i>(please specify)</i>		0,00

^(a) Use the definitions for fractions as specified in the IPCC Guidelines (Volume 3. Reference Manual, pp. 4.92-4.113) as elaborated by the IPCC good practice guidance (pp. 4.54-4.74).

National Inventory Report 2007 - ICELAND

Inventory 2005
Submission 2007 v1.1
ICELAND

TABLE 4.E SECTORAL BACKGROUND DATA FOR AGRICULTURE
Prescribed Burning of Savannas
(Sheet 1 of 1)

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	ACTIVITY DATA AND OTHER RELATED INFORMATION					IMPLIED EMISSION FACTORS		EMISSIONS	
	Area of savanna burned (k ha/yr)	Average above-ground biomass density (t dm/ha)	Fraction of savanna burned	Biomass burned (Gg dm)	Nitrogen fraction in biomass	CH ₄	N ₂ O	CH ₄	N ₂ O
						(kg/t dm)		(Gg)	
(specify ecological zone)								NA	NA

Additional information

	Living Biomass	Dead Biomass
Fraction of above-ground biomass	0,00	0,00
Fraction oxidized	0,00	0,00
Carbon fraction	0,00	0,00

Documentation box:

Parties should provide detailed explanations on the Agriculture sector in Chapter 6: Agriculture (CRF sector 4) of the NIR. Use this documentation box to provide references to relevant sections of the NIR if any additional information and/or further details are needed to understand the content of this table.

National Inventory Report 2007 - ICELAND

TABLE 4.F. SECTORAL BACKGROUND DATA FOR AGRICULTURE

Field Burning of Agricultural Residues

(Sheet 1 of 1)

Inventory 2005

Submission 2007 v1.1

ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	ACTIVITY DATA AND OTHER RELATED INFORMATION							IMPLIED EMISSION FACTORS		EMISSIONS		
	Crop production (t)	Residue/ Crop ratio	Dry matter (dm) fraction of residue	Fraction burned in fields	Fraction oxidized	Total biomass burned (Gg dm)	C fraction of residue	N-C ratio in biomass residues	CH ₄	N ₂ O	CH ₄	N ₂ O
									(kg/t dm)		(Gg)	
1. Cereals											NA,NO	NA,NO
Wheat		0,00	0,00	0,00	0,00		0,00	NO	NO	NO	NO	NO
Barley		0,00	0,00	0,00	0,00		0,00	NO	NO	NO	NO	NO
Maize		0,00	0,00	0,00	0,00		0,00	NO	NO	NO	NO	NO
Oats		0,00	0,00	0,00	0,00		0,00	NO	NO	NO	NO	NO
Rye		0,00	0,00	0,00	0,00		0,00	NO	NO	NO	NO	NO
Rice		0,00	0,00	0,00	0,00		0,00	NO	NO	NO	NO	NO
Other (please specify)											NA	NA
2. Pulses											NA,NO	NA,NO
Dry bean		0,00	0,00	0,00	0,00		0,00	NO	NO	NO	NO	NO
Peas		0,00	0,00	0,00	0,00		0,00	NO	NO	NO	NO	NO
Soybeans		0,00	0,00	0,00	0,00		0,00	NO	NO	NO	NO	NO
Other (please specify)											NA	NA
3 Tubers and Roots											NA,NO	NA,NO
Potatoes		0,00	0,00	0,00	0,00		0,00	NO	NO	NO	NO	NO
Other (please specify)											NA	NA
4 Sugar Cane		0,00	0,00	0,00	0,00		0,00	NO	NO	NO	NO	NO
5 Other (please specify)											NA	NA

Documentation box:

Parties should provide detailed explanations on the Agriculture sector in Chapter 6: Agriculture (CRF sector 4) of the NIR. Use this documentation box to provide references to relevant sections of the NIR if any additional information and/or further details are needed to understand the content of this table.

National Inventory Report 2007 - ICELAND

**TABLE 5 SECTORAL REPORT FOR LAND USE, LAND-USE CHANGE AND FORESTRY
(Sheet 1 of 1)**

Inventory 2005
Submission 2007 v1.1
ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	Net CO ₂	CH ₄ ⁽²⁾	N ₂ O ⁽²⁾	NO _x	CO	NM VOC
	emissions/removals ^{(1),(2)}					
(Gg)						
Total Land-Use Categories	1,286,10	2,34	1,35	NA,NE	NA,NE	NA,NE
A. Forest Land	-126,27	NA,NE,NO	0,00	NE	NE	NE
1. Forest Land remaining Forest Land	-118,48	NA,NE,NO	0,00	NE	NE	NE
2. Land converted to Forest Land	-7,78	NE,NO	0,00	NE	NE	NE
B. Cropland	3,46	NE,NO	NE,NO	NE	NE	NE
1. Cropland remaining Cropland	3,46	NE	NE	NE	NE	NE
2. Land converted to Cropland	NE,NO	NE,NO	NE,NO	NE	NE	NE
C. Grassland	1,800,74	NE	NE	NE	NE	NE
1. Grassland remaining Grassland	1,800,74	NE	NE	NE	NE	NE
2. Land converted to Grassland	NE,NO	NE	NE	NE	NE	NE
D. Wetlands	141,42	2,34	0,07	NA,NE	NA,NE	NA,NE
1. Wetlands remaining Wetlands ⁽³⁾	141,42	2,34	0,07	NA	NA	NA
2. Land converted to Wetlands	NE,NO	NE,NO	NE,NO	NE	NE	NE
E. Settlements	NE	NE	NE	NE	NE	NE
1. Settlements remaining Settlements ⁽³⁾	NE	NE	NE	NE	NE	NE
2. Land converted to Settlements	NE	NE	NE	NE	NE	NE
F. Other Land	NE	NE	NE	NE	NE	NE
1. Other Land remaining Other Land ⁽⁴⁾	NE	NE	NE	NE	NE	NE
2. Land converted to Other Land	NE	NE	NE	NE	NE	NE
G. Other (please specify)⁽⁵⁾	-533,24	NA,NE,NO	1,28	NE	NE	NE
<i>Harvested Wood Products⁽⁶⁾</i>						
Revegetation	-533,24	NE,NO	0,02	NE	NE	NE
Grassland organic soil	NE,NO	NA,NE,NO	1,26	NE	NE	NE
Information items⁽⁷⁾						
Forest Land converted to other Land-Use Categories	NE	NE	NE	NE	NE	NE
Grassland converted to other Land-Use Categories	NE	NE	NE	NE	NE	NE

⁽¹⁾ According to the Revised 1996 IPCC Guidelines, for the purposes of reporting, the signs for removals are always negative (-) and for emissions positive (+).

⁽²⁾ For each land-use category and sub-category, this table sums net CO₂ emissions and removals shown in tables 5.A to 5.F, and the CO₂, CH₄ and N₂O emissions showing in tables 5(I) to 5(V).

⁽³⁾ Parties may decide not to prepare estimates for these categories contained in appendices 3a.3 and 3a.4 of the IPCC good practice guidance for LULUCF, although they may do so if they wish.

⁽⁴⁾ This land-use category is to allow the total of identified land area to match the national area.

⁽⁵⁾ The total for category 5.G Other includes items specified only under category 5.G in this table as well as sources and sinks specified in category 5.G in tables 5(I) to 5(V).

⁽⁶⁾ Parties may decide not to prepare estimates for this category contained in appendix 3a.1 of the IPCC good practice guidance for LULUCF, although they may do so if they wish and report in this row.

⁽⁷⁾ These items are listed for information only and will not be added to the totals, because they are already included in subcategories 5.A.2 to 5.F.2.

<p>Documentation box:</p> <ul style="list-style-type: none"> Parties should provide detailed explanations on the Land Use, Land-Use Change and Forestry sector in Chapter 7: Land Use, Land-Use Change and Forestry (CRF sector 5) of the NIR. Use this documentation box to provide references to relevant sections of the NIR if any additional information and/or further details are needed to understand the content of this table. If estimates are reported under 5.G Other, use this documentation box to provide information regarding activities covered under this category and to provide reference to the section in the NIR where background information can be found.
5.A Forest Land: Land conversion due to afforestation only reported as aggregated estimate. All afforestation assumed on grassland.
5.C.1 Grassland remaining Grassland: Extensive drainage of wetland has taken place in Iceland mostly in the period from 1940- 85. Information on this drainage is not geographically identifiable. All this drainage is therefore classified as grassland although some of it should be classified to landuse the category Cropland.
5.D Wetlands: CH ₄ and N ₂ O emissions from reservoirs
5.G Revegetation: The CO ₂ sink of revegetation is reported here. Background information on revegetation are provided in NIR (Ch7).
5.G 5(H) Non-CO ₂ emissions from drainage of soils and wetlands: Extensive drainage of wetland has taken place in Iceland mostly in the period from 1940- 85. Information on this drainage is not geographically identifiable. All this drainage is therefore classified as grassland although some of it should be classified to the landuse category Cropland. Accordingly N ₂ O emission from drained cultivated organic soils (Histosols, Andic Histosol and Gleyic Andosol) is included here.

National Inventory Report 2007 - ICELAND

TABLE 5.A SECTORAL BACKGROUND DATA FOR LAND USE, LAND-USE CHANGE AND FORESTRY
Forest Land
 (Sheet 1 of 1)

Inventory 2005
 Submission 2007 v1.1
 ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES		ACTIVITY DATA		IMPLIED CARBON-STOCK-CHANGE FACTORS						CHANGES IN CARBON STOCK						Net CO ₂ emissions/removals ^{(8) (9)}	
Land-Use Category	Sub-division ⁽¹⁾	Area ⁽²⁾ (kha)	Area of organic soil ⁽³⁾ (kha)	Carbon stock change in living biomass per area ⁽⁴⁾			Net carbon stock change in dead organic matter per area ⁽⁴⁾	Net carbon stock change in soils per area ⁽⁴⁾		Carbon stock change in living biomass ^{(3) (4)}			Net carbon stock change in dead organic matter ⁽⁴⁾	Net carbon stock change in soils ^{(4) (6)}			(Gg)
				Gains	Losses	Net change		Mineral soils ⁽⁵⁾	Organic soils	Gains	Losses	Net change		Mineral soils	Organic soils ⁽⁷⁾		
				(Mg C/ha)						(Gg C)							
A. Total Forest Land		54,20	3,76	0,65	NE,NO	0,65	NE,NO	NE,NO	NE,NO	-0,16	35,04	NE,NO	35,04	NE,NO	NE,NO	-0,60	-126,27
1. Forest Land remaining Forest Land		52,40	3,53	0,63	NE	0,63	NE	NE	NE	-0,16	32,88	NE	32,88	NE	NE	-0,57	-118,48
	Native Birch	25,00	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
	Plantations	27,40	3,53	1,20	NE	1,20	NE	NE	NE	-0,16	32,88	NE	32,88	NE	NE	-0,57	-118,48
2. Land converted to Forest Land ⁽¹⁰⁾		1,80	0,23	1,20	NE,NO	1,20	NE,NO	NE,NO	NE,NO	-0,16	2,16	NE,NO	2,16	NE,NO	NE,NO	-0,04	-7,78
2.1 Cropland converted to Forest Land		NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
2.2 Grassland converted to Forest Land		1,80	0,23	1,20	NE	1,20	NE	NE	NE	-0,16	2,16	NE	2,16	NE	NE	-0,04	-7,78
	New Plantations (ID=	1,80	0,23	1,20	NE	1,20	NE	NE	NE	-0,16	2,16	NE	2,16	NE	NE	-0,04	-7,78
2.3 Wetlands converted to Forest Land		NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
2.4 Settlements converted to Forest Land		NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
2.5 Other Land converted to Forest Land		NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE

- ⁽¹⁾ Land categories may be further divided according to climate zone, management system, soil type, vegetation type, tree species, ecological zone or national land classification.
- ⁽²⁾ The total area of the subcategories, in accordance with the sub-division used, should be entered here. For lands converted to Forest Land report the cumulative area remaining in the category in the reporting year.
- ⁽³⁾ Carbon stock gains and losses should be listed separately except in cases where, due to the methods used, it is technically impossible to separate information on gains and losses.
- ⁽⁴⁾ The signs for estimates of gains in carbon stocks are positive (+) and of losses in carbon stocks are negative (-).
- ⁽⁵⁾ Implied carbon-stock-change factors for mineral soils are calculated by dividing the net C stock change estimate for mineral soil by the difference between the area and the area of organic soil.
- ⁽⁶⁾ When Parties are estimating fluxes for organic soils but cannot separate these fluxes from mineral soils, these fluxes should be reported under mineral soils.
- ⁽⁷⁾ The value reported for organic soils is estimated as a flux. For consistency with other entries in this column, these fluxes should be expressed in the unit required in this column, i.e. in Gg C.
- ⁽⁸⁾ According to the Revised 1996 IPCC Guidelines, for the purposes of reporting, the signs for removals are always negative (-) and for emissions positive (+). Net changes in carbon stocks are converted to CO₂ by multiplying C by 44/12 and changing the sign for net CO₂ removals to be negative (-) and for net CO₂ emissions to be positive (+). Note that carbon stock changes in a single pool are not necessarily equal to emissions or removals, because some carbon stock changes result from carbon transfers among pools rather than exchanges with the atmosphere.
- ⁽⁹⁾ Where Parties directly estimate emissions and removals rather than carbon stock changes, they may report emissions/removals directly in this column and use notation keys in the stock change columns.
- ⁽¹⁰⁾ A Party may report aggregate estimates for all conversions of land to forest land when data are not available to report them separately. A Party should specify in the documentation box which types of land conversion are included. Separate estimates for grassland conversion should be provided in table 5 as an information item.

Documentation box:
 Parties should provide detailed explanations on the Land Use, Land-Use Change and Forestry sector in Chapter 7: Land Use, Land-Use Change and Forestry (CRF sector 5) of the NIR. Use this documentation box to provide references to relevant sections of the NIR if any additional information and/or further details are needed to understand the content of this table.

National Inventory Report 2007 - ICELAND

TABLE 5.B SECTORAL BACKGROUND DATA FOR LAND USE, LAND-USE CHANGE AND FORESTRY

Cropland
(Sheet 1 of 1)

Inventory 2005
Submission 2007 v1.1
ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES		ACTIVITY DATA		IMPLIED CARBON-STOCK-CHANGE FACTORS						CHANGES IN CARBON STOCK						Net CO ₂ emissions/removals ^{(10) (11)}	
Land-Use Category	Sub-division ⁽¹⁾	Area ⁽²⁾ (kha)	Area of organic soil ⁽²⁾ (kha)	Carbon stock change in living biomass per area ⁽³⁾ (4)			Net carbon stock change in dead organic matter per area ⁽⁴⁾	Net carbon stock change in soils per area ⁽⁴⁾		Carbon stock change in living biomass ^{(3), (4), (6)}			Net carbon stock change in dead organic matter ^{(4) (7)}	Net carbon stock change in soils ^{(4) (8)}			(Gg)
				Gains	Losses	Net change		Mineral soils ⁽⁵⁾	Organic soils	Gains	Losses	Net change		Mineral soils	Organic soils ⁽⁹⁾		
				(Mg C/ha)						(Gg C)							
B. Total Cropland		129.00	NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NE,NO	IE,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NE,NO	IE,NE,NO	IE,NA,NE,NO	
1. Cropland remaining Cropland		129.00	NE	NA	NA	NA	NA	NE	IE	NA	NA	NA	NA	NE	IE	IE,NA,NE	
2. Land converted to Cropland ⁽¹²⁾		NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	
2.1 Forest Land converted to Cropland		NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	
2.2 Grassland converted to Cropland		NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	
2.3 Wetlands converted to Cropland		NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	
2.4 Settlements converted to Cropland		NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	
2.5 Other Land converted to Cropland		NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	

- ⁽¹⁾ Land categories may be further divided according to climate zone, management system, soil type, vegetation type, tree species, ecological zone or national land classification.
- ⁽²⁾ The total area of the subcategories, in accordance with the sub-division used, should be entered here. For lands converted to Cropland report the cumulative area remaining in the category in the reporting year.
- ⁽³⁾ Carbon stock gains and losses should be listed separately except in cases where, due to the methods used, it is technically impossible to separate information on gains and losses.
- ⁽⁴⁾ The signs for estimates of gains in carbon stocks are positive (+) and of losses in carbon stocks are negative (-).
- ⁽⁵⁾ Implied carbon-stock-change factors for mineral soils are calculated by dividing the net C stock change estimate for mineral soil by the difference between the area and the area of organic soil.
- ⁽⁶⁾ For category 5.B.1 Cropland remaining Cropland this column only includes changes in perennial woody biomass.
- ⁽⁷⁾ No reporting on dead organic matter pools is required for category 5.B.1. Cropland remaining Cropland.
- ⁽⁸⁾ When Parties are estimating fluxes for organic soils but cannot separate these fluxes from mineral soils, these fluxes should be reported under mineral soils.
- ⁽⁹⁾ The value reported for organic soils is estimated as a flux. For consistency with other entries in this column, these fluxes should be expressed in the unit required in this column, i.e. in Gg C.
- ⁽¹⁰⁾ According to the Revised 1996 IPCC Guidelines, for the purposes of reporting, the signs for removals are always negative (-) and for emissions positive (+). Net changes in carbon stocks are converted to CO₂ by multiplying C by 44/12 and changing the sign for net CO₂ removals to be negative (-) and for net CO₂ emissions to be positive (+). Note that carbon stock changes in a single pool are not necessarily equal to emissions or removals, because some carbon stock changes result from carbon transfers among pools rather than exchanges with the atmosphere.
- ⁽¹¹⁾ Where Parties directly estimate emissions and removals rather than carbon stock changes, they may report emissions/removals directly in this column and use notation keys in the stock change columns.
- ⁽¹²⁾ A Party may report aggregate estimates for all land conversions to cropland, when data are not available to report them separately. A Party should specify in the documentation box which types of land conversion are included. Separate estimates for forest land and grassland conversion should be provided in table 5 as an information item.

Documentation box:
Parties should provide detailed explanations on the Land Use, Land-Use Change and Forestry sector in Chapter 7: Land Use, Land-Use Change and Forestry (CRF sector 5) of the NIR. Use this documentation box to provide references to relevant sections of the NIR if any additional information and/or further details are needed to understand the content of this table.

National Inventory Report 2007 - ICELAND

TABLE 5.C SECTORAL BACKGROUND DATA FOR LAND USE, LAND-USE CHANGE AND FORESTRY

Inventory 2005

Grassland

Submission 2007 v1.1

(Sheet 1 of 1)

ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES		ACTIVITY DATA		IMPLIED CARBON-STOCK-CHANGE FACTORS						CHANGES IN CARBON STOCK						Net CO ₂ emissions/removals ^{(10) (11)}
Land-Use Category	Sub-division ⁽¹⁾	Area ⁽²⁾ (kha)	Area of organic soil ⁽²⁾ (kha)	Carbon stock change in living biomass per area ⁽⁴⁾			Net carbon stock change in dead organic matter per area ⁽⁴⁾	Net carbon stock change in soils per area ⁽⁴⁾		Carbon stock change in living biomass ^{(3), (4), (6)}			Net carbon stock change in dead organic matter ^{(4) (7)}	Net carbon stock change in soils ^{(4) (8)}		
				Gains	Losses	Net change		Mineral soils ⁽⁵⁾	Organic soils	Gains	Losses	Net change		Mineral soils	Organic soils ⁽⁹⁾	
				(Mg C/ha)						(Gg C)						
C. Total Grassland		3,843.90	446.47	NE,NO	NE,NO	NE,NO	NA,NE,NO	NE,NO	-1,10	NE,NO	NE,NO	NE,NO	NA,NE,NO	NE,NO	-491,11	1,800,74
1. Grassland remaining Grassland		3,843.90	446.47	NE	NE	NE	NA	NE	-1,10	NE	NE	NE	NA	NE	-491,11	1,800,74
2. Land converted to Grassland ⁽¹²⁾		NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO
2.1 Forest Land converted to Grassland		NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
2.2 Cropland converted to Grassland		NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
2.3 Wetlands converted to Grassland		NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
2.4 Settlements converted to Grassland		NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
2.5 Other Land converted to Grassland		NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE

⁽¹⁾ Land categories may be further divided according to climate zone, management system, soil type, vegetation type, tree species, ecological zone or national land classification.

⁽²⁾ The total area of the subcategories, in accordance with the sub-division used, should be entered here. For lands converted to Grassland report the cumulative area remaining in the category in the reporting year.

⁽³⁾ Carbon stock gains and losses should be listed separately except in cases where, due to the methods used, it is technically impossible to separate information on gains and losses.

⁽⁴⁾ The signs for estimates of gains in carbon stocks are positive (+) and of losses in carbon stocks are negative (-).

⁽⁵⁾ Implied carbon-stock-change factors for mineral soils are calculated by dividing the net C stock change estimate for mineral soil by the difference between the area and the area of organic soil.

⁽⁶⁾ For category 5.C.1 Grassland remaining Grassland this column only includes changes in perennial woody biomass.

⁽⁷⁾ No reporting on dead organic matter pools is required for category 5.C.1 Grassland remaining Grassland.

⁽⁸⁾ When Parties are estimating fluxes for organic soils but cannot separate these fluxes from mineral soils, these fluxes should be reported under mineral soils.

⁽⁹⁾ The value reported for organic soils is estimated as a flux. For consistency with other entries in this column, these fluxes should be expressed in the unit required in this column, i.e. in Gg C.

⁽¹⁰⁾ According to the Revised 1996 IPCC Guidelines, for the purposes of reporting, the signs for removals are always negative (-) and for emissions positive (+). Net changes in carbon stocks are converted to CO₂ by multiplying C by 44/12 and changing the sign for net CO₂ removals to be negative (-) and for net CO₂ emissions to be positive (+). Note that carbon stock changes in a single pool are not necessarily equal to emissions or removals, because some carbon stock changes result from carbon transfers among pools rather than exchanges with the atmosphere.

⁽¹¹⁾ Where Parties directly estimate emissions and removals rather than carbon stock changes, they may report emissions/removals directly in this column and use notation keys in the stock change columns.

⁽¹²⁾ A Party may report aggregate estimates for all land conversions to grassland, when data are not available to report them separately. A Party should specify in the documentation box which types of land conversion are included. Separate estimates for forest land conversion should be provided in table 5 as an information item.

Documentation box:

Parties should provide detailed explanations on the Land Use, Land-Use Change and Forestry sector in Chapter 7: Land Use, Land-Use Change and Forestry (CRF sector 5) of the NIR. Use this documentation box to provide references to relevant sections of the NIR if any additional information and/or further details are needed to understand the content of this table.

National Inventory Report 2007 - ICELAND

TABLE 5.D SECTORAL BACKGROUND DATA FOR LAND USE, LAND-USE CHANGE AND FORESTRY
Wetlands
(Sheet 1 of 1)

Inventory 2005
 Submission 2007 v1.1
 ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES		ACTIVITY DATA	IMPLIED CARBON-STOCK-CHANGE FACTORS					CHANGES IN CARBON STOCK					Net CO ₂ emissions/removals ⁽⁵⁾⁽⁶⁾
Land-Use Category	Sub-division ⁽¹⁾	Area ⁽²⁾ (kha)	Carbon stock change in living biomass per area ⁽³⁾ (4)			Net carbon stock change in dead organic matter per area ⁽⁴⁾	Net carbon stock change in soils per area ⁽⁴⁾	Carbon stock change in living biomass ⁽³⁾⁽⁴⁾			Net carbon stock change in dead organic matter ⁽⁴⁾	Net carbon stock change in soils ⁽⁴⁾	
			Gains	Losses	Net change			Gains	Losses	Net change			
			(Mg C/ha)					(Gg C)					
D. Total Wetlands		876,49	NE,NO	NE,NO	NE,NO	NE,NO	-0,04	NE,NO	NE,NO	NE,NO	NE,NO	-38,57	141,42
1. Wetlands remaining Wetlands ⁽⁷⁾		876,49	NE	NE	NE	NE	-0,04	NE	NE	NE	NE	-38,57	141,42
	Lakes and rivers	183,90	NE	NE	NE	NE	NO	NE	NE	NE	NE	NO	NE,NO
	Peatland	667,59	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
	Reservoirs	25,00	NE	NE	NE	NE	-1,54	NE	NE	NE	NE	-38,57	141,42
2. Land converted to Wetlands ⁽⁸⁾		NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO
	2.1 Forest Land converted to Wetlands	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
	2.2 Cropland converted to Wetlands	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
	2.3 Grassland converted to Wetlands	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
	2.4 Settlements converted to Wetlands	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
	2.5 Other Land converted to Wetlands	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE

- ⁽¹⁾ Land categories may be further divided according to climate zone, management system, soil type, vegetation type, tree species, ecological zone or national land classification.
- ⁽²⁾ The total area of the subcategories, in accordance with the sub-division used, should be entered here. For lands converted to Wetlands report the cumulative area remaining in the category in the reporting year.
- ⁽³⁾ Carbon stock gains and losses should be listed separately except in cases where, due to the methods used, it is technically impossible to separate information on gains and losses.
- ⁽⁴⁾ The signs for estimates of gains in carbon stocks are positive (+) and of losses in carbon stocks are negative (-).
- ⁽⁵⁾ According to the Revised 1996 IPCC Guidelines, for the purposes of reporting, the signs for removals are always negative (-) and for emissions positive (+). Net changes in carbon stocks are converted to CO₂ by multiplying C by 44/12 and changing the sign for net CO₂ removals to be negative (-) and for net CO₂ emissions to be positive (+). Note that carbon stock changes in a single pool are not necessarily equal to emissions or removals, because some carbon stock changes result from carbon transfers among pools rather than exchanges with the atmosphere.
- ⁽⁶⁾ Where Parties directly estimate emissions and removals rather than carbon stock changes, they may report emissions/removals directly in this column and use notation keys in the stock change columns.
- ⁽⁷⁾ Parties may decide not to prepare estimates for this category contained in appendix 3a.3 of the IPCC good practice guidance for LULUCF, although they may do so if they wish.
- ⁽⁸⁾ A Party may report aggregate estimates for all land conversions to wetlands, when data are not available to report them separately. A Party should specify in the documentation box which types of land conversion are included. Separate estimates for forest land and grassland conversion should be provided in table 5 as an information item.

Documentation box:
 Parties should provide detailed explanations on the Land Use, Land-Use Change and Forestry in Chapter 7: Land Use, Land-Use Change and Forestry (CRF sector 5) of the NIR. Use this documentation box to provide references to relevant sections of the NIR if any additional information and/or further details are needed to understand the content of this table.

National Inventory Report 2007 - ICELAND

TABLE 5.E SECTORAL BACKGROUND DATA FOR LAND USE, LAND-USE CHANGE AND FORESTRY Settlements
(Sheet 1 of 1)

Inventory 2005
Submission 2007 v1.1
ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES		ACTIVITY DATA	IMPLIED CARBON-STOCK-CHANGE FACTORS					CHANGES IN CARBON STOCK					Net CO ₂ emissions/removals ^{(6) (7)}
Land-Use Category	Sub-division ⁽¹⁾	Area ⁽²⁾ (kha)	Carbon stock change in living biomass per area ⁽³⁾ (4)			Net carbon stock change in dead organic matter per area ⁽⁴⁾	Net carbon stock change in soils per area ⁽⁴⁾	Carbon stock change in living biomass ^{(3), (4), (5)}			Net carbon stock change in dead organic matter ⁽⁴⁾	Net carbon stock change in soils ⁽⁴⁾	
			Gains	Losses	Net change			Gains	Losses	Net change			
			(Mg C/ha)					(Gg C)					
E. Total Settlements		68,45	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
1. Settlements remaining Settlements ⁽⁸⁾		68,45	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
2. Land converted to Settlements ⁽⁹⁾		NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
2.1 Forest Land converted to Settlements		NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
2.2 Cropland converted to Settlements		NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
2.3 Grassland converted to Settlements		NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
2.4 Wetlands converted to Settlements		NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
2.5 Other Land converted to Settlements		NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE

- ⁽¹⁾ Land categories may be further divided according to climate zone, management system, soil type, vegetation type, tree species, ecological zone or national land classification.
- ⁽²⁾ The total area of the subcategories, in accordance with the sub-division used, should be entered here. For lands converted to Settlements report the cumulative area remaining in the category in the reporting year.
- ⁽³⁾ Carbon stock gains and losses should be listed separately except in cases where, due to the methods used, it is technically impossible to separate information on gains and losses.
- ⁽⁴⁾ The signs for estimates of gains in carbon stocks are positive (+) and of losses in carbon stocks are negative (-).
- ⁽⁵⁾ For category 5.E.1 Settlements remaining Settlements this column only includes changes in perennial woody biomass.
- ⁽⁶⁾ According to the Revised 1996 IPCC Guidelines, for the purposes of reporting, the signs for removals are always negative (-) and for emissions positive (+). Net changes in carbon stocks are converted to CO₂ by multiplying C by 44/12 and changing the sign for net CO₂ removals to be negative (-) and for net CO₂ emissions to be positive (+). Note that carbon stock changes in a single pool are not necessarily equal to emissions or removals, because some carbon stock changes result from carbon transfers among pools rather than exchanges with the atmosphere.
- ⁽⁷⁾ Where Parties directly estimate emissions and removals rather than carbon stock changes, they may report emissions/removals directly in this column and use notation keys in the stock change columns.
- ⁽⁸⁾ Parties may decide not to prepare estimates for this category contained in appendix 3a.4 of the IPCC good practice guidance for LULUCF, although they may do so if they wish.
- ⁽⁹⁾ A Party may report aggregate estimates for all land conversions to settlements, when data are not available to report them separately. A Party should specify in the documentation box which types of land conversion are included. Separate estimates for forest land and grassland conversion should be provided in table 5 as an information item.

Documentation box:
Parties should provide detailed explanations on the Land Use, Land-Use Change and Forestry sector in Chapter 7: Land Use, Land-Use Change and Forestry (CRF sector 5) of the NIR. Use this documentation box to provide references to relevant sections of the NIR if any additional information and/or further details are needed to understand the content of this table.

National Inventory Report 2007 - ICELAND

TABLE 5.F. SECTORAL BACKGROUND DATA FOR LAND USE, LAND-USE CHANGE AND FORESTRY

Other land
(Sheet 1 of 1)

Inventory 2005
Submission 2007 v1.1
ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES		ACTIVITY DATA	IMPLIED CARBON-STOCK-CHANGE FACTORS					CHANGES IN CARBON STOCK					Net CO ₂ emissions/removals ⁽⁵⁾⁽⁶⁾
Land-Use Category	Sub-division ⁽¹⁾	Area ⁽²⁾ (kha)	Carbon stock change in living biomass per area ⁽³⁾ (4)			Net carbon stock change in dead organic matter per area ⁽⁴⁾	Net carbon stock change in soils per area ⁽⁴⁾	Carbon stock change in living biomass ⁽³⁾⁽⁴⁾			Net carbon stock change in dead organic matter ⁽⁴⁾	Net carbon stock change in soils ⁽⁴⁾	
			Gains	Losses	Net change			Gains	Losses	Net change			
			(Mg C/ha)						(Gg C)				
F. Total Other Land		5,310.06	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
1. Other Land remaining Other Land ⁽⁷⁾		5,310.06											
2. Land converted to Other Land ⁽⁸⁾		NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
2.1 Forest Land converted to Other Land		NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
2.2 Cropland converted to Other Land		NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
2.3 Grassland converted to Other Land		NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
2.4 Wetlands converted to Other Land		NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
2.5 Settlements converted to Other Land		NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE

⁽¹⁾ Land categories may be further divided according to climate zone, management system, soil type, vegetation type, tree species, ecological zone or national land classification.

⁽²⁾ The total area of the subcategories, in accordance with the sub-division used, should be entered here. For lands converted to Other Land report the cumulative area remaining in the category in the reporting year.

⁽³⁾ Carbon stock gains and losses should be listed separately except in cases where, due to the methods used, it is technically impossible to separate information on gains and losses.

⁽⁴⁾ The signs for estimates of gains in carbon stocks are positive (+) and of losses in carbon stocks are negative (-).

⁽⁵⁾ According to the Revised 1996 IPCC Guidelines, for the purposes of reporting, the signs for removals are always negative (-) and for emissions positive (+). Net changes in carbon stocks are converted to CO₂ by multiplying C by 44/12 and changing the sign for net CO₂ removals to be negative (-) and for net CO₂ emissions to be positive (+). Note that carbon stock changes in a single pool are not necessarily equal to emissions or removals, because some carbon stock changes result from carbon transfers among pools rather than exchanges with the atmosphere.

⁽⁶⁾ Where Parties directly estimate emissions and removals rather than carbon stock changes, they may report emissions/removals directly in this column and use notation keys in the stock change columns.

⁽⁷⁾ This land-use category is to allow the total of identified land area to match the national area.

⁽⁸⁾ A Party may report aggregate estimates for all land conversions to other land, when data are not available to report them separately. A Party should specify in the documentation box which types of land conversion are included. Separate estimates for forest land and grassland conversion should be provided in table 5 as an information item.

Documentation box:

Parties should provide detailed explanations on the Land Use, Land-Use Change and Forestry sector in Chapter 7: Land Use, Land-Use Change and Forestry (CRF sector 5) of the NIR. Use this documentation box to provide references to relevant sections of the NIR if any additional information and/or further details are needed to understand the content of this table.

National Inventory Report 2007 - ICELAND

Inventory 2005
Submission 2007 v1.1
ICELAND

TABLE 5 (I) SECTORAL BACKGROUND DATA FOR LAND USE, LAND-USE CHANGE AND FORESTRY
Direct N₂O emissions from N fertilization⁽¹⁾ of Forest Land and Other
(Sheet 1 of 1)

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	ACTIVITY DATA	IMPLIED EMISSION FACTORS	EMISSIONS ⁽⁴⁾
Land-Use Category ⁽²⁾	Total amount of fertilizer applied (Gg N/yr)	N ₂ O-N emissions per unit of fertilizer (kg N ₂ O-N/kg N) ⁽³⁾	N ₂ O (Gg)
Total for all Land Use Categories	0,83	0,01	0,02
A. Forest Land ⁽⁵⁾⁽⁶⁾	0,02	0,01	0,00
1. Forest Land remaining Forest Land	0,00	0,01	0,00
2. Land converted to Forest Land	0,02	0,01	0,00
G. Other (please specify)			
Revegetation	0,81	0,01	0,02
Grassland organic soil	NA	NA	NA

⁽¹⁾ Direct N₂O emissions from fertilization are estimated using equations 3.2.17 and 3.2.18 of the IPCC good practice guidance for LULUCF based on the amounts of fertilizers applied to forest land.

⁽²⁾ N₂O emissions from N fertilization of cropland and grassland are reported in the Agriculture sector; therefore only Forest Land is included in this table.

⁽³⁾ In the calculation of the implied emission factor, N₂O emissions are converted to N₂O-N by multiplying by 28/44.

⁽⁴⁾ Emissions are reported with a positive sign.

⁽⁵⁾ If a Party is not able to separate the fertilizer applied to forest land from that applied to agriculture, it may report all N₂O emissions from fertilization in the Agriculture sector. This should be explicitly indicated in the documentation box.

⁽⁶⁾ A Party may report aggregate estimates for all N fertilization on forest land in the category Forest Land remaining Forest Land when data are not available to report Forest Land remaining Forest Land and Land converted to Forest Land separately.

Documentation box:

Parties should provide detailed explanations on the Land Use, Land-Use Change and Forestry sector in Chapter 7: Land Use, Land-Use Change and Forestry (CRF sector 5) of the NIR. Use this documentation box to provide references to relevant sections of the NIR if any additional information and/or further details are needed to understand the content of this table.

5 5(I) Direct N2O emissions from N fertilization of Forest Land and Other:All N-fertilizers are included in Agricultural sector. Data not available for amount of fertilisers used in forestry or revegetation.

5.A.1 5(I) Direct N2O emissions from N fertilization of Forest Land and Other/2005:No separation of fertilizers used in forestry and agriculture available for this year

National Inventory Report 2007 - ICELAND

TABLE 5 (II) SECTORAL BACKGROUND DATA FOR LAND USE, LAND-USE CHANGE AND FORESTRY

Non-CO₂ emissions from drainage of soils and wetlands⁽¹⁾

(Sheet 1 of 1)

Inventory 2005

Submission 2007 v1.1

ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES		ACTIVITY DATA	IMPLIED EMISSION FACTORS		EMISSIONS ⁽⁵⁾	
Land-Use Category ⁽²⁾	Sub-division ⁽³⁾	Area (kha)	N ₂ O-N per area ⁽⁴⁾ (kg N ₂ O-N/ha)	CH ₄ per area (kg CH ₄ /ha)	N ₂ O	CH ₄
					(Gg)	
Total all Land-Use Categories					1,27	NA,NE,NO
A. Forest Land⁽⁶⁾			0,60	NA,NE	0,00	NA,NE
Organic Soil		3,53	0,60	NA	0,00	NA
Mineral Soil		NE	NE	NE	NE	NE
D. Wetlands			NE	NE	NE	NE
Peatland ⁽⁷⁾		NE	NE	NE	NE	NE
Flooded Lands ⁽⁷⁾		NE	NE	NE	NE	NE
G. Other (please specify)					1,26	NA,NO
Revegetation		NO	NO	NO	NO	NO
Organic Soil		NO	NO	NO	NO	NO
Mineral Soil		NO	NO	NO	NO	NO
Grassland organic soil		446,47	1,80	NA,NO	1,26	NA,NO
Organic Soil		446,47	1,80	NA	1,26	NA
Mineral Soil		NO	NO	NO	NO	NO

⁽¹⁾ Parties may decide not to prepare estimates for these categories contained in appendices 3a.2 and 3a.3 of the IPCC good practice guidance for LULUCF, although they may do so if they wish.

⁽²⁾ N₂O emissions from drained cropland and grassland soils are covered in the Agriculture tables of the CRF under Cultivation of Histosols.

⁽³⁾ A Party should report further disaggregations of drained soils corresponding to the methods used. Tier 1 disaggregates soils into "nutrient rich" and "nutrient poor" areas, whereas higher-tier methods can further disaggregate into different peatland types, soil fertility or tree species.

⁽⁴⁾ In the calculation of the implied emission factor, N₂O emissions are converted to N₂O-N by multiplying by 28/44.

⁽⁵⁾ Emissions are reported with a positive sign.

⁽⁶⁾ In table 5, these emissions will be added to 5.A.1 Forest Land remaining Forest Land.

⁽⁷⁾ In table 5, these emissions will be added to 5.D.2 Land converted to Wetlands.

Documentation box:

Parties should provide detailed explanations on the Land Use, Land-Use Change and Forestry sector in Chapter 7: Land Use, Land-Use Change and Forestry (CRF sector 5) of the NIR. Use this documentation box to provide references to relevant sections of the NIR if any additional information and/or further details are needed to understand the content of this table.

National Inventory Report 2007 - ICELAND

TABLE 5 (III) SECTORAL BACKGROUND DATA FOR LAND USE, LAND-USE CHANGE AND FORESTRY
N₂O emissions from disturbance associated with land-use conversion to cropland ⁽¹⁾
(Sheet 1 of 1)

Inventory 2005
 Submission 2007 v1.1
 ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	ACTIVITY DATA	IMPLIED EMISSION FACTORS	EMISSIONS ⁽⁴⁾
Land-Use Category ⁽²⁾	Land area converted	N ₂ O-N emissions per area converted ⁽³⁾	N ₂ O
	(kha)	(kg N ₂ O-N/ha)	(Gg)
Total all Land-Use Categories ⁽⁵⁾	NE	NE	NE
B. Cropland	NE	NE	NE
2. Lands converted to Cropland ⁽⁶⁾	NE	Value is sum of Cells: B14, B17, B20, B23.	NE
Organic Soils	NE		NE
Mineral Soils	NE		NE
2.1 Forest Land converted to Cropland	NE	NE	NE
Organic Soils	NE	NE	NE
Mineral Soils	NE	NE	NE
2.2 Grassland converted to Cropland	NE	NE	NE
Organic Soils	NE	NE	NE
Mineral Soils	NE	NE	NE
2.3 Wetlands converted to Cropland ⁽⁷⁾	NE	NE	NE
Organic Soils	NE	NE	NE
Mineral Soils	NE	NE	NE
2.5 Other Land converted to Cropland	NE	NE	NE
Organic Soils	NE	NE	NE
Mineral Soils	NE	NE	NE
G. Other (please specify)			
Revegetation	NE	NE	NE
Organic Soils	NE	NE	NE
Mineral Soils	NE	NE	NE
Grassland organic soil	NE	NE	NE
Organic Soils	NE	NE	NE
Mineral Soils	NE	NE	NE

⁽¹⁾ Methodologies for N₂O emissions from disturbance associated with land-use conversion are based on equations 3.3.14 and 3.3.15 of the IPCC good practice guidance for LULUCF. N₂O emissions from fertilization in the preceding land use and new land use should not be reported.

⁽²⁾ According to the IPCC good practice guidance for LULUCF, N₂O emissions from disturbance of soils are only relevant for land conversions to cropland. N₂O emissions from Cropland remaining Cropland are included in the Agriculture sector of the good practice guidance. The good practice guidance provides methodologies only for mineral soils.

⁽³⁾ In the calculation of the implied emission factor, N₂O emissions are converted to N₂O-N by multiplying by 28/44.

⁽⁴⁾ Emissions are reported with a positive sign.

⁽⁵⁾ Parties can separate between organic and mineral soils, if they have data available.

⁽⁶⁾ If activity data cannot be disaggregated to all initial land uses, Parties may report some initial land uses aggregated under Other Land converted to Cropland (indicate in the documentation box what this category includes).

⁽⁷⁾ Parties should avoid double counting with N₂O emissions from drainage and from cultivation of organic soils reported in Agriculture under Cultivation of Histosols.

Documentation box:

Parties should provide detailed explanations on the Land Use, Land-Use Change and Forestry sector in Chapter 7: Land Use, Land-Use Change and Forestry (CRF Sector 5) of the NIR. Use this documentation box to provide references to relevant sections of the NIR if any additional information and/or further details are needed to understand the content of this table.

5 5(III) N₂O emissions from disturbance associated with land-use conversion to cropland: Information on land converted to cropland not available although occurring. Land has been converted to cropland from grassland, other land and wetland. No estimates on area converted available.

National Inventory Report 2007 - ICELAND

TABLE 5 (IV) SECTORAL BACKGROUND DATA FOR LAND USE, LAND-USE CHANGE AND FORESTRY
CO₂ emissions from agricultural lime application ⁽¹⁾
(Sheet 1 of 1)

Inventory 2005
 Submission 2007 v1.1
 ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	ACTIVITY DATA	IMPLIED EMISSION FACTORS	EMISSIONS ⁽³⁾
Land-Use Category	Total amount of lime applied (Mg/yr)	CO ₂ -C per unit of lime ⁽²⁾ (Mg CO ₂ -C /Mg)	CO ₂ (Gg)
Total all Land-Use Categories ^{(4), (5), (6)}	8.483,81	0,11	3,46
B. Cropland ^{(6) (7)}	8.483,81	0,11	3,46
Limestone CaCO ₃	2.662,09	0,12	1,17
Dolomite CaMg(CO ₃) ₂	135,60	0,07	0,03
shellsand (90% CaCO ₃)	5.686,12	0,11	2,25
C. Grassland ^{(6) (8)}	NO	NO	NO
Limestone CaCO ₃	NO	NO	NO
Dolomite CaMg(CO ₃) ₂	NO	NO	NO
G. Other (please specify) ^{(6) (9)}			
Revegetation	NA,NO	C,NA	C,NA
Limestone CaCO ₃	NO	C	C
Dolomite CaMg(CO ₃) ₂	NA	NA	NA
Grassland organic soil	NO	NO	NO
Limestone CaCO ₃	NO	NO	NO
Dolomite CaMg(CO ₃) ₂	NO	NO	NO

⁽¹⁾ CO₂ emissions from agricultural lime application are addressed in equations 3.3.6 and 3.4.11 of the IPCC good practice guidance for LULUCF.

⁽²⁾ The implied emission factor is expressed in unit of carbon to facilitate comparison with published emission factors.

⁽³⁾ Emissions are reported with a positive sign.

⁽⁴⁾ If Parties are not able to separate liming application for different land-use categories, they should include liming for all land-use categories in the category 5.G Other.

⁽⁵⁾ Parties that are able to provide data for lime application to forest land should provide this information under 5.G Other and specify in the documentation box that forest land application is included in this category.

⁽⁶⁾ A Party may report aggregate estimates for total lime applications when data are not available for limestone and dolomite.

⁽⁷⁾ In table 5, these CO₂ emissions will be added to 5.B.1 Cropland remaining Cropland.

⁽⁸⁾ In table 5, these CO₂ emissions will be added to 5.C.1 Grassland remaining Grassland.

⁽⁹⁾ If a Party has data broken down to limestone and dolomite at national level, it can report these data under 5.G Other.

Documentation box:

Parties should provide detailed explanations on the Land Use, Land-Use Change and Forestry sector in Chapter 7: Land Use, Land-Use Change and Forestry (CRF sector 5) of the NIR. Use this documentation box to provide references to relevant sections of the NIR if any additional information and/or further details are needed to understand the content of this table.

5.5(IV) CO₂ emissions from agricultural lime application:Liming information only available on national level

National Inventory Report 2007 - ICELAND

TABLE 5 (V) SECTORAL BACKGROUND DATA FOR LAND USE, LAND-USE CHANGE AND FORESTRY
Biomass Burning⁽¹⁾
(Sheet 1 of 1)

Inventory 2005
 Submission 2007 v1.1
 ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	ACTIVITY DATA			IMPLIED EMISSION FACTOR			EMISSIONS ⁽⁵⁾		
	Description ⁽³⁾	Unit (ha or kg dm)	Values	CO ₂	CH ₄	N ₂ O	CO ₂ ⁽⁴⁾	CH ₄	N ₂ O
Land-Use Category ⁽²⁾				(Mg/activity data unit)			(Gg)		
Total for Land-Use Categories	Area burned	ha	NA,NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO
A. Forest Land			NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO
1. Forest land remaining Forest Land			NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO
Controlled Burning	(specify)		NO	NO	NO	NO	NO	NO	NO
Wildfires	(specify)		NE	NE	NE	NE	NE	NE	NE
2. Land converted to Forest Land			NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO
Controlled Burning	(specify)		NO	NO	NO	NO	NO	NO	NO
Wildfires	(specify)		NE	NE	NE	NE	NE	NE	NE
B. Cropland	Area burned	ha	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO
1. Cropland remaining Cropland ⁽⁶⁾			NE	NE	NE	NE	NE	NE	NE
Controlled Burning	Area burned	ha	NE	NE	NE	NE	NE	NE	NE
Wildfires	Area burned	ha	NE	NE	NE	NE	NE	NE	NE
2. Land converted to Cropland			NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO
Controlled Burning	(specify)		NO	NO	NO	NO	NO	NO	NO
Wildfires	(specify)		NE	NE	NE	NE	NE	NE	NE
2.1. Forest Land converted to Cropland			NO	NO	NO	NO	NO	NO	NO
Controlled Burning	(specify)		NO	NO	NO	NO	NO	NO	NO
Wildfires	(specify)		NO	NO	NO	NO	NO	NO	NO
C. Grassland			NE	NE	NE	NE	NE	NE	NE
1. Grassland remaining grassland ⁽⁷⁾			NE	NE	NE	NE	NE	NE	NE
Controlled Burning	(specify)		NE	NE	NE	NE	NE	NE	NE
Wildfires	(specify)		NE	NE	NE	NE	NE	NE	NE
2. Land converted to Grassland			NE	NE	NE	NE	NE	NE	NE
Controlled Burning	(specify)		NE	NE	NE	NE	NE	NE	NE
Wildfires	(specify)		NE	NE	NE	NE	NE	NE	NE
2.1. Forest Land converted to Grassland			NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO
Controlled Burning	(specify)		NO	NO	NO	NO	NO	NO	NO
Wildfires	(specify)		NO	NE	NE	NE	NE	NE	NE
D. Wetlands	Area burned	ha	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO
1. Wetlands remaining Wetlands ⁽⁸⁾			NE	NE	NE	NE	NE	NE	NE
Controlled Burning	Area burned	ha	NE	NE	NE	NE	NE	NE	NE
Wildfires	Area burned	ha	NE	NE	NE	NE	NE	NE	NE
2. Land converted to Wetlands			NO	NO	NO	NO	NO	NO	NO
Controlled Burning	Area burned	ha	NO	NO	NO	NO	NO	NO	NO
Wildfires	(specify)		NO	NO	NO	NO	NO	NO	NO
2.1. Forest Land converted to Wetlands			NO	NO	NO	NO	NO	NO	NO
Controlled Burning	(specify)		NO	NO	NO	NO	NO	NO	NO
Wildfires	(specify)		NO	NO	NO	NO	NO	NO	NO
E. Settlements⁽⁸⁾			NE	NE	NE	NE	NE	NE	NE
F. Other Land⁽⁹⁾	Area burned	ha	NA	NE	NE	NE	NE	NE	NE
G. Other (please specify)									
Revegetation			NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO
Controlled Burning			NO	NO	NO	NO	NO	NO	NO
Wildfires			NE	NE	NE	NE	NE	NE	NE
Grassland organic soil			NE	NE	NE	NE	NE	NE	NE
Controlled Burning			NE	NE	NE	NE	NE	NE	NE
Wildfires			NE	NE	NE	NE	NE	NE	NE

⁽¹⁾ Methodological guidance on burning can be found in sections 3.2.1.4 and 3.4.1.3 of the IPCC good practice guidance for LULUCF.
⁽²⁾ Parties should report both controlled/prescribed burning and wildfires emissions, where appropriate, in a separate manner.
⁽³⁾ For each category activity data should be selected between area burned or biomass burned. Units for area will be ha and for biomass burned kg dm. The implied emission factor will refer to the selected activity data with an automatic change in the units.
⁽⁴⁾ If CO₂ emissions from biomass burning are not already included in tables 5.A - 5.F, they should be reported here. This should be clearly documented in the documentation box and in the NIR. Double counting should be avoided. Parties that include all carbon stock changes in the carbon stock tables (5.A, 5.B, 5.C, 5.D, 5.E and 5.F), should report IE (included elsewhere) in this column.
⁽⁵⁾ Emissions are reported with a positive sign.
⁽⁶⁾ In-situ above-ground woody biomass burning is reported here. Agricultural residue burning is reported in the Agriculture sector.
⁽⁷⁾ Includes only emissions from controlled biomass burning on grasslands outside the tropics (prescribed savanna burning is reported under the Agriculture sector).
⁽⁸⁾ Parties may decide not to prepare estimates for these categories contained in appendices 3a.2, 3a.3 and 3a.4 of the IPCC good practice guidance for LULUCF, although they may do so if they wish.
⁽⁹⁾ This land-use category is to allow the total of identified land area to match the national area.

Documentation box:
 Parties should provide detailed explanations on the Land Use, Land-Use Change and Forestry sector in Chapter 7: Land Use, Land-Use Change and Forestry (CRF sector 5) of the NIR. Use this documentation box to provide references to relevant sections of the NIR if any additional information and/or further details are needed to understand the content of this table.

5.5(V) Biomass Burning: Biomass burning taking place but data not available on total area or biomass burned.

National Inventory Report 2007 - ICELAND

TABLE 6 SECTORAL REPORT FOR WASTE
(Sheet 1 of 1)

Inventory 2005
Submission 2007 v1.1
ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	CO ₂ ⁽¹⁾	CH ₄	N ₂ O	NO _x	CO	NMVOC	SO ₂
	(Gg)						
Total Waste	0,03	7,60	0,02	0,00	0,00	0,00	0,00
A. Solid Waste Disposal on Land	NE,NO	7,33		NE	NE	NE	
1. Managed Waste Disposal on Land	NE	3,55		NE	NE	NE	
2. Unmanaged Waste Disposal Sites	NE,NO	3,36		NE	NE	NE	
3. Other (as specified in table 6.A)		0,41		NE	NE	NE	
Uncategorized		0,41		NE	NE	NE	
B. Waste Water Handling		0,27	0,02	NA,NE	NA,NE	NA,NE	
1. Industrial Wastewater		NE	NE	NE	NE	NE	
2. Domestic and Commercial Waste Water		0,27	0,02	NE	NE	NE	
3. Other (as specified in table 6.B)		NA	NA	NA	NA	NA	
C. Waste Incineration	0,03	NE	0,00	0,00	0,00	0,00	0,00
D. Other (please specify)	NA	NA	NA	NA	NA	NA	NA

⁽¹⁾ CO₂ emissions from source categories Solid waste disposal on land and Waste incineration should only be included if they derive from non-biological or inorganic waste sources.

Documentation box:

- Parties should provide detailed explanations on the waste sector in Chapter 8: Waste (CRF sector 6) of the NIR. Use this documentation box to provide references to relevant sections of the NIR if any additional information and/or further details are needed to understand the content of this table.
- If estimates are reported under "6.D Other", use this documentation box to provide information regarding activities covered under this category and to provide reference to the section in the NIR where background information can be found.

National Inventory Report 2007 - ICELAND

TABLE 6.A SECTORAL BACKGROUND DATA FOR WASTE

Solid Waste Disposal

(Sheet 1 of 1)

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	ACTIVITY DATA AND OTHER RELATED INFORMATION			IMPLIED EMISSION FACTOR		EMISSIONS		
	Annual MSW at the SWDS (Gg)	MCF	DOC degraded %	CH ₄ ⁽¹⁾	CO ₂	CH ₄		CO ₂ ⁽⁴⁾
				(t/t MSW)		Emissions ⁽²⁾	Recovery ⁽³⁾	
					(Gg)			
1 Managed Waste Disposal on Land	180,00	1,00	0,00	0,03	NE	3,55	2,40	NE
2 Unmanaged Waste Disposal Sites			0,00		NE,NO	3,36	NO	NE,NO
a. Deep (>5 m)	NO	NO	NO	NO	NO	NO	NO	NO
b. Shallow (<5 m)	172,75	0,40	0,00	0,02	NE	3,36	NO	NE
3 Other (<i>please specify</i>)						0,41	NO	
Uncategorized	NO	0,60	0,00	NO	NO	0,41	NO	

Note: MSW - Municipal Solid Waste, SWDS - Solid Waste Disposal Site, MCF - Methane Correction Factor, DOC - Degradable Organic Carbon (IPCC Guidelines (Volume 3. Reference Manual, section 6.2.4)).

MSW includes household waste, yard/garden waste, commercial/market waste and organic industrial solid waste. MSW should not include inorganic industrial waste such as construction or demolition materials.

⁽¹⁾ The CH₄ implied emission factor (IEF) is calculated on the basis of gross CH₄ emissions, as follows: IEF = (CH₄ emissions + CH₄ recovered)/annual MSW at the SWDS.

⁽²⁾ Actual emissions (after recovery).

⁽³⁾ CH₄ recovered and flared or utilized.

⁽⁴⁾ Under Solid Waste Disposal, CO₂ emissions should be reported only when the disposed waste is combusted at the disposal site as a management practice. CO₂ emissions from non-biogenic wastes are included in the total emissions, whereas the CO₂ emissions from biogenic wastes are not included in the total emissions.

TABLE 6.C SECTORAL BACKGROUND DATA FOR WASTE

Waste Incineration

(Sheet 1 of 1)

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	ACTIVITY DATA Amount of incinerated wastes (Gg)	IMPLIED EMISSION FACTOR			EMISSIONS		
		CO ₂	CH ₄	N ₂ O	CO ₂ ⁽¹⁾	CH ₄	N ₂ O
		(kg/t waste)			(Gg)		
Waste Incineration	0,05				0,03	NE	0,00
a. Biogenic ⁽¹⁾	0,04	995,24	NE	0,10	0,04	NE	0,00
b. Other (non-biogenic - <i>please specify</i>) ^{(1),(2)}					0,03	NE	0,00
Plastics and other non-biogenic waste	0,01	3.483,33	NE	0,10	0,03	NE	0,00

⁽¹⁾ Under Solid Waste Disposal, CO₂ emissions should be reported only when the disposed waste is combusted at the disposal site as a management practice. CO₂ emissions from non-biogenic wastes are included in the total emissions, while the CO₂ emissions from biogenic wastes are not included in the total emissions.

⁽²⁾ Enter under this source category all types of non-biogenic wastes, such as plastics.

Note: Only emissions from waste incineration without energy recovery are to be reported in the Waste sector. Emissions from incineration with energy recovery are to be reported in the Energy sector, as Other Fuels (see IPCC good practice guidance, page 5.23).

Documentation box:

- Parties should provide detailed explanations on the waste sector in Chapter 8: Waste (CRF sector 6) of the NIR. Use this documentation box to provide references to relevant sections of the NIR if any additional information and/or further
- Parties that use country-specific models should provide a reference in the documentation box to the relevant section in the NIR where these models are described, and fill in only the relevant cells of tables 6.A and 6.C.
- Provide a reference to the relevant section in the NIR, in particular with regard to:
 - (a) A population size (total or urban population) used in the calculations and the rationale for doing so;
 - (b) The composition of landfilled waste;
 - (c) In relation to the amount of incinerated wastes (specify whether the reported data relate to wet or dry matter).

National Inventory Report 2007 - ICELAND

Inventory 2005
Submission 2007 v1.1
ICELAND

TABLE 6.B SECTORAL BACKGROUND DATA FOR WASTE
Waste Water Handling
(Sheet 1 of 2)

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	ACTIVITY DATA AND RELATED INFORMATION ⁽¹⁾		IMPLIED EMISSION FACTOR		EMISSIONS		
	Total organic product	CH ₄ ⁽²⁾	N ₂ O ⁽³⁾	CH ₄		N ₂ O ⁽³⁾	
				Emissions ⁽⁴⁾	Recovery ⁽⁵⁾		
(Gg DC ⁽¹⁾ /yr)	(kg/kg DC)	(Gg)					
1. Industrial Waste Water				NE	NE	NE	
a. Waste Water	NE	NE	NE	NE	NE	NE	
b. Sludge	NE	NE	NE	NE	NE	NE	
2. Domestic and Commercial Wastewater				0,27	NE,NO	0,02	
a. Waste Water	NE	NE	NE	0,27	NO	NE	
b. Sludge	NE	NE	NE	NE	NE	NE	
3. Other (please specify) ⁽⁶⁾				NA	NA	NA	

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	ACTIVITY DATA AND OTHER RELATED INFORMATION			IMPLIED EMISSION FACTOR	EMISSIONS
	Population (1000s)	Protein consumption (kg/person/yr)	N fraction (kg N/kg protein)	N ₂ O (kg N ₂ O-N/kg sewage N produced)	N ₂ O (Gg)
N ₂ O from human sewage ⁽³⁾	299,40	31,76	0,16	0,01	0,02

⁽¹⁾ DC - degradable organic component. DC indicators are COD (Chemical Oxygen Demand) for industrial waste water and BOD (Biochemical Oxygen Demand) for Domestic/Commercial waste water/sludge (IPCC Guidelines (Volume 3. Reference Manual, pp. 6.14, 6.18)).

⁽²⁾ The CH₄ implied emission factor (IEF) is calculated on the basis of gross CH₄ emissions, as follows: IEF = (CH₄ emissions + CH₄ recovered or flared) / total organic product.

⁽³⁾ Parties using methods other than those from the IPCC for estimating N₂O emissions from human sewage or waste-water treatment should provide aggregate data in this table.

⁽⁴⁾ Actual emissions (after recovery).

⁽⁵⁾ CH₄ recovered and flared or utilized.

⁽⁶⁾ Use the cells below to specify each activity covered under "6.B.3 Other". Note that under each reported activity, data for waste water and sludge are to be reported separately.

Documentation box:

- Parties should provide detailed explanations on the Waste sector in Chapter 8: Waste (CRF sector 6) of the NIR. Use this documentation box to provide references to relevant sections of the NIR if any additional information and/or further details are needed to understand the content of this table.
- Regarding the estimates for N₂O from human sewage, specify whether total or urban population is used in the calculations and the rationale for doing so. Provide explanation in the documentation box.
- Parties using methods other than those from the IPCC for estimating N₂O emissions from human sewage or waste-water treatment should provide, in the NIR, corresponding information on methods, activity data and emission factors used, and should provide a reference to the relevant section of the NIR in this documentation box.

TABLE 6.B SECTORAL BACKGROUND DATA FOR WASTE
Waste Water Handling
(Sheet 2 of 2)

Inventory 2005
 Submission 2007 v1.1
 ICELAND

Additional information

	Domestic	Industrial
Total waste water (m ³):	NE	NE
Treated waste water (%):	60,70	NE

Waste-water streams:	Waste-water output (m ³)	DC (kg COD/m ³)
Industrial waste water	NE	NE
Iron and steel		
Non-ferrous		
Fertilizers		
Food and beverage		
Paper and pulp		
Organic chemicals		
Other (please specify)		
Textile		
Rubber		
Poultry		
Wood and wood production		
Wool Scouring		
Other agricultural		
Chemical		
Dairy Processing		
Electricity, steam, water production		
Leather industry		
Leather and Skins		
Iron and steel		
Meat industry		
Fuels		
Machinery and equipment		
Mining and quarrying		
DC (kg BOD/1000 person/yr)		
Domestic and Commercial	NE	
Other (please specify)		

Handling systems:	Industrial waste water treated (%)	Industrial sludge treated (%)	Domestic waste water treated (%)	Domestic sludge treated (%)
Aerobic	0,00	0,00	0,00	0,00
Anaerobic	0,00	0,00	0,00	0,00
Other (please specify)	0,00	0,00	0,00	0,00

National Inventory Report 2007 - ICELAND

SUMMARY 1.A SUMMARY REPORT FOR NATIONAL GREENHOUSE GAS INVENTORIES (IPCC TABLE 7A) (Sheet 1 of 3)

Inventory 2005
Submission 2007 v1.1
ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	Net CO ₂	CH ₄	N ₂ O	HFCs ⁽¹⁾		PFCs ⁽¹⁾		SF ₆		NO _x	CO	NMVOC	SO ₂
	emissions/removals			P	A	P	A	P	A				
	(Gg)			CO ₂ equivalent (Gg)						(Gg)			
Total National Emissions and Removals	4,158,24	22,15	2,35	76,74	NA,NE,NO	NA,NE,NO	26,09	0,00	NA,NE,NO	27,83	19,58	8,29	40,53
1. Energy	1,913,62	0,16	0,24							26,65	19,35	4,06	2,48
A. Fuel Combustion	1,880,77												
Reference Approach ⁽²⁾	1,880,77												
Sectoral Approach ⁽²⁾	1,913,62	0,16	0,24							26,65	19,35	4,06	2,48
1. Energy Industries	22,60	0,00	0,00							0,24	0,07	0,00	0,05
2. Manufacturing Industries and Construction	442,28	0,02	0,11							4,52	1,35	0,59	1,23
3. Transport	716,38	0,08	0,12							4,66	16,21	2,97	0,10
4. Other Sectors	716,26	0,07	0,02							17,18	1,72	0,50	0,93
5. Other	16,11	0,00	0,00							0,04	0,00	0,00	0,16
B. Fugitive Emissions from Fuels	NA,NE,NO	NA,NE,NO	NA,NO							NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
1. Solid Fuels	NA,NO	NA,NO	NA,NO							NA,NO	NA,NO	NA,NO	NA,NO
2. Oil and Natural Gas	NA,NE,NO	NA,NE,NO	NA,NO							NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
2. Industrial Processes	835,11	0,05	NA,NE,NO	76,74	NA,NE,NO	NA,NE,NO	26,09	0,00	NA,NE,NO	1,18	0,23	0,10	5,68
A. Mineral Products	55,10	NA,NE	NA,NE							0,02	0,05	0,00	0,03
B. Chemical Industry		NE,NO	NE,NO	NA	NA	NA	NA	NA	NA		NE,NO	NE,NO	NE,NO
C. Metal Production	780,00	0,05	NA				26,09		NA,NO	1,17	0,18	0,09	5,65
D. Other Production ⁽³⁾	NE									NO	NO	NE,NO	NO
E. Production of Halocarbons and SF ₆					NA,NO		NA,NO		NA,NO				
F. Consumption of Halocarbons and SF ₆				76,74	NA,NE,NO	NE,NO	NA,NE,NO	0,00	NA,NE,NO				
G. Other	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

Note: A = Actual emissions based on Tier 2 approach of the IPCC Guidelines.
P = Potential emissions based on Tier 1 approach of the IPCC Guidelines.

Note: All footnotes for this table are given at the end of the table on sheet 3.

National Inventory Report 2007 - ICELAND

SUMMARY 1.A SUMMARY REPORT FOR NATIONAL GREENHOUSE GAS INVENTORIES (IPCC TABLE 7A) (Sheet 2 of 3)

Inventory 2005
Submission 2007 v1.1
ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	Net CO ₂ emissions/removals	CH ₄	N ₂ O	HFCs ⁽¹⁾		PFCs ⁽¹⁾		SF ₆		NO _x	CO	NMVOC	SO ₂
				P	A	P	A	P	A				
	(Gg)				CO ₂ equivalent (Gg)						(Gg)		
3. Solvent and Other Product Use	NE		0,01							NE	NE	4,14	NE
4. Agriculture		12,00	0,72							NA,NO	NA,NO	NA,NE,NO	NO
A. Enteric Fermentation		11,02											
B. Manure Management		0,97	0,08									NE,NO	
C. Rice Cultivation		NA,NO										NA,NO	
D. Agricultural Soils ⁽⁴⁾		NA,NE	0,64									NA,NE	
E. Prescribed Burning of Savannas		NA	NA							NO	NO	NO	
F. Field Burning of Agricultural Residues		NA,NO	NA,NO							NA,NO	NA,NO	NA,NO	
G. Other		NA	NA							NA	NA	NA	NO
5. Land Use, Land-Use Change and Forestry	⁽⁵⁾ 1.286,10	2,34	1,35							NA,NE	NA,NE	NA,NE	NE
A. Forest Land	⁽⁵⁾ -126,27	NA,NE,NO	0,00							NE	NE	NE	
B. Cropland	⁽⁵⁾ 3,46	NE,NO	NE,NO							NE	NE	NE	
C. Grassland	⁽⁵⁾ 1.800,74	NE	NE							NE	NE	NE	
D. Wetlands	⁽⁵⁾ 141,42	2,34	0,07							NA,NE	NA,NE	NA,NE	
E. Settlements	⁽⁵⁾ NE	NE	NE							NE	NE	NE	
F. Other Land	⁽⁵⁾ NE	NE	NE							NE	NE	NE	
G. Other	⁽⁵⁾ -533,24	NA,NE,NO	1,28							NE	NE	NE	NE
6. Waste	0,03	7,60	0,02							0,00	0,00	0,00	0,00
A. Solid Waste Disposal on Land	⁽⁶⁾ NE,NO	7,33								NE	NE	NE	
B. Waste-water Handling		0,27	0,02							NA,NE	NA,NE	NA,NE	
C. Waste Incineration	⁽⁶⁾ 0,03	NE	0,00							0,00	0,00	0,00	0,00
D. Other		NA	NA							NA	NA	NA	NA
7. Other (please specify)⁽⁷⁾	123,38	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	32,38
Geothermal Energy	123,38	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	32,38

Note: All footnotes for this table are given at the end of the table on sheet 3.

National Inventory Report 2007 - ICELAND

SUMMARY 1.A SUMMARY REPORT FOR NATIONAL GREENHOUSE GAS INVENTORIES (IPCC TABLE 7A) (Sheet 3 of 3)

Inventory 2007
Submission 2007 v1.0
ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	Net CO ₂	CH ₄	N ₂ O	HFCs		PFCs		SF ₆		NO _x	CO	NM VOC	SO ₂
	emissions/removals			P	A	P	A	P	A				
	(Gg)			CO ₂ equivalent (Gg)				(Gg)					
Memo Items: ⁽⁸⁾													
International Bunkers	407,16	0,00	0,01										
Aviation	407,16	0,00	0,01										
Marine	NA,NE,NO	NA,NE,NO	NA,NE,NO										
Multilateral Operations	NO	NO	NO							NO	NO	NO	NO
CO₂ Emissions from Biomass	NA,NO												

⁽¹⁾ The emissions of HFCs and PFCs are to be expressed as CO₂ equivalent emissions. Data on disaggregated emissions of HFCs and PFCs are to be provided in Table 2(II) of this common reporting format.

⁽²⁾ For verification purposes, countries are asked to report the results of their calculations using the Reference approach and to explain any differences with the Sectoral approach in the documentation box to Table 1.A.(c). For estimating national total emissions, the results from the Sectoral approach should be used, where possible.

⁽³⁾ Other Production includes Pulp and Paper and Food and Drink Production.

⁽⁴⁾ Parties which previously reported CO₂ from soils in the Agriculture sector should note this in the NIR.

⁽⁵⁾ For the purposes of reporting, the signs for removals are always negative (-) and for emissions positive (+).

⁽⁶⁾ CO₂ from source categories Solid Waste Disposal on Land and Waste Incineration should only be included if it stems from non-biogenic or inorganic waste streams. Only emissions from Waste Incineration Without Energy Recovery are to be reported in the Waste sector, whereas emissions from Incineration With Energy Recovery are to be reported in the Energy sector.

⁽⁷⁾ If reporting any country-specific source category under sector "7. Other", detailed explanations should be provided in Chapter 9: Other (CRF sector 7) of the NIR.

⁽⁸⁾ Countries are asked to report emissions from international aviation and marine bunkers and multilateral operations, as well as CO₂ emissions from biomass, under Memo Items. These emissions should not be included in the national total emissions from the energy sector. Amounts of biomass used as fuel are included in the national energy consumption but the corresponding CO₂ emissions are not included in the national total as it is assumed that the biomass is produced in a sustainable manner. If the biomass is harvested at an unsustainable rate, net CO₂ emissions are accounted for as a loss of biomass stocks in the Land Use, Land-use Change and Forestry sector.

National Inventory Report 2007 - ICELAND

SUMMARY 1.B SHORT SUMMARY REPORT FOR NATIONAL GREENHOUSE GAS INVENTORIES (IPCC TABLE 7B) (Sheet 1 of 1)

Inventory 2005
Submission 2007 v1.1
ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	Net CO ₂	CH ₄	N ₂ O	HFCs ⁽¹⁾		PFCs ⁽¹⁾		SF ₆		NO _x	CO	NM VOC	SO ₂
	emissions/removals			P	A	P	A	P	A				
	(Gg)			CO ₂ equivalent (Gg)				(Gg)					
Total National Emissions and Removals	4,158,24	22,15	2,35	76,74	NA,NE,NO	NA,NE,NO	26,09	0,00	NA,NE,NO	27,83	19,58	8,29	40,53
1. Energy	1,913,62	0,16	0,24							26,65	19,35	4,06	2,48
A. Fuel Combustion	Reference Approach ⁽²⁾	1,880,77											
	Sectoral Approach ⁽²⁾	1,913,62	0,16	0,24						26,65	19,35	4,06	2,48
B. Fugitive Emissions from Fuels		NA,NE,NO	NA,NE,NO	NA,NO						NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
2. Industrial Processes	835,11	0,05	NA,NE,NO	76,74	NA,NE,NO	NA,NE,NO	26,09	0,00	NA,NE,NO	1,18	0,23	0,10	5,68
3. Solvent and Other Product Use	NE		0,01							NE	NE	4,14	NE
4. Agriculture⁽³⁾		12,00	0,72							NA,NO	NA,NO	NA,NE,NO	NO
5. Land Use, Land-Use Change and Forestry⁽⁴⁾	1,286,10	2,34	1,35							NA,NE	NA,NE	NA,NE	NE
6. Waste	0,03	7,60	0,02							0,00	0,00	0,00	0,00
7. Other	123,38	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	32,38
Memo Items:⁽⁵⁾													
International Bunkers	407,16	0,00	0,01										
Aviation	407,16	0,00	0,01										
Marine	NA,NE,NO	NA,NE,NO	NA,NE,NO										
Multilateral Operations	NO	NO	NO							NO	NO	NO	NO
CO₂ Emissions from Biomass	NA,NO												

Note: A = Actual emissions based on Tier 2 approach of the IPCC Guidelines.
P = Potential emissions based on Tier 1 approach of the IPCC Guidelines.

⁽¹⁾ The emissions of HFCs and PFCs are to be expressed as CO₂ equivalent emissions. Data on disaggregated emissions of HFCs and PFCs are to be provided in Table 2(II) of this common reporting format.

⁽²⁾ For verification purposes, countries are asked to report the results of their calculations using the Reference approach and to explain any differences with the Sectoral approach in the documentation box to Table 1.A.(c). For estimating national total emissions, the result from the Sectoral approach should be used, where possible.

⁽³⁾ Parties which previously reported CO₂ from soils in the Agriculture sector should note this in the NIR.

⁽⁴⁾ For the purposes of reporting, the signs for removals are always negative (-) and for emissions positive (+).

⁽⁵⁾ Countries are asked to report emissions from international aviation and marine bunkers and multilateral operations, as well as CO₂ emissions from biomass, under Memo Items. These emissions should not be included in the national total emissions from the energy sector. Amounts of biomass used as fuel are included in the national energy consumption but the corresponding CO₂ emissions are not included in the national total as it is assumed that the biomass is produced in a sustainable manner. If the biomass is harvested at an unsustainable rate, net CO₂ emissions are accounted for as a loss of biomass stocks in the Land Use, Land-use Change and Forestry sector.

SUMMARY 2 SUMMARY REPORT FOR CO₂ EQUIVALENT EMISSIONS
(Sheet 1 of 1)

Inventory 2005
Submission 2007 v1.1
ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	CO ₂ ⁽¹⁾	CH ₄	N ₂ O	HFCs ⁽²⁾	PFCs ⁽²⁾	SF ₆ ⁽²⁾	Total
	CO ₂ equivalent (Gg)						
Total (Net Emissions)⁽¹⁾	4,158,24	465,07	728,47	76,74	26,09	5,38	5,459,98
1. Energy	1,913,62	3,38	75,22				1,992,22
A. Fuel Combustion (Sectoral Approach)	1,913,62	3,38	75,22				1,992,22
1. Energy Industries	22,60	0,01	0,56				23,17
2. Manufacturing Industries and Construction	442,28	0,37	32,73				475,38
3. Transport	716,38	1,58	35,95				753,91
4. Other Sectors	716,26	1,41	5,94				723,61
5. Other	16,11	0,01	0,04				16,15
B. Fugitive Emissions from Fuels	NA,NE,NO	NA,NE,NO	NA,NO				NA,NE,NO
1. Solid Fuels	NA,NO	NA,NO	NA,NO				NA,NO
2. Oil and Natural Gas	NA,NE,NO	NA,NE,NO	NA,NO				NA,NE,NO
2. Industrial Processes	835,11	0,98	NA,NE,NO	76,74	26,09	5,38	944,30
A. Mineral Products	55,10	NA,NE	NA,NE				55,10
B. Chemical Industry		NE,NO	NE,NO	NA	NA	NA	NA,NE,NO
C. Metal Production	780,00	0,98	NA	NA	26,09	NA,NO	807,08
D. Other Production	NE						NE
E. Production of Halocarbons and SF ₆				NA,NO	NA,NO	NA,NO	NA,NO
F. Consumption of Halocarbons and SF ₆ ⁽²⁾				76,74	NA,NE,NO	5,38	82,12
G. Other	NA	NA	NA	NA	NA	NA	NA
3. Solvent and Other Product Use	NE		3,29				3,29
4. Agriculture		251,94	223,28				475,21
A. Enteric Fermentation		231,49					231,49
B. Manure Management		20,44	25,72				46,16
C. Rice Cultivation		NA,NO					NA,NO
D. Agricultural Soils ⁽³⁾		NA,NE	197,56				197,56
E. Prescribed Burning of Savannas		NA	NA				NA
F. Field Burning of Agricultural Residues		NA,NO	NA,NO				NA,NO
G. Other		NA	NA				NA
5. Land Use, Land-Use Change and Forestry⁽¹⁾	1,286,10	49,14	419,27				1,754,51
A. Forest Land	-126,27	NA,NE,NO	1,14				-125,13
B. Cropland	3,46	NE,NO	NE,NO				3,46
C. Grassland	1,800,74	NE	NE				1,800,74
D. Wetlands	141,42	49,14	21,70				212,26
E. Settlements	NE	NE	NE				NE
F. Other Land	NE	NE	NE				NE
G. Other	-533,24	NA,NE,NO	396,43				-136,81
6. Waste	0,03	159,63	7,41				167,06
A. Solid Waste Disposal on Land	NE,NO	154,00					154,00
B. Waste-water Handling		5,63	7,41				13,04
C. Waste Incineration	0,03	NE	0,00				0,03
D. Other	NA	NA	NA				NA
7. Other (as specified in Summary 1.A)	123,38	NA	NA	NA	NA	NA	123,38
Memo Items:⁽⁴⁾							
International Bunkers	407,16	0,06	3,57				410,79
Aviation	407,16	0,06	3,57				410,79
Marine	NA,NE,NO	NA,NE,NO	NA,NE,NO				NA,NE,NO
Multilateral Operations	NO	NO	NO				NO
CO₂ Emissions from Biomass	NA,NO						NA,NO
Total CO ₂ Equivalent Emissions without Land Use, Land-Use Change and Forestry							3,705,47
Total CO ₂ Equivalent Emissions with Land Use, Land-Use Change and Forestry							5,459,98

⁽¹⁾ For CO₂ from Land Use, Land-use Change and Forestry the net emissions/removals are to be reported. For the purposes of reporting, the signs for removals are always negative (-) and for emissions positive (+).

⁽²⁾ Actual emissions should be included in the national totals. If no actual emissions were reported, potential emissions should be included.

⁽³⁾ Parties which previously reported CO₂ from soils in the Agriculture sector should note this in the NIR.

⁽⁴⁾ See footnote 8 to table Summary 1.A.

National Inventory Report 2007 - ICELAND

SUMMARY 3 SUMMARY REPORT FOR METHODS AND EMISSION FACTORS USED (Sheet 1 of 2)

Inventory 2005
Submission 2007 v1.1
ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	CO ₂		CH ₄		N ₂ O		HFCs		PFCs		SF ₆	
	Method applied	Emission factor	Method applied	Emission factor	Method applied	Emission factor	Method applied	Emission factor	Method applied	Emission factor	Method applied	Emission factor
1. Energy												
A. Fuel Combustion												
1. Energy Industries												
2. Manufacturing Industries and Construction												
3. Transport												
4. Other Sectors												
5. Other												
B. Fugitive Emissions from Fuels	NA	NA	NA	NA	NA	NA						
1. Solid Fuels	NA	NA	NA	NA	NA	NA						
2. Oil and Natural Gas	NA	NA	NA	NA	NA	NA						
2. Industrial Processes							NA	NA			NA	NA
A. Mineral Products			NA	NA	NA	NA						
B. Chemical Industry			NA	NA	NA	NA					NA	NA
C. Metal Production					NA	NA						
D. Other Production												
E. Production of Halocarbons and SF ₆							NA	NA	NA	NA	NA	NA
F. Consumption of Halocarbons and SF ₆							NA	NA	NA	NA	NA	NA
G. Other	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

Use the following notation keys to specify the method applied:

D (IPCC default)	T1a, T1b, T1c (IPCC Tier 1a, Tier 1b and Tier 1c, respectively)	CR (CORINAIR)
RA (Reference Approach)	T2 (IPCC Tier 2)	CS (Country Specific)
T1 (IPCC Tier 1)	T3 (IPCC Tier 3)	OTH (Other)

If using more than one method within one source category, list all the relevant methods. Explanations regarding country-specific methods, other methods or any modifications to the default IPCC methods, as well as information

Use the following notation keys to specify the emission factor used:

D (IPCC default)	CS (Country Specific)	OTH (Other)
CR (CORINAIR)	PS (Plant Specific)	

Where a mix of emission factors has been used, list all the methods in the relevant cells and give further explanations in the documentation box. Also use the documentation box to explain the use of notation OTH.

National Inventory Report 2007 - ICELAND

SUMMARY 3 SUMMARY REPORT FOR METHODS AND EMISSION FACTORS USED (Sheet 2 of 2)

Inventory 2005
Submission 2007 v1.1
ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	CO ₂		CH ₄		N ₂ O		HFCs		PFCs		SF ₆	
	Method applied	Emission factor	Method applied	Emission factor	Method applied	Emission factor	Method applied	Emission factor	Method applied	Emission factor	Method applied	Emission factor
3. Solvent and Other Product Use	NA	NA										
4. Agriculture												
A. Enteric Fermentation												
B. Manure Management												
C. Rice Cultivation			NA	NA								
D. Agricultural Soils			NA	NA								
E. Prescribed Burning of Savannas			NA	NA	NA	NA						
F. Field Burning of Agricultural Residues			NA	NA	NA	NA						
G. Other			NA	NA	NA	NA						
5. Land Use, Land-Use Change and Forestry	CS,T1	CS,D			T1	D						
A. Forest Land			NA	NA	T1	D						
B. Cropland	CS,T1	CS,D	NA	NA	NA	NA						
C. Grassland			NA	NA	NA	NA						
D. Wetlands												
E. Settlements	NA	NA	NA	NA	NA	NA						
F. Other Land	NA	NA	NA	NA	NA	NA						
G. Other			NA	NA	T1	D						
6. Waste												
A. Solid Waste Disposal on Land	NA	NA										
B. Waste-water Handling												
C. Waste Incineration			NA	NA								
D. Other	NA	NA	NA	NA	NA	NA						
7. Other (as specified in Summary 1.A)	CS		NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

Use the following notation keys to specify the method applied:

D (IPCC default)	T1a, T1b, T1c (IPCC Tier 1a, Tier 1b and Tier 1c, respectively)	CR (CORINAIR)
RA (Reference Approach)	T2 (IPCC Tier 2)	CS (Country Specific)
T1 (IPCC Tier 1)	T3 (IPCC Tier 3)	OTH (Other)

If using more than one method within one source category, list all the relevant methods. Explanations regarding country-specific methods, other methods or any modifications to the default IPCC methods, as well as information regarding the use of different methods per

Use the following notation keys to specify the emission factor used:

D (IPCC default)	CS (Country Specific)	OTH (Other)
CR (CORINAIR)	PS (Plant Specific)	

Where a mix of emission factors has been used, list all the methods in the relevant cells and give further explanations in the documentation box. Also use the documentation box to explain the use of notation OTH.

Documentation box:

- Parties should provide the full information on methodological issues, such as methods and emission factors used, in the relevant sections of Chapters 3 to 9 (see section 2.2 of each of Chapters 3 - 9) of the NIR. Use this documentation box to provide references to relevant sections of the NIR if any additional information and further details are needed to understand the content of this table.
- Where a mix of methods/emission factors has been used within one source category, use this documentation box to specify those methods/emission factors for the various sub-sources where they have been applied.
- Where the notation OTH (Other) has been entered in this table, use this documentation box to specify those other methods/emission factors.

TABLE 7 SUMMARY OVERVIEW FOR KEY CATEGORIES
(Sheet 1 of 1)

KEY CATEGORIES OF EMISSIONS AND REMOVALS	Gas	Criteria used for key source identification			Key category excluding LULUCF ⁽¹⁾	Key category including LULUCF ⁽¹⁾	Comments ⁽²⁾
		L	T	Q			
Specify key categories according to the national level of disaggregation used:							

Note: L = Level assessment; T = Trend assessment; Q = Qualitative assessment.

⁽¹⁾ The term "key categories" refers to both the key source categories as addressed in the IPCC good practice guidance and the key categories as addressed in the IPCC good practice guidance for LULUCF.

⁽²⁾ For estimating key categories Parties may chose the disaggregation level presented as an example in table 7.1 of the IPCC good practice guidance (page 7.6) and table 5.4.1 (page 5.31) of the IPCC good practice guidance for LULUCF, the level used in table Summary 1.A of the common reporting format or any other disaggregation level that the Party used to determine its key categories.

Documentation box:

Parties should provide the full information on methodologies used for identifying key categories and the quantitative results from the level and trend assessments (according to tables 7.1–7.3 of the IPCC good practice guidance and tables 5.4.1–5.4.3 of the IPCC good practice guidance for LULUCF) in Annex 1 to the NIR.

National Inventory Report 2007 - ICELAND

TABLE 8(a) RECALCULATION - RECALCULATED DATA
(Sheet 1 of 2)

Recalculated year: Inventory 2005
Submission 2007 v1.1
ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	CO ₂						CH ₄						N ₂ O					
	Previous submission	Latest submission	Difference	Difference ⁽¹⁾	Impact of recalculation on total emissions excluding LULUCF ⁽²⁾	Impact of recalculation on total emissions including LULUCF ⁽²⁾	Previous submission	Latest submission	Difference	Difference ⁽¹⁾	Impact of recalculation on total emissions excluding LULUCF ⁽²⁾	Impact of recalculation on total emissions including LULUCF ⁽²⁾	Previous submission	Latest submission	Difference	Difference ⁽¹⁾	Impact of recalculation on total emissions excluding LULUCF ⁽²⁾	Impact of recalculation on total emissions including LULUCF ⁽²⁾
	CO ₂ equivalent (Gg)						CO ₂ equivalent (Gg)						CO ₂ equivalent (Gg)					
	%						%						%					
Total National Emissions and Removals		4,158.24					465.07						728.47					
1. Energy		1,913.62					3.28						75.22					
1.A. Fuel Combustion Activities		1,913.62					3.28						75.22					
1.A.1. Energy Industries		22.60					0.01						0.56					
1.A.2. Manufacturing Industries and Construction		442.28					0.37						32.73					
1.A.3. Transport		716.38					1.58						35.95					
1.A.4. Other Sectors		716.26					1.41						5.94					
1.A.5. Other		16.11					0.01						0.04					
1.B. Fugitive Emissions from Fuels		NA,NE,NO					NA,NE,NO						NA,NO					
1.B.1. Solid fuel		NA,NO					NA,NO						NA,NO					
1.B.2. Oil and Natural Gas		NA,NE,NO					NA,NE,NO						NA,NO					
2. Industrial Processes		835.11					0.98						NA,NE,NO					
2.A. Mineral Products		55.10					NA,NE						NA,NE					
2.B. Chemical Industry							NE,NO						NE,NO					
2.C. Metal Production		780.00					0.98						NA					
2.D. Other Production		NE																
2.G. Other		NA					NA						NA					
3. Solvent and Other Product Use		NE											3.29					
4. Agriculture							251.94						223.28					
4.A. Enteric Fermentation							231.49											
4.B. Manure Management							20.44						25.72					
4.C. Rice Cultivation							NA,NO											
4.D. Agricultural Soils ⁽⁴⁾							NA,NE						197.56					
4.E. Prescribed Burning of Savannas							NA						NA					
4.F. Field Burning of Agricultural Residues							NA,NO						NA,NO					
4.G. Other							NA						NA					
5. Land Use, Land-Use Change and Forestry (net)⁽⁵⁾		1,286.10					49.14						419.27					
5.A. Forest Land		-126.27					NA,NE,NO						1.14					
5.B. Cropland		3.46					NE,NO						NE,NO					
5.C. Grassland		1,800.74					NE						NE					
5.D. Wetlands		141.42					49.14						21.70					
5.E. Settlements		NE					NE						NE					
5.F. Other Land		NE					NE						NE					
5.G. Other		-533.24					NA,NE,NO						396.43					

Note: All footnotes for this table are given at the end of the table on sheet 2.

National Inventory Report 2007 - ICELAND

TABLE 8(a) RECALCULATION - RECALCULATED DATA
(Sheet 2 of 2)

Recalculated year: Inventory 2005
Submission 2007 v1.1
ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	CO ₂						CH ₄						N ₂ O					
	Previous submission	Latest submission	Difference	Difference ⁽¹⁾	Impact of recalculation on total emissions excluding LULUCF ⁽²⁾	Impact of recalculation on total emissions including LULUCF ⁽³⁾	Previous submission	Latest submission	Difference	Difference ⁽¹⁾	Impact of recalculation on total emissions excluding LULUCF ⁽²⁾	Impact of recalculation on total emissions including LULUCF ⁽³⁾	Previous submission	Latest submission	Difference	Difference ⁽¹⁾	Impact of recalculation on total emissions excluding LULUCF ⁽²⁾	Impact of recalculation on total emissions including LULUCF ⁽³⁾
	CO ₂ equivalent (Gg)						CO ₂ equivalent (Gg)						CO ₂ equivalent (Gg)					
6. Waste		0,03					159,63							7,41				
6.A. Solid Waste Disposal on Land		NE,NO					154,00											
6.B. Waste-water Handling							5,63							7,41				
6.C. Waste Incineration		0,03					NE							0,00				
6.D. Other		NA					NA							NA				
7. Other (as specified in Summary I.A)		123,38					NA							NA				
Memo Items:																		
International Bankers		407,16					0,06							3,57				
Multilateral Operations		NO					NO							NO				
CO ₂ Emissions from Biomass		NA,NO																

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	HFCs						PFCs						SF ₆					
	Previous submission	Latest submission	Difference	Difference ⁽¹⁾	Impact of recalculation on total emissions excluding LULUCF ⁽²⁾	Impact of recalculation on total emissions including LULUCF ⁽³⁾	Previous submission	Latest submission	Difference	Difference ⁽¹⁾	Impact of recalculation on total emissions excluding LULUCF ⁽²⁾	Impact of recalculation on total emissions including LULUCF ⁽³⁾	Previous submission	Latest submission	Difference	Difference ⁽¹⁾	Impact of recalculation on total emissions excluding LULUCF ⁽²⁾	Impact of recalculation on total emissions including LULUCF ⁽³⁾
	CO ₂ equivalent (Gg)						CO ₂ equivalent (Gg)						CO ₂ equivalent (Gg)					
Total Actual Emissions		NA,NE,NO					26,09							NA,NE,NO				
2.C.3. Aluminium Production							26,09											
2.E. Production of Halocarbons and SF ₆		NA,NO					NA,NO							NA,NO				
2.F. Consumption of Halocarbons and SF ₆		NA,NE,NO					NA,NE,NO							NA,NE,NO				
2.G. Other		NA					NA							NA				
Potential Emissions from Consumption of HFCs/PFCs and SF ₆		76,74					NE,NO							5,38				

	Previous submission	Latest submission	Difference	Difference ⁽¹⁾
	CO ₂ equivalent (Gg)			
Total CO ₂ Equivalent Emissions with Land Use, Land-Use Change and Forestry		5.459,98		
Total CO ₂ Equivalent Emissions without Land Use, Land-Use Change and Forestry		3.623,35		

⁽¹⁾ Estimate the percentage change due to recalculation with respect to the previous submission (percentage change = 100 x [(LS-PS)/PS], where LS = latest submission and PS = previous submission. All cases of recalculation of the estimate of the source/sink category should be addressed and explained in table 8(b).
⁽²⁾ Total emissions refer to total aggregate GHG emissions expressed in terms of CO₂ equivalent, excluding GHGs from the LULUCF sector. The impact of the recalculation on the total emissions is calculated as follows: impact of recalculation (%) = 100 x [(source (LS) - source (PS))/total emissions (LS)], where LS = latest submission, PS = previous submission.
⁽³⁾ Total emissions refer to total aggregate GHG emissions expressed in terms of CO₂ equivalent, including GHGs from the LULUCF sector. The impact of the recalculation on the total emissions is calculated as follows: impact of recalculation (%) = 100 x [(source (LS) - source (PS))/total emissions (LS)], where LS = latest submission, PS = previous submission.
⁽⁴⁾ Parties which previously reported CO₂ from soils in the Agriculture sector should note this in the NIR.
⁽⁵⁾ Net CO₂ emissions/removals to be reported.

Documentation box:
Parties should provide detailed information on recalculations in Chapter 10: Recalculations and Improvements, and in the relevant sections of Chapters 3 to 9 (see section 2.5 of each of Chapters 3 - 9) of the NIR. Use this documentation box to provide references to relevant sections of the NIR if any additional information and further details are needed to understand the content of this table.

TABLE 8(b) RECALCULATION - EXPLANATORY INFORMATION
(Sheet 1 of 1)

Specify the sector and source/sink category ⁽¹⁾ where changes in estimates have occurred:	GHG	RECALCULATION DUE TO				
		CHANGES IN:			Addition/removal/ reallocation of source/sink categories	Other changes in data (e.g. statistical or editorial changes, correction of errors)
		Methods ⁽²⁾	Emission factors ⁽²⁾	Activity data ⁽²⁾		

⁽¹⁾ Enter the identification code of the source/sink category (e.g. 1.B.1) in the first column and the name of the category (e.g. Fugitive Emissions from Solid Fuels) in the second column of the table. Note that the source categories entered in this table should match those used in table 8(a).
⁽²⁾ Explain changes in methods, emission factors and activity data that have resulted in recalculation of the estimate of the source/sink as indicated in table 8(a). Include changes in the assumptions and coefficients in the Methods column.

Documentation box:
Parties should provide the full information on recalculations in Chapter 10: Recalculations and Improvements, and in the relevant sections of Chapters 3 to 9 (see section 2.5 of each of Chapters 3 to 9) of the NIR. Use this documentation box to provide references to relevant sections of the NIR if any additional information and further details are needed to understand the content of this table. References should point particularly to the sections of the NIR in which justifications of the changes as to improvements in the accuracy, completeness and consistency of the inventory are reported.

National Inventory Report 2007 - ICELAND

TABLE 9(b) COMPLETENESS - INFORMATION ON ADDITIONAL GREENHOUSE GASES
(Sheet 1 of 1)

Inventory 2005
 Submission 2007 v1.1
 ICELAND

Additional GHG emissions reported ⁽¹⁾						
GHG	Source category	Emissions (Gg)	Estimated GWP value (100-year horizon)	Emissions CO ₂ equivalent (Gg)	Reference to the source of GWP value	Explanation

⁽¹⁾ Parties are encouraged to provide information on emissions of greenhouse gases whose GWP values have not yet been agreed upon by the COP. Include such gases in this table if they are considered in the submitted inventory. Provide additional information on the estimation methods used.

Documentation box:
 Parties should provide detailed information regarding completeness of the inventory in the NIR (Chapter 1.8: General Assessment of the Completeness, and Annex 5). Use this documentation box to provide references to relevant sections of the NIR if any additional information and further details are needed to understand the content of this table.

National Inventory Report 2007 - ICELAND

TABLE 10 EMISSION TRENDS
CO₂
(Part 1 of 2)

Inventory 2005
Submission 2007 v1.1
ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	Base year (1990)	1991	1992	1993	1994	1995	1996	1997	1998	1999
	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)
1. Energy	1.672,56	1.627,13	1.752,38	1.812,36	1.774,87	1.777,72	1.868,40	1.915,47	1.877,33	1.906,72
A. Fuel Combustion (Sectoral Approach)	1.672,56	1.627,13	1.752,38	1.812,36	1.774,87	1.777,72	1.868,40	1.915,47	1.877,33	1.906,72
1. Energy Industries	20,70	22,28	21,29	22,35	22,22	24,61	20,00	15,27	37,64	20,64
2. Manufacturing Industries and Construction	361,05	285,42	338,63	365,40	344,24	358,05	398,89	467,81	441,43	466,69
3. Transport	600,13	611,43	621,54	622,17	624,79	600,44	590,81	602,47	605,24	626,84
4. Other Sectors	690,56	707,87	770,13	801,03	783,53	793,00	858,33	829,89	788,06	788,18
5. Other	0,12	0,14	0,78	1,42	0,10	1,62	0,38	0,03	4,95	4,36
B. Fugitive Emissions from Fuels	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
1. Solid Fuels	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO
2. Oil and Natural Gas	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
2. Industrial Processes	392,66	359,37	362,43	409,86	410,71	427,14	426,21	484,91	512,73	659,07
A. Mineral Products	52,34	48,71	45,74	39,73	37,45	37,96	41,87	46,64	54,49	61,52
B. Chemical Industry	0,36	0,31	0,25	0,24	0,35	0,46	0,40	0,44	0,40	0,43
C. Metal Production	339,96	310,34	316,43	369,89	372,91	388,72	383,94	437,83	457,84	597,12
D. Other Production	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
E. Production of Halocarbons and SF ₆										
F. Consumption of Halocarbons and SF ₆										
G. Other	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
3. Solvent and Other Product Use	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
4. Agriculture										
A. Enteric Fermentation										
B. Manure Management										
C. Rice Cultivation										
D. Agricultural Soils										
E. Prescribed Burning of Savannas										
F. Field Burning of Agricultural Residues										
G. Other										
5. Land Use, Land-Use Change and Forestry⁽²⁾	1.622,57	1.606,92	1.587,62	1.572,71	1.554,32	1.532,82	1.513,92	1.493,34	1.467,40	1.438,74
A. Forest Land	-33,30	-39,35	-45,84	-52,32	-58,38	-64,00	-69,19	-73,95	-79,57	-86,05
B. Cropland	IE,NA,NE,NO	IE,NA,NE,NO	IE,NA,NE,NO	IE,NA,NE,NO	IE,NA,NE,NO	IE,NA,NE,NO	IE,NA,NE,NO	IE,NA,NE,NO	IE,NA,NE,NO	IE,NA,NE,NO
C. Grassland	1.811,55	1.811,00	1.810,27	1.809,50	1.808,69	1.807,96	1.807,30	1.806,68	1.806,09	1.805,43
D. Wetlands	141,42	141,42	141,42	141,42	141,42	141,42	141,42	141,42	141,42	141,42
E. Settlements	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
F. Other Land	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
G. Other	-297,11	-306,16	-318,23	-325,89	-337,42	-352,57	-365,61	-380,81	-400,55	-422,06
6. Waste	18,84	18,69	18,19	15,49	14,27	12,59	11,28	10,87	9,21	7,53
A. Solid Waste Disposal on Land	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
B. Waste-water Handling										
C. Waste Incineration	18,84	18,69	18,19	15,49	14,27	12,59	11,28	10,87	9,21	7,53
D. Other	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
7. Other (as specified in Summary 1.A)	66,63	66,63	66,63	66,63	66,63	81,68	82,18	71,38	94,08	122,96
Total CO₂ emissions including net CO₂ from LULUCF	3.773,27	3.678,74	3.787,24	3.877,05	3.820,81	3.831,95	3.902,00	3.975,97	3.960,75	4.135,02
Total CO₂ emissions excluding net CO₂ from LULUCF	2.150,70	2.071,82	2.199,62	2.304,35	2.266,49	2.299,13	2.388,07	2.482,63	2.493,35	2.696,28
Memo Items:										
International Bunkers	318,65	259,64	263,56	293,02	307,10	380,15	395,45	440,80	514,67	527,25
Aviation	219,65	221,99	203,62	195,64	213,62	236,15	271,51	292,12	338,13	363,37
Marine	99,00	37,65	59,95	97,38	93,49	144,00	123,95	148,68	176,54	163,88
Multilateral Operations	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
CO₂ Emissions from Biomass	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO

Note: All footnotes for this table are given at the end of the table on sheet 5.

National Inventory Report 2007 - ICELAND

TABLE 10 EMISSION TRENDS
CO₂
(Part 2 of 2)

Inventory 2005
Submission 2007 v1.1
ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	2000	2001	2002	2003	2004	2005	Change from base to latest reported year
	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)	%
1. Energy	1,808.95	1,781.97	1,854.70	1,797.28	1,890.14	1,913.62	14.41
A. Fuel Combustion (Sectoral Approach)	1,808.95	1,781.97	1,854.70	1,797.28	1,890.14	1,913.62	14.41
1. Energy Industries	14.41	14.54	15.69	14.63	19.85	22.60	9.17
2. Manufacturing Industries and Construction	419.46	451.59	452.83	424.87	450.59	442.28	22.50
3. Transport	629.42	640.06	643.65	666.71	678.56	716.38	19.37
4. Other Sectors	741.05	656.26	720.24	675.62	725.28	716.26	3.72
5. Other	4.61	19.53	22.30	15.45	15.85	16.11	13,315.75
B. Fugitive Emissions from Fuels	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	0.00
1. Solid Fuels	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	0.00
2. Oil and Natural Gas	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	0.00
2. Industrial Processes	765.57	803.55	822.27	824.33	846.01	835.11	112.68
A. Mineral Products	65.57	58.77	40.56	33.08	50.93	55.10	5.27
B. Chemical Industry	0.41	0.49	0.45	0.48	0.39		-100.00
C. Metal Production	699.60	744.28	781.25	790.78	794.69	780.00	129.44
D. Other Production	NE	NE	NE	NE	NE	NE	0.00
E. Production of Halocarbons and SF ₆							
F. Consumption of Halocarbons and SF ₆							
G. Other	NA	NA	NA	NA	NA	NA	0.00
3. Solvent and Other Product Use	NE	NE	NE	NE	NE	NE	0.00
4. Agriculture							
A. Enteric Fermentation							
B. Manure Management							
C. Rice Cultivation							
D. Agricultural Soils							
E. Prescribed Burning of Savannas							
F. Field Burning of Agricultural Residues							
G. Other							
5. Land Use, Land-Use Change and Forestry⁽²⁾	1,413.75	1,396.31	1,375.51	1,350.97	1,315.80	1,286.10	-20.74
A. Forest Land	-92.11	-97.73	-104.65	-111.14	-118.49	-126.27	279.22
B. Cropland	IE,NA,NE,NO	IE,NA,NE,NO	IE,NA,NE,NO	2.36	2.52	3.46	100.00
C. Grassland	1,804.66	1,803.93	1,803.23	1,802.42	1,801.62	1,800.74	-0.60
D. Wetlands	141.42	141.42	141.42	141.42	141.42	141.42	0.00
E. Settlements	NE	NE	NE	NE	NE	NE	0.00
F. Other Land	NE	NE	NE	NE	NE	NE	0.00
G. Other	-440.23	-451.32	-464.49	-484.11	-511.27	-533.24	79.48
6. Waste	7.08	6.57	6.10	5.20	2.44	0.03	-99.86
A. Solid Waste Disposal on Land	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NE,NO	0.00
B. Waste-water Handling							
C. Waste Incineration	7.08	6.57	6.10	5.20	2.44	0.03	-99.86
D. Other	NA	NA	NA	NA	NA	NA	0.00
7. Other (as specified in Summary 1.A)	163.48	154.48	159.35	137.89	124.08	123.38	85.17
Total CO₂ emissions including net CO₂ from LULUCF	4,158.84	4,142.87	4,217.93	4,115.67	4,178.48	4,158.24	10.20
Total CO₂ emissions excluding net CO₂ from LULUCF	2,745.09	2,746.56	2,842.41	2,764.70	2,862.68	2,872.13	33.54
Memo Items:							
International Bunkers	626.29	498.17	517.17	509.59	597.30	407.16	27.78
Aviation	407.74	349.13	309.85	330.02	370.26	407.16	85.37
Marine	218.55	149.04	207.32	179.57	227.04	NA,NE,NO	-100.00
Multilateral Operations	NO	NO	NO	NO	NO	NO	0.00
CO₂ Emissions from Biomass	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	0.00

Note: All footnotes for this table are given at the end of the table on sheet 5.

National Inventory Report 2007 - ICELAND

TABLE 10 EMISSION TRENDS
CH₄
(Part 1 of 2)

Inventory 2005
Submission 2007 v1.1
ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	Base year (1990)	1991	1992	1993	1994	1995	1996	1997	1998	1999
	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)
1. Energy	0,22	0,23	0,24	0,24	0,24	0,22	0,23	0,20	0,20	0,17
A. Fuel Combustion (Sectoral Approach)	0,22	0,23	0,24	0,24	0,24	0,22	0,23	0,20	0,20	0,17
1. Energy Industries	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
2. Manufacturing Industries and Construction	0,01	0,01	0,01	0,01	0,01	0,01	0,01	0,02	0,02	0,02
3. Transport	0,15	0,15	0,16	0,16	0,16	0,13	0,13	0,11	0,11	0,08
4. Other Sectors	0,06	0,06	0,07	0,07	0,07	0,07	0,08	0,08	0,07	0,07
5. Other	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
B. Fugitive Emissions from Fuels	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
1. Solid Fuels	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO
2. Oil and Natural Gas	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
2. Industrial Processes	0,03	0,02	0,02	0,03	0,03	0,03	0,03	0,03	0,02	0,03
A. Mineral Products	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE
B. Chemical Industry	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO
C. Metal Production	0,03	0,02	0,02	0,03	0,03	0,03	0,03	0,03	0,02	0,03
D. Other Production										
E. Production of Halocarbons and SF ₆										
F. Consumption of Halocarbons and SF ₆										
G. Other	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
3. Solvent and Other Product Use										
4. Agriculture	13,97	13,67	13,26	13,18	13,25	12,85	12,96	13,12	13,28	13,16
A. Enteric Fermentation	12,85	12,57	12,18	12,11	12,19	11,79	11,90	12,06	12,20	12,09
B. Manure Management	1,11	1,10	1,08	1,07	1,07	1,06	1,06	1,07	1,08	1,07
C. Rice Cultivation	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO
D. Agricultural Soils	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE
E. Prescribed Burning of Savannas	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
F. Field Burning of Agricultural Residues	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO
G. Other	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
5. Land Use, Land-Use Change and Forestry	2,34	2,34	2,34	2,34	2,34	2,34	2,34	2,34	2,34	2,34
A. Forest Land	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
B. Cropland	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO
C. Grassland	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
D. Wetlands	2,34	2,34	2,34	2,34	2,34	2,34	2,34	2,34	2,34	2,34
E. Settlements	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
F. Other Land	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
G. Other	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
6. Waste	5,47	5,66	5,84	5,99	6,14	6,29	6,54	6,70	6,85	7,06
A. Solid Waste Disposal on Land	5,41	5,60	5,77	5,92	6,07	6,22	6,47	6,63	6,77	6,95
B. Waste-water Handling	0,06	0,06	0,06	0,06	0,07	0,07	0,07	0,07	0,09	0,11
C. Waste Incineration	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
D. Other	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
7. Other (as specified in Summary I.A)	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Total CH₄ emissions including CH₄ from LULUCF	22,03	21,93	21,70	21,78	22,00	21,73	22,10	22,40	22,70	22,76
Total CH₄ emissions excluding CH₄ from LULUCF	19,69	19,59	19,36	19,44	19,66	19,39	19,76	20,06	20,36	20,42
Memo Items:										
International Bunkers	0,01	0,01	0,01	0,01	0,01	0,02	0,01	0,02	0,02	0,02
Aviation	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Marine	0,01	0,00	0,01	0,01	0,01	0,01	0,01	0,01	0,02	0,02
Multilateral Operations	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
CO₂ Emissions from Biomass										

Note: All footnotes for this table are given at the end of the table on sheet 5.

National Inventory Report 2007 - ICELAND

TABLE 10 EMISSION TRENDS
CH₄
(Part 2 of 2)

Inventory 2005
Submission 2007 v1.1
ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	2000	2001	2002	2003	2004	2005	Change from base to latest reported year
	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)	%
1. Energy	0,17	0,16	0,17	0,17	0,17	0,16	-27,68
A. Fuel Combustion (Sectoral Approach)	0,17	0,16	0,17	0,17	0,17	0,16	-27,68
1. Energy Industries	0,00	0,00	0,00	0,00	0,00	0,00	-10,27
2. Manufacturing Industries and Construction	0,02	0,02	0,02	0,02	0,02	0,02	47,42
3. Transport	0,08	0,08	0,08	0,09	0,09	0,08	-48,72
4. Other Sectors	0,07	0,06	0,07	0,06	0,07	0,07	6,07
5. Other	0,00	0,00	0,00	0,00	0,00	0,00	13.433,66
B. Fugitive Emissions from Fuels	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	0,00
1. Solid Fuels	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	0,00
2. Oil and Natural Gas	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	0,00
2. Industrial Processes	0,04	0,04	0,05	0,04	0,05	0,05	61,01
A. Mineral Products	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE	0,00
B. Chemical Industry	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	0,00
C. Metal Production	0,04	0,04	0,05	0,04	0,05	0,05	61,01
D. Other Production							
E. Production of Halocarbons and SF ₆							
F. Consumption of Halocarbons and SF ₆							
G. Other	NA	NA	NA	NA	NA	NA	0,00
3. Solvent and Other Product Use							
4. Agriculture	12,60	12,56	12,26	12,05	11,86	12,00	-14,10
A. Enteric Fermentation	11,56	11,53	11,27	11,08	10,91	11,02	-14,24
B. Manure Management	1,04	1,03	0,99	0,97	0,95	0,97	-12,56
C. Rice Cultivation	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	0,00
D. Agricultural Soils	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE	0,00
E. Prescribed Burning of Savannas	NA	NA	NA	NA	NA	NA	0,00
F. Field Burning of Agricultural Residues	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	0,00
G. Other	NA	NA	NA	NA	NA	NA	0,00
5. Land Use, Land-Use Change and Forestry	2,34	2,34	2,34	2,34	2,34	2,34	0,00
A. Forest Land	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	0,00
B. Cropland	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	0,00
C. Grassland	NE	NE	NE	NE	NE	NE	0,00
D. Wetlands	2,34	2,34	2,34	2,34	2,34	2,34	0,00
E. Settlements	NE	NE	NE	NE	NE	NE	0,00
F. Other Land	NE	NE	NE	NE	NE	NE	0,00
G. Other	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	0,00
6. Waste	7,32	7,65	7,39	7,56	7,50	7,60	39,00
A. Solid Waste Disposal on Land	7,17	7,50	7,13	7,30	7,23	7,33	35,59
B. Waste-water Handling	0,15	0,15	0,26	0,26	0,26	0,27	343,26
C. Waste Incineration	NE	NE	NE	NE	NE	NE	0,00
D. Other	NA	NA	NA	NA	NA	NA	0,00
7. Other (as specified in Summary 1.A)	NA	NA	NA	NA	NA	NA	0,00
Total CH₄ emissions including CH₄ from LULUCF	22,46	22,76	22,20	22,16	21,92	22,15	0,54
Total CH₄ emissions excluding CH₄ from LULUCF	20,12	20,42	19,86	19,82	19,58	19,81	0,60
Memo Items:							
International Bunkers	0,02	0,02	0,02	0,02	0,02	0,00	-73,85
Aviation	0,00	0,00	0,00	0,00	0,00	0,00	85,35
Marine	0,02	0,01	0,02	0,02	0,02	NA,NE,NO	-100,00
Multilateral Operations	NO	NO	NO	NO	NO	NO	0,00
CO₂ Emissions from Biomass							

Note: All footnotes for this table are given at the end of the table on sheet 5.

National Inventory Report 2007 - ICELAND

TABLE 10 EMISSION TRENDS
N₂O
(Part 1 of 2)

Inventory 2005
Submission 2007 v1.1
ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	Base year (1990)	1991	1992	1993	1994	1995	1996	1997	1998	1999
	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)
1. Energy	0.09	0.08	0.08	0.09	0.09	0.12	0.12	0.16	0.16	0.19
A. Fuel Combustion (Sectoral Approach)	0.09	0.08	0.08	0.09	0.09	0.12	0.12	0.16	0.16	0.19
1. Energy Industries	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
2. Manufacturing Industries and Construction	0.05	0.05	0.05	0.05	0.05	0.06	0.06	0.07	0.07	0.08
3. Transport	0.02	0.02	0.02	0.02	0.02	0.04	0.04	0.06	0.06	0.09
4. Other Sectors	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
5. Other	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
B. Fugitive Emissions from Fuels	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO
1. Solid Fuels	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO
2. Oil and Natural Gas	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO
2. Industrial Processes	0.16	0.15	0.14	0.14	0.14	0.14	0.16	0.13	0.12	0.12
A. Mineral Products	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE
B. Chemical Industry	0.16	0.15	0.14	0.14	0.14	0.14	0.16	0.13	0.12	0.12
C. Metal Production	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
D. Other Production										
E. Production of Halocarbons and SF ₆										
F. Consumption of Halocarbons and SF ₆										
G. Other	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
3. Solvent and Other Product Use	0.02	0.02	0.02	0.02	0.01	0.02	0.02	0.02	0.02	0.02
4. Agriculture	0.89	0.86	0.81	0.83	0.85	0.81	0.84	0.83	0.83	0.86
A. Enteric Fermentation										
B. Manure Management	0.11	0.10	0.10	0.10	0.10	0.09	0.09	0.09	0.09	0.09
C. Rice Cultivation										
D. Agricultural Soils	0.78	0.76	0.71	0.73	0.75	0.72	0.75	0.74	0.74	0.77
E. Prescribed Burning of Savannas	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
F. Field Burning of Agricultural Residues	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO
G. Other	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
5. Land Use, Land-Use Change and Forestry	1.35	1.35	1.35	1.34	1.34	1.35	1.34	1.35	1.35	1.35
A. Forest Land	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
B. Cropland	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO
C. Grassland	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
D. Wetlands	0.07	0.07	0.07	0.07	0.07	0.07	0.07	0.07	0.07	0.07
E. Settlements	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
F. Other Land	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
G. Other	1.28	1.28	1.28	1.27	1.27	1.27	1.27	1.27	1.28	1.27
6. Waste	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
A. Solid Waste Disposal on Land										
B. Waste-water Handling	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
C. Waste Incineration	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
D. Other	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
7. Other (as specified in Summary 1.A)	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Total N₂O emissions including N₂O from LULUCF	2.52	2.49	2.42	2.44	2.46	2.45	2.51	2.50	2.50	2.56
Total N₂O emissions excluding N₂O from LULUCF	1.17	1.14	1.07	1.10	1.12	1.10	1.16	1.16	1.15	1.21
Memo Items:										
International Bunkers	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Aviation	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Marine	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Multilateral Operations	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
CO₂ Emissions from Biomass										

Note: All footnotes for this table are given at the end of the table on sheet 5.

National Inventory Report 2007 - ICELAND

TABLE 10 EMISSION TRENDS
N₂O
(Part 2 of 2)

Inventory 2005
 Submission 2007 v1.1
 ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	2000	2001	2002	2003	2004	2005	Change from base to latest reported year
	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)	%
1. Energy	0.19	0.19	0.19	0.20	0.22	0.24	181.62
A. Fuel Combustion (Sectoral Approach)	0.19	0.19	0.19	0.20	0.22	0.24	181.62
1. Energy Industries	0.00	0.00	0.00	0.00	0.00	0.00	1,284.85
2. Manufacturing Industries and Construction	0.08	0.08	0.08	0.08	0.10	0.11	105.63
3. Transport	0.09	0.09	0.09	0.10	0.10	0.12	596.46
4. Other Sectors	0.02	0.02	0.02	0.02	0.02	0.02	6.24
5. Other	0.00	0.00	0.00	0.00	0.00	0.00	13,433.66
B. Fugitive Emissions from Fuels	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	0.00
1. Solid Fuels	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	0.00
2. Oil and Natural Gas	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	0.00
2. Industrial Processes	0.06	0.05	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	-100.00
A. Mineral Products	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE	0.00
B. Chemical Industry	0.06	0.05	NE,NO	NE,NO	NE,NO	NE,NO	-100.00
C. Metal Production	NA	NA	NA	NA	NA	NA	0.00
D. Other Production							
E. Production of Halocarbons and SF ₆							
F. Consumption of Halocarbons and SF ₆							
G. Other	NA	NA	NA	NA	NA	NA	0.00
3. Solvent and Other Product Use	0.01	0.01	0.01	0.01	0.01	0.01	-45.19
4. Agriculture	0.84	0.83	0.78	0.74	0.72	0.72	-18.70
A. Enteric Fermentation							
B. Manure Management	0.09	0.09	0.08	0.08	0.08	0.08	-21.38
C. Rice Cultivation							
D. Agricultural Soils	0.76	0.75	0.69	0.66	0.64	0.64	-18.34
E. Prescribed Burning of Savannas	NA	NA	NA	NA	NA	NA	0.00
F. Field Burning of Agricultural Residues	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	0.00
G. Other	NA	NA	NA	NA	NA	NA	0.00
5. Land Use, Land-Use Change and Forestry	1.35	1.35	1.35	1.35	1.35	1.35	0.35
A. Forest Land	0.00	0.00	0.00	0.00	0.00	0.00	357.23
B. Cropland	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	0.00
C. Grassland	NE	NE	NE	NE	NE	NE	0.00
D. Wetlands	0.07	0.07	0.07	0.07	0.07	0.07	0.00
E. Settlements	NE	NE	NE	NE	NE	NE	0.00
F. Other Land	NE	NE	NE	NE	NE	NE	0.00
G. Other	1.27	1.27	1.27	1.28	1.28	1.28	0.14
6. Waste	0.02	0.02	0.02	0.02	0.02	0.02	0.47
A. Solid Waste Disposal on Land							
B. Waste-water Handling	0.02	0.02	0.02	0.02	0.02	0.02	17.09
C. Waste Incineration	0.00	0.00	0.00	0.00	0.00	0.00	-99.86
D. Other	NA	NA	NA	NA	NA	NA	0.00
7. Other (as specified in Summary 1.A)	NA	NA	NA	NA	NA	NA	0.00
Total N₂O emissions including N₂O from LULUCF	2.48	2.46	2.35	2.33	2.32	2.35	-6.72
Total N₂O emissions excluding N₂O from LULUCF	1.13	1.11	1.00	0.98	0.97	1.00	-14.84
Memo Items:							
International Bankers	0.02	0.01	0.01	0.01	0.02	0.01	29.17
Aviation	0.01	0.01	0.01	0.01	0.01	0.01	85.35
Marine	0.01	0.00	0.01	0.00	0.01	NA,NE,NO	-100.00
Multilateral Operations	NO	NO	NO	NO	NO	NO	0.00
CO₂ Emissions from Biomass							

Note: All footnotes for this table are given at the end of the table on sheet 5.

National Inventory Report 2007 - ICELAND

TABLE 10 EMISSION TRENDS
HFCs, PFCs and SF₆
 (Part 1 of 2)

Inventory 2005
 Submission 2007 v1.1
 ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	Base year (1990)	1991	1992	1993	1994	1995	1996	1997	1998	1999
	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)
Emissions of HFCs⁽³⁾ - (Gg CO₂ equivalent)	NA,NE,NO	NA,NE,NO	0,47	1,56	3,12	25,01	28,56	37,46	63,90	59,40
HFC-23	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
HFC-32	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
HFC-41	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
HFC-43-10msec	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
HFC-125	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
HFC-134	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
HFC-134a	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
HFC-152a	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
HFC-143	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
HFC-143a	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
HFC-227ea	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
HFC-236fa	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
HFC-245ca	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
Unspecified mix of listed HFCs ⁽⁴⁾ - (Gg CO ₂ equivalent)	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO
Emissions of PFCs⁽³⁾ - (Gg CO₂ equivalent)	419,63	348,34	155,28	74,86	44,57	58,84	25,15	82,36	180,13	173,21
CF ₄	0,05	0,05	0,02	0,01	0,01	0,01	0,00	0,01	0,02	0,02
C ₂ F ₆	0,01	0,01	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
C ₃ F ₈	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
C ₄ F ₁₀	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
i-C ₄ F ₈	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
C ₃ F ₁₂	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
C ₆ F ₁₄	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
Unspecified mix of listed PFCs ⁽⁴⁾ - (Gg CO ₂ equivalent)	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO
Emissions of SF₆⁽³⁾ - (Gg CO₂ equivalent)	5,38	5,38	5,38	5,38	5,38	5,38	5,38	5,38	5,38	5,38
SF ₆	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO

Note: All footnotes for this table are given at the end of the table on sheet 5.

TABLE 10 EMISSION TRENDS
HFCs, PFCs and SF₆
 (Part 2 of 2)

Inventory 2005
 Submission 2007 v1.1
 ICELAND

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	2000	2001	2002	2003	2004	2005	Change from base to latest reported year
	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)	%
Emissions of HFCs⁴³ - (Gg CO₂ equivalent)	32.28	53.78	35.16	69.35	58.40	76.74	100.00
HFC-23	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	0.00
HFC-32	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	0.00
HFC-41	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	0.00
HFC-43-10mcc	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	0.00
HFC-125	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	0.00
HFC-134	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	0.00
HFC-134a	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	0.00
HFC-152a	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	0.00
HFC-143	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	0.00
HFC-143a	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	0.00
HFC-227ea	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	0.00
HFC-236fa	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	0.00
HFC-245ca	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	0.00
Unspecified mix of listed HFCs ⁴³ - (Gg CO ₂ equivalent)	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	0.00
Emissions of PFCs⁴³ - (Gg CO₂ equivalent)	127.16	91.66	72.54	59.78	38.58	26.09	-93.78
CF ₄	0.02	0.01	0.01	0.01	0.01	0.00	-93.78
C ₂ F ₆	0.00	0.00	0.00	0.00	0.00	0.00	-93.78
C ₃ F ₈	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	0.00
C ₄ F ₁₀	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	0.00
c-C ₄ F ₁₀	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	0.00
C ₅ F ₁₂	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	0.00
C ₆ F ₁₄	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	0.00
Unspecified mix of listed PFCs ⁴³ - (Gg CO ₂ equivalent)	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	0.00
Emissions of SF₆⁴³ - (Gg CO₂ equivalent)	5.38	5.38	5.38	5.38	5.38	5.38	0.00
SF ₆	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	0.00

Note: All footnotes for this table are given at the end of the table on sheet 5.

National Inventory Report 2007 - ICELAND

**TABLE 10 EMISSION TRENDS
SUMMARY
(Part 1 of 2)**

Inventory 2005
Submission 2007 v1.1
ICELAND

GREENHOUSE GAS EMISSIONS	Base year (1990)	1991	1992	1993	1994	1995	1996	1997	1998	1999
	CO ₂ equivalent (Gg)									
CO ₂ emissions including net CO ₂ from LULUCF	3,773.27	3,678.74	3,787.24	3,877.05	3,820.81	3,831.95	3,902.00	3,975.97	3,960.75	4,135.02
CO ₂ emissions excluding net CO ₂ from LULUCF	2,150.70	2,071.82	2,199.62	2,304.35	2,266.49	2,299.13	2,388.07	2,482.63	2,493.35	2,696.28
CH ₄ emissions including CH ₄ from LULUCF	462.57	460.44	455.76	457.29	462.02	456.24	464.07	470.30	476.67	477.96
CH ₄ emissions excluding CH ₄ from LULUCF	413.43	411.30	406.62	408.15	412.88	407.10	414.93	421.16	427.53	428.82
N ₂ O emissions including N ₂ O from LULUCF	780.91	770.74	749.31	757.88	762.47	759.49	777.09	775.64	773.66	793.24
N ₂ O emissions excluding N ₂ O from LULUCF	363.09	352.60	331.49	341.12	345.94	342.38	360.29	358.41	355.91	375.85
HFCs	NA,NE,NO	NA,NE,NO	0.47	1.56	3.12	25.01	28.56	37.46	63.90	59.40
PFCs	419.63	348.34	155.28	74.86	44.57	58.84	25.15	82.36	180.13	173.21
SF ₆	5.38	5.38	5.38	5.38	5.38	5.38	5.38	5.38	5.38	5.38
Total (including LULUCF)	5,441.76	5,263.64	5,153.44	5,174.02	5,098.36	5,136.90	5,202.24	5,347.11	5,460.49	5,644.21
Total (excluding LULUCF)	3,352.23	3,189.44	3,098.85	3,135.41	3,078.37	3,137.84	3,222.37	3,387.39	3,526.20	3,738.94

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	Base year (1990)	1991	1992	1993	1994	1995	1996	1997	1998	1999
	CO ₂ equivalent (Gg)									
1. Energy	1,703.95	1,658.08	1,783.26	1,844.83	1,807.47	1,820.16	1,910.98	1,968.26	1,930.53	1,970.72
2. Industrial Processes	866.64	760.40	565.92	536.28	508.68	559.12	535.15	651.81	798.42	933.91
3. Solvent and Other Product Use	6.00	4.87	4.77	4.71	3.88	4.71	4.71	4.71	4.96	4.68
4. Agriculture	567.95	554.49	530.03	534.34	541.16	520.15	533.34	532.29	537.79	543.57
5. Land Use, Land-Use Change and Forestry ⁽⁵⁾	2,089.53	2,074.20	2,054.59	2,038.61	2,019.98	1,999.07	1,979.87	1,959.72	1,934.29	1,905.27
6. Waste	141.06	144.97	148.23	148.61	150.57	152.02	156.00	158.93	160.42	163.10
7. Other	66.63	66.63	66.63	66.63	66.63	81.68	82.18	71.38	94.08	122.96
Total (including LULUCF)⁽⁵⁾	5,441.76	5,263.64	5,153.44	5,174.02	5,098.36	5,136.90	5,202.24	5,347.11	5,460.49	5,644.21

⁽¹⁾ The column "Base year" should be filled in only by those Parties with economies in transition that use a base year different from 1990 in accordance with the relevant decisions of the COP. For these Parties, this different base year is used to calculate the percentage change in the final column of this table.

⁽²⁾ Fill in net emissions/removals as reported in table Summary 1.A. For the purposes of reporting, the signs for removals are always negative (-) and for emissions positive (+).

⁽³⁾ Enter actual emissions estimates. If only potential emissions estimates are available, these should be reported in this table and an indication for this be provided in the documentation box. Only in these rows are the emissions expressed as CO₂ equivalent emissions.

⁽⁴⁾ In accordance with the UNFCCC reporting guidelines, HFC and PFC emissions should be reported for each relevant chemical. However, if it is not possible to report values for each chemical (i.e. mixtures, confidential data, lack of disaggregation), this row could be used for reporting aggregate figures for HFCs and PFCs, respectively. Note that the unit used for this row is Gg of CO₂ equivalent and that appropriate notation keys should be entered in the cells for the individual chemicals.

⁽⁵⁾ Includes net CO₂, CH₄ and N₂O from LULUCF.

**TABLE 10 EMISSION TRENDS
SUMMARY
(Part 2 of 2)**

Inventory 2005
Submission 2007 v1.1
ICELAND

GREENHOUSE GAS EMISSIONS	2000	2001	2002	2003	2004	2005	Change from base to latest reported year
	CO ₂ equivalent (Gg)	(%)					
CO ₂ emissions including net CO ₂ from LULUCF	4.158,84	4.142,87	4.217,93	4.115,67	4.178,48	4.158,24	10,20
CO ₂ emissions excluding net CO ₂ from LULUCF	2.745,09	2.746,56	2.842,41	2.764,70	2.862,68	2.872,13	33,54
CH ₄ emissions including CH ₄ from LULUCF	471,74	477,91	466,28	465,44	460,25	465,07	0,54
CH ₄ emissions excluding CH ₄ from LULUCF	422,60	428,77	417,14	416,30	411,11	415,93	0,60
N ₂ O emissions including N ₂ O from LULUCF	768,63	762,13	728,49	720,93	720,66	728,47	-6,72
N ₂ O emissions excluding N ₂ O from LULUCF	351,64	345,08	311,33	302,45	301,80	309,20	-14,84
HFCs	32,28	53,78	35,16	69,35	58,40	76,74	100,00
PFCs	127,16	91,66	72,54	59,78	38,58	26,09	-93,78
SF ₆	5,38	5,38	5,38	5,38	5,38	5,38	0,00
Total (including LULUCF)	5.564,03	5.533,72	5.525,78	5.436,54	5.461,75	5.459,98	0,33
Total (excluding LULUCF)	3.684,15	3.671,23	3.683,96	3.617,96	3.677,94	3.705,47	10,54

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	2000	2001	2002	2003	2004	2005	Change from base to latest reported year
	CO ₂ equivalent (Gg)	(%)					
1. Energy	1.872,62	1.844,66	1.916,81	1.861,99	1.961,45	1.992,22	16,92
2. Industrial Processes	949,96	971,41	936,31	959,77	949,33	944,30	8,96
3. Solvent and Other Product Use	4,53	4,03	4,03	3,72	3,41	3,29	-45,19
4. Agriculture	525,48	521,94	498,70	483,10	472,42	475,21	-16,33
5. Land Use, Land-Use Change and Forestry ⁽⁵⁾	1.879,88	1.862,49	1.841,81	1.818,59	1.783,81	1.754,51	-16,03
6. Waste	168,09	174,71	168,77	171,48	167,25	167,06	18,43
7. Other	163,48	154,48	159,35	137,89	124,08	123,38	85,17
Total (including LULUCF)⁽⁵⁾	5.564,03	5.533,72	5.525,78	5.436,54	5.461,75	5.459,98	0,33

⁽¹⁾ The column "Base year" should be filled in only by those Parties with economies in transition that use a base year different from 1990 in accordance with the relevant decisions of the COP. For these Parties, this different base year is used to calculate the percentage change in the final column of this table.

⁽²⁾ Fill in net emissions/removals as reported in table Summary I.A. For the purposes of reporting, the signs for removals are always negative (-) and for emissions positive (+).

⁽³⁾ Enter actual emissions estimates. If only potential emissions estimates are available, these should be reported in this table and an indication for this be provided in the documentation box. Only in these rows are the emissions expressed as CO₂ equivalent emissions.

⁽⁴⁾ In accordance with the UNFCCC reporting guidelines, HFC and PFC emissions should be reported for each relevant chemical. However, if it is not possible to report values for each chemical (i.e. mixtures, confidential data, lack of disaggregation), this row could be used for reporting aggregate figures for HFCs and PFCs, respectively. Note that the unit used for this row is Gg of CO₂ equivalent and that appropriate notation keys should be entered in the cells for the individual chemicals.

⁽⁵⁾ Includes net CO₂, CH₄ and N₂O from LULUCF.

Documentation box:

- Parties should provide detailed explanations on emissions trends in Chapter 2: Trends in Greenhouse Gas Emissions and, as appropriate, in the corresponding Chapters 3 - 9 of the NIR. Use this documentation box to provide references to relevant sections of the NIR if any additional information and further details are needed to understand the content of this table.
- Use the documentation box to provide explanations if potential emissions are reported.